Modeled structure of the 75-kDa neurotrophin receptor

BARBARA S. CHAPMAN AND IRWIN D. KUNTZ

University of California, San Francisco, California 94143-0446

(RECEIVED April 11, 1995; ACCEPTED June 15, 1995)

Abstract

Motifs in ligand-binding domains of the neurotrophin (NTR) and lymphotoxin (TNFR-I) receptors define a family of receptors that mediates programmed cell death. We have explored relationships of architecture and function in this family through a molecular model of NTR, also called p75^{NGFR} or LANR. Modeling by homology took advantage of four modular subdomains in the crystal structure of TNFR-I that also occur in NTR. Hypothetical complexes between the model and a ligand structure (for nerve growth factor, NGF) were then examined using docking software. NTR appears to bind in the dimer interface of NGF, making two sets of contacts. NTR subdomains III and IV provide the ligand-contact surfaces, in contrast to TNFR, in which subdomains II and III contact TNF- β . NTR subdomain II appears to have been evolutionarily modified, potentially contributing to an interface between receptor subunits. These and other specific predictions of the model will require experimental confirmation.

Keywords: homology modeling; ligand binding; neurotrophin receptor; NGF; NGFR/TNFR family; protein-protein docking

Growth factors and cytokines act in concert to regulate cell survival or programmed cell death, with multiple factors influencing each cell type at different stages of differentiation (Korsching, 1993; Raff et al., 1993). The network provides cues during development and maintains tissue homeostasis. Much recent work has been directed toward unraveling the complex signaling that maintains it. During evolution of this signaling system, a small number of receptor families has come to bind diverse ligands, a situation not unexpected in elaboration of closely coordinated functions. The history of these receptor families and their ligands is likely to be found imbedded in amino acid sequence and three-dimensional structure (Bazan, 1991). Insights into this coordinated system may be gained by comparing related receptors that bind different ligands and by evaluating the functional implications of structural adaptation.

One of these families is defined by the neurotrophin (NT) and lymphotoxin (TNF) receptors (Smith et al., 1990). Ten members and their ligands have been discovered (Beutler & Van Huffel, 1994), although specific functions are known for only a few. Crystal structures have been published for three ligands, TNF- α (cachectin) (Eck & Sprang, 1989), TNF- β (lymphotoxin- α) (Eck et al., 1992), and NGF (nerve growth factor) (McDonald et al., 1991). One ligand-receptor complex, TNF- β with TNFR-I, has

also been solved (Banner et al., 1993). Although the NTRs and TNFRs are structurally related, their ligands come from two different families (Daopin et al., 1992; McDonald & Hendrickson, 1993).

The 75-kDa neurotrophin receptor (NTR), also called p75^{NGFR} or LANR (low-affinity neurotrophin receptor), mediates programmed cell death in neurons deprived of neurotrophic factors (Lee et al., 1992, 1994; Rabizadeh et al., 1993; Barrett & Bartlett, 1994). It binds all four known neurotrophins, which are about 50% identical in amino acid sequence and similar in structure (Rodríguez-Tébar et al., 1992; Davies et al., 1993; Narhi et al., 1993). The neurotrophin receptor has recently been shown to associate with tyrosine kinase receptors trkA, trkB, and trkC to form high-affinity NGF-binding complexes (Hantzopoulos et al., 1994; Mahadeo et al., 1994; Wolf et al., 1995). Modeling NTR provides an unusual opportunity to explore a versatile architecture capable of association with receptor subunits from separate superfamilies and able to bind diverse ligands.

Advances in computational techniques, molecular biology, and structural analysis have made it feasible to construct useful models of macromolecular complexes (Ring & Cohen, 1993; Johnson et al., 1994). These models allow design of additional biochemical and mutagenesis studies to unravel structure/function relationships. For example, molecular models developed from crystallographic and biochemical data have been useful in analysis of protein-protein complexes such as ubiquinol:cytochrome c2 oxidoreductase (bc-complex) (Crofts et al., 1992) and

Reprint requests to: Barbara S. Chapman, Department of Pharmaceutical Chemistry, Box 0446, University of California, San Francisco, California 94143-0446; e-mail: bchapman@cgl.ucsf.edu.

the actin-myosin complex (Rayment et al., 1993). A model of the cytokine interleukin-6 (IL-6), prepared from the structurally related bovine granulocyte colony-stimulating factor, has been used to design antagonists for the IL-6 receptor (Savino et al., 1994). Our laboratory has used a docking procedure to regenerate molecular complexes from separately crystallized components (Shoichet & Kuntz, 1991; Shoichet et al., 1992).

In this paper we employ the crystal structure of the 55-kDa TNF receptor (Banner et al., 1993) to prepare a molecular model for the neurotrophin receptor. Modeling by homology was facilitated by the modular architecture of these receptors, whose extracellular domains are each composed of four folded subdomains (Johnson et al., 1986; Banner et al., 1993). We have investigated molecular interactions between the neurotrophin receptor model and the crystal structure of NGF (McDonald et al., 1991) using DOCK3.5 drug design software (Shoichet & Kuntz, 1991; Meng et al., 1992, 1993). The crystal structure of a TNF- β trimer in complex with its receptor revealed that TNFR subunits bind in the interface between ligand subunits (Banner et al., 1993). An analogous receptor-binding site on NGF was evaluated using data from mutagenesis and biochemical modification (Rosenberg et al., 1986; Ibáñez et al., 1990, 1991, 1992, 1993; Drinkwater et al., 1991, 1993). Several potential binding surfaces were explored on the receptor model by docking into the crystal structure of NGF (Shoichet & Kuntz, 1991; Meng et al., 1992). Likely contact residues were found in subdomains III and IV of NTR, as predicted by mutagenesis and antibody-binding studies. The model complex also revealed that subdomain II of NTR is probably not involved in ligand contact but appears to be specialized for a new function.

Results

Shared patterns and motifs in TNFR and NTR

Primary sequences of the extracellular domains from receptors in this family can be aligned using groups of 6 cysteine residues within repeating sequences of approximately 40 residues (Smith et al., 1990). The 55-kDa lymphotoxin receptor (TNFR) and 75-kDa neurotrophin receptor (NTR) each contain four of these repeating units, shown in Figure 1A (Johnson et al., 1986; Loetscher et al., 1990). The first crystal structure for a member of the NTR/TNFR family revealed that each repeat folds into a compact structure having three disulfide bonds, two hydrogen bonds, and one packed aromatic side chain (Banner et al., 1993). These are the structurally important residues. Informed by the three-dimensional structure, the alignment shows that, despite an overall sequence identity of 29%, more than 80% of structurally important residues are conserved (Fig. 1A). These structural residues form a scaffolding for three loops in each folded unit (Banner et al., 1993). Nearly all of the sequence variability resides in these loops, labeled *a*, *b*, and *c* in Figure 1A. Distribution of substitutions into loop regions of closely related receptors has been observed, for example, in the prolactin and growth hormone receptors (Somers et al., 1994).

Modeling strategies

The NTR model was created by dividing one of the TNFR crystal structure subunits into its four subdomains, retaining the ladder-like scaffold composed of three disulfide bridges and

their connecting residues, replacing each variable loop, and re-assembling the four-unit structure. Three problems were addressed. First, a good algorithm was needed to perform the loop replacements. The most conservative approach required aligning residues of each loop so that a single gap or insertion was placed at the maximum distance from structural elements. These structural elements were then used as anchors for the loop replacement function in SYBYL (Tripos Associates, St. Louis, Missouri). Given a range of residues on the amino and carboxyl sides of the gap or insertion, the program searches a structural database for coordinates of main-chain atoms that match at anchoring α carbons. The algorithm produces a set of possible main chains that can be evaluated for fit to anchoring residues and for homology with the desired loop sequence.

The second problem was to model the fourth subdomain of NTR, given that its homologue in the crystal structure was incomplete. The second TNFR subdomain was chosen as the template for NTR subdomain IV on the basis of sequence homology. TNFR subdomain II and NTR subdomain IV align with a single gap in loop *b* and 30% sequence identity (Fig. 1B). The next best alignment was with TNFR subdomain III, showing two gaps and 16% sequence identity.

The third problem was to resolve steric conflicts created by substitutions of side chains and loop backbones. Conflicts were resolved within SYBYL by a two-step procedure. Working in short segments from more constrained to less constrained regions, alternative rotamers were tested at each position until most van der Waals contacts were eliminated. Then a molecular mechanics procedure was applied to shift atom positions and bond torsion angles, leading to a relaxed, low-energy structure (Weiner et al., 1986).

Building folded units of NTR

The four subdomains of the neurotrophin receptor were modeled and energy minimized following the strategies outlined above. Properties of the crystal template and model subdomains are compared in Table 1. Homologous folded units were similar in subdomain length and location of structurally conserved residues (evaluated as distances between matched α carbons). Total energies after minimization were generally higher for the model than for the crystal subdomains, due to higher torsional and bond-stretching energy, less favorable van der Waals interactions, and fewer electrostatic combinations. The modeled subdomains of NTR are shown in Figure 1C.

As suggested by the sequence alignment, subdomain I of the NTR conforms to the N-terminal repeat motif that characterizes members of the receptor family (Smith et al., 1990). There are no spacer residues between the second and third cysteines, leading to a shortened cylindrical shape relative to the other three subdomains. The NTR subdomain I is shorter by five residues than its TNFR homolog and differs by substitution of both hydrogen bond partners (Lys replaces Thr in the scaffold and Gln replaces Asp in loop *c*, shown in the upper left of Fig. 1C). These changes result in an RMS deviation (RMSD) of 0.77 Å/atom at structurally conserved α carbons and in a higher internal energy (Table 1).

Sequence changes in subdomain II include four hydrophobic side chains for charged side chains of TNFR. These substitutions dramatically alter both the hydrophobicity and electrostatic character of subdomain II. The net effect of these replacements

Fig. 1. Sequence and structure of folded subdomains of TNFR and NTR. **A:** Amino acid sequences of TNFR-I and NTR aligned using structurally important residues. Disulfide bridges are indicated by brackets at the top. Structurally important residues are labeled at the top of each column with the symbol \$. Differences between human and rat NTR are marked with asterisks. **B:** TNFR subdomain II aligned with NTR subdomain IV. **C:** Tube and stick representation of four modeled subdomains of the 75-kDa NTR. Main-chain atoms are shown as tubes and structurally important side chains are displayed as sticks. Each folded structure has three disulfide bridges, one packed aromatic ring, and two side chains for potential hydrogen bond partners. The three loops of each fold are labeled *a*, *b*, and *c*. Upper left, subdomain I; lower left, subdomain II; upper right, subdomain III; lower right, subdomain IV. **D:** Space-filling representation of subdomain II of NTR (right) compared with subdomain II of TNFR (left). Color code: light gray, hydrophobic side chains; dark gray, polar or charged side chains; black, packed aromatic ring. Side chains of substituted hydrophobic for polar residues are labeled.

is to convert the net charge from +1 in TNFR to -5 in NTR and increase the number of hydrophobic side chains from 9 in TNFR to 14 in NTR. Insertion of Leu at the beginning of loop *a* led to a new trajectory for the loop and to packing of the Phe 45 aromatic ring on the outside, rather than the inside of the folded structure (Fig. 1C, lower left). Analogous insertions occur in subdomain II of the human Fas and CD27 receptors (Camerini et al., 1991; Itoh et al., 1991). Displacement of Phe 45 relative to Phe 60 of TNFR produced an RMSD of 2.25 Å/atom for α carbons of the nine structurally important residues of the subdomain (Fig. 1A). However, RMSD fell to 0.73 Å/atom when Phe 45 and Phe 60 were excluded from the calculation (Table 1). These modifications also introduced hydrophobic patches onto the surface of this subdomain, shown in Figure 1D.

Subdomain III in TNFR is characterized by a number of aromatic side chains in loop *a* (Fig. 1A; Fig. 1C, upper right).

The neurotrophin receptor sequence preserves three of the five aromatic residues in the amino-terminal half of its shortened *a* loop. Loop *c* of the subdomain is highly conserved, where the two sequences share six of seven residues. Substitution of Arg for the hydrogen bonding Ser residue in the scaffold slightly changed internal packing, but the modeled structure remained very close in height (28 Å), RMS distance (0.41 Å/atom), and energy (Table 1).

NTR subdomain IV was based on structural information from subdomain II of TNFR in order to extend the subdomain beyond loop *a*. The modeled structure (Fig. 1C, lower right) displays the canonical fold, including hydrogen bonding of the scaffold and packed aromatic ring. The minimized structure relaxed to an energy of -125.2 kcal/mol and gave an RMSD of 0.32 Å/atom from the template at structurally conserved α carbons (Table 1).

Table 1. Neurotrophin receptor model subdomains compared with TNFR structure

Measurement	Subdomain I	Subdomain II	Subdomain III	Subdomain IV
Domain length (Å)				
TNFR	23.1	30.6	28.0	Incomplete
NTR	23.4	28.2	28.0	27.9
RMSD C α ^a (Å/atom)				
NTR versus TNFR	0.77	2.25 ^b	0.41	0.32 ^c
Linker ϕ , ψ				
TNFR	−122°, 159° −72°, 134°	−165°, −153°	−114°, 119°	
NTR	−143°, 147° −70°, 132°	−64°, 172°	−98°, 105° −62°, 145°	
Energy ^d (kcal/mol)				
TNFR	−89.2	−119.8	−129.6	Incomplete
NTR	−65.9	−54.9	−108.9	−125.2

^a Pairwise distances between α carbons of the model matched with α carbons of the TNFR-I crystal structure were calculated for the six Cys residues, the structural aromatic residue in loop *a*, and the two hydrogen bonding partners.

^b Matching eight residues (not including Phe 60 and Phe 45), RMS error = 0.73 Å/atom.

^c Compared with domain II, which was used as template.

^d After minimizing individual domains using the AMBER all-atom force field.

Assembly and refinement of the NTR model

The full NTR structure was assembled by linking its component subdomains and setting ϕ and ψ angles of the linking residues. Angles from the TNFR crystal structure were used except at the junction of subdomains II and III. Here Arg substitutes for Gly in a forbidden conformation (Ramachandran & Sassiakharan, 1968). ϕ and ψ angles were adjusted interactively in SYBYL by computing the overall energy at various settings and selecting angles producing the fewest van der Waals conflicts (lowest total energy). The structure was refined through local energy minimization to resolve van der Waals conflicts in the subdomain interface zones, and the entire molecule was then minimized. We obtained a minimum energy of −460.6 kcal/mol, compared with −495.6 kcal/mol for the minimized TNFR molecule. The two structures are shown in Figure 2A.

Main-chain torsion angles of the refined structure were analyzed in a Ramachandran plot (Ramachandran & Sassiakharan, 1968) and compared with angles in the TNFR crystal structure (Fig. 2B). Fewer residues of the NTR model were in ideal conformations compared with the crystal structure. Eleven of these strained bonds were concentrated in NTR subdomain II loop *a* (five), loop *b* (three), and loop *c* (three). To see if the structure could be improved, residues having nonideal torsions were excised and replaced using the loop search module of SYBYL. The strategy either did not lead to significant change or resulted in moving the problem to another bond.

To explore the possibility that the structure was trapped in a local energy minimum, we subjected the model to a molecular dynamics procedure. The model was warmed to 300 K for 10 ps, then energy minimized. Independent dynamic runs produced models having total energies of −579.0 and −612.6 kcal/mol after minimization. Both of these structures were lower in energy than the minimized crystal structure but showed significant disruptions in folding within subdomains (Table 2). RMS differences at structurally important α carbons increased from an average of 0.94 Å/atom for the original model to more than 2 Å/atom for the quenched models.

Much of the observed strain and high internal energy resulted from limitations in the ability of current software to accurately predict loop structure (Ring & Cohen, 1993). This problem is most apparent in the modeled structure of subdomain II, owing to the low sequence homology in its loops. The version of the model that was minimized without molecular dynamics was used for the docking experiments described below.

Analysis of receptor–ligand interactions by molecular docking

One measure of the model's validity is how realistically it interacts with one of its ligands. We used the program DOCK3.5 to explore the geometry of the complex between our model of NTR and NGF (McDonald et al., 1991). DOCK drug design software was developed to identify molecules having shapes complemen-

Table 2. Relationship of scaffold atoms in the NTR model to the TNFR-I crystal structure before and after molecular dynamics simulation

Subdomain	RMS difference versus TNFR (Å/atom) ^a		
	NTR model initial	10 ps at 300 K #1	10 ps at 300 K #2
I	0.77	2.46	2.15
II	2.25	3.46	2.90
III	0.41	1.81	1.42
IV ^b	0.32	1.41	1.85
All	0.94	2.29	2.08

^a RMS errors calculated for distances between α carbons of nine structurally important residues in each subdomain, including six Cys residues, two hydrogen-bonding partners, and one packed aromatic.

^b Because subdomain IV of TNFR is incomplete in the crystal structure, distances were calculated versus subdomain II of TNFR.

A

Fig. 2. Assembled structure of NTR model. **A:** Space-filling representations of TNFR R subunit (left) and the NTR model (right). Subdomains are color coded: red, subdomain I; yellow, subdomain II; green, subdomain III; blue, subdomain IV; orange, residues linking subdomains. **B:** Ramachandran plots of ϕ and ψ angles. TNFR R subunit structure (left); NTR model (right).

B

tary to surface features of proteins (Kuntz et al., 1982). An important application of this software has been to predict binding modes of small molecules and polypeptide ligands (Shoichet & Kuntz, 1991; Meng et al., 1992, 1993).

Having coordinates for the TNFR structure in complex with its ligand TNF- β provided a way to test the docking method. By regenerating the TNF/TNFR crystal complex from its separated components, a procedure and set of parameters were established for evaluating the NTR model complex with NGF.

The TNF/TNFR crystal complex is composed of a TNF- β trimer with three TNFR subunits bound at the interfaces between

ligand subunits (Banner et al., 1993). A test system was prepared from the A and C subunits of TNF- β and subunit R of the TNF receptor, which contacts their interface. A representation of the solvent-accessible surface was prepared using the program DMS (Connolly, 1983). Matching points for the docking site were generated with SPHGEN (Kuntz et al., 1982) and selected for proximity to crystallographic contact residues using molecular graphics tools in MIDAS (Ferrin et al., 1988). Similarly, a set of matching points close to crystallographically identified contacts was chosen for the TNFR R subunit. Scoring grids for the docking site were prepared with CHEMGRID (Meng et al.,

1992) to provide force-field scores for orientations of the TNFR subunit found by DOCK in the docking site. Parameters were adjusted through a series of docking runs that produced force-field scores within 10% of the crystal complex, and that oriented the TNFR molecule within an RMS distance of 1.3 Å/atom from its crystallographic position (data not shown). These parameters were used for the NTR-NGF study.

A likely binding site for NTR was identified on NGF using a list of residues generated from binding analyses of NGF derivatives and mutants (Rosenberg et al., 1986; Ibáñez et al., 1990, 1992; Drinkwater et al., 1991, 1993). Known contact residues were then distributed along a dimer interface of the NGF crystal structure on the assumption that receptors of a family might have similar modes of ligand binding. This assumption was consistent with biological evidence that heterodimers formed by protomers from different neurotrophins show 10-fold lower activity than homodimers in neuronal survival assays (Jungbluth et al., 1994).

Biochemically important residues presented two clear binding surfaces when arranged along the groove formed by NGF protomers (Fig. 3A). This molecular surface was converted to a docking site, and force-field scoring grids were computed as for the TNF/TNFR test complex. Three structural hypotheses were tested for an NTR-NGF complex: Hypothesis 1, contacts formed by acidic residues of subdomains III and IV on the "outside" face of the NTR model (Fig. 2A); hypothesis 2, contacts from NTR residues in homologous positions to TNFR ligand contacts; and hypothesis 3, contacts near residues of NTR subdomains III and IV on the "inside" of the angle formed by the subdomains of NTR.

These hypotheses were tested by forming complexes with DOCK3.5 (Meng et al., 1993). Shown in Figure 3B are force-field scores plotted against RMS distance of each orientation relative to the arbitrary input orientation. Hypothesis 1, acidic contacts of subdomains III and IV, produced 303 orientations in the docking site (Fig. 3B, top panel). When viewed with MIDAS, more than 80% of these appeared to touch both clusters of residues in the dimer interface of NGF. The principal family (indicated by the arrow) among these orientations produced the highest observed force-field score (−42.8). This score was composed of an electrostatic score of −17.0, a van der Waals repulsion component of 24.5, and a van der Waals attraction score of −50.3.

Hypothesis 2, the contact surface homologous to the TNFR binding site, yielded 138 possible orientations. Viewed with MIDAS, approximately 40% of these orientations interacted with both binding-residue clusters on NGF. The best score was relatively poor (−35.0) and the principal family of orientations did not include the best score (Fig. 3B, middle panel).

For hypothesis 3, contacts representing the inside surface of subdomains III and IV, DOCK found 193 orientations, none of which was simultaneously in contact with both zones in the NGF binding site (Fig. 3B, bottom panel). The results of testing these three models showed that the largest number of orientations, the highest force-field score, and the best alignment in the docking site were obtained using hypothesis 1, contacts in the acidic outer surface of subdomains III and IV.

Several of the highest-scoring orientations were viewed in complex with NGF to evaluate the number of biochemically identified contacts made by the docked receptor model. Potential binding interactions were defined as atoms of NGF posi-

tioned within a radius of 3.5 Å from atoms of the receptor. The best orientation found using the TNFR-homologous surface appeared to be close to NGF side chains not identified in mutagenesis studies. In contrast, the major family of acidic-surface orientations had side chains close to the presumptive binding residues of NGF. One of these orientations is shown in complex with NGF in Figure 3C.

NGF side chains within a contact radius of 3.5 Å from the NTR model included Lys 32, Lys 34, Lys 95, and Arg 100, which contribute strongly to NTR binding (Ibáñez et al., 1992). The C-terminal end of NTR subdomain IV contacted the NGF surface formed by Ile 31, Lys 32, Lys 34, Lys 88, and Arg 100. The N-terminal loop of subdomain III appeared to interact with residues near the N- and C-termini of NGF, including Gly 10, Ser 113, and Arg 114 (Drinkwater et al., 1993). In all, more than half of the NGF residues at the modeled NTR/NGF interface have been implicated by biochemistry and mutagenesis (Rosenberg et al., 1986; Ibáñez et al., 1990, 1992; Drinkwater et al., 1991, 1993).

Ligand contacts of the receptor included Thr 91, Arg 93, Glu 95, Arg 150, Glu 151, Cys 152, and Thr 153. These residues are found on the third and fourth subdomains of NTR, consistent with deletion studies and the location of ligand-blocking monoclonal antibody epitopes (Welcher et al., 1991; Yan & Chao, 1991; Baldwin et al., 1992; Baldwin & Shooter, 1994). NTRs from three species have been sequenced (Johnson et al., 1986; Radeke et al., 1987; Large et al., 1989; Heuer et al., 1990). Predicted binding site residues Thr 91, Glu 95, Arg 150, Glu 151, and Cys 152 are either identical or have very conservative replacements in human, rat, and chicken receptors. Arg 93 is changed to Ser and Thr 153 to Lys in chicken NTR. Species specificity of all known monoclonal antibodies recognizing the NTR binding site suggests that precise conservation of binding site residues may not be critical.

Species specificities of two NGF-blocking monoclonal antibodies, ME20.4 for the human receptor (Marano et al., 1987) and MC192 for the rat receptor (Chandler et al., 1984), permit identification of probable epitopes in the binding site by comparing sequences of the human and rat receptors (Johnson et al., 1986; Radeke et al., 1987). Six amino acid differences found in NTR subdomains III and IV are marked in Figure 1A. Figure 4 illustrates the overlap between the probable residues of the antibody epitopes and the contact residues suggested by DOCK3.5.

Discussion

Insights from molecular modeling and docking

The amount of detailed information that can be extracted from our modeling effort depends on the robustness of the model and the accuracy of the docking simulation. Validity of the NTR model can be judged both by its consistency with experimental evidence and its ability to interact with the NGF structure in a plausible way. We have shown that NTR subdomains III and IV interact with NGF in the model complex, consistent with data derived from receptor mutagenesis (Sehgal et al., 1989; Welcher et al., 1991; Yan & Chao, 1991; Baldwin et al., 1992; Barker & Murphy, 1992; Baldwin & Shooter, 1994, 1995). Having placed biochemical and mutagenesis data for NGF in structural context by analogy with the receptor binding surface of TNF- β , we have shown that the model of NTR fits into the presumptive

Fig. 3. Docking the NTR model structure with its ligand, NGF. **A:** Space-filling representation of murine NGF dimer. Protomers are colored light and dark green, with disulfide bridges in yellow. Side chains of residues identified by mutagenesis to affect interactions between NGF and NTR are colored blue. These residues are Phe 12, Ile 31, Lys 32, Lys 34, Arg 50, Tyr 52, Arg 69, Lys 74, Lys 88, Lys 95, Arg 100, and 112–115. **B:** Orientations of NTR in the NGF docking site identified by DOCK3.5. The force-field score (FF score) for each orientation is plotted against RMS distance from an arbitrary starting orientation of NTR. Top, surface including acidic residues from the "outside" of the angle formed by subdomains II and III; middle, surface corresponding to TNFR ligand contacts; bottom, surface from "inside" the angle formed by subdomains II and III. NTR structural homologues of TNFR contact residues are Val 48, Ala 51, Thr 52, Pro 54, Lys 56, Pro 57, Glu 60, Val 62, Gly 63, Leu 64, Gln 87, Asp 88, and Arg 93. **C:** Views of the NGF dimer docked with the modeled NTR in a space-filling representation. Color coding of NTR domains is the same as in Figure 2, except that linker residues are not separately colored.

Fig. 4. Ligand contacts between NTR and NGF predicted by docking. Space-filling model of NTR. Residues in contact (i.e., within a radius of 3.5 Å) with biochemically identified residues of NGF are colored pink. Residues potentially forming epitopes for species-specific monoclonal antibodies are colored magenta. These include Thr 90 (Glu in rat), Arg 93 (His), Arg 98 (Ser), Ala 102 (Val), Asp 123 (Glu), and Arg 154 (Pro). The Arg 93 side chain, identified in both docking and epitope analysis, is shown in rose.

binding site on NGF (Rosenberg et al., 1986; Ibáñez et al., 1990, 1991, 1992, 1993; Drinkwater et al., 1991, 1993; Banner et al., 1993). This complex is formed in a stereospecific way, with docking scores that are more than 75% of the regenerated TNF/TNFR crystal complex. It makes contact with more than 50% of experimentally defined receptor binding side chains on NGF. This docking mode leaves open the possibility of two NTR molecules binding a single NGF dimer, as is occasionally observed experimentally.

Limitations of the docking method and the model contribute to low force-field scores and tentative interactions with NGF. Several large approximations affect the quality of the result, including the inability of DOCK3.5 to emphasize hydrogen bonds. Based on tests with the TNF/TNFR crystal complex, the NTR/NGF model complex is likely to be at least 1.3 Å/atom (and possibly much further) away from the true relationship between receptor and ligand. It is clear that receptor/ligand contacts are predominantly with one subunit of NGF, and some probable contacts on the second NGF subunit are unaccounted for in the docked complex. Further refinement of the model, with particular attention to subdomain II, should be done to take advantage of advances in methodology. Given the strengths and uncertainties inherent in this model, a cautious analysis is warranted.

Versatility of modular architecture and interactions between structural units

The first description of the structural fold characteristic of the NTR/TNFR family anticipated that relatives could be modeled by homology (Banner et al., 1993). The secondary structure of these modules is composed entirely of loops connected by disulfide bridges. The cylindrical unit is stabilized by one major hydrophobic interaction and a pair of hydrogen bonds. Although average amino acid identity is only 29% in a structure-based alignment, identities rise to 81% at structurally important residues (Fig. 1A). Thus, the scaffold has been conserved, although loop sequences have undergone extensive evolution from an ancestral subdomain. Loops probably contain most of the structural information for specific ligand binding. Comparison of NTR and TNFR is particularly informative in this regard because their ligands are structurally unrelated (Daopin et al., 1992; McDonald & Hendrickson, 1993). A docking analysis, presented here, has confirmed biochemical data suggesting that loops of units III and IV of NTR participate in ligand binding. This contrasts with the TNF receptor, which binds TNF- β through contacts on subdomains II and III (Banner et al., 1993).

Although the subdomains fold independently, significant interactions at the interfaces of adjacent units are clearly predicted by the model, which reveals tightly stacked cylinders (Fig. 2A). Loops forming the lower and upper edges of adjacent subdomains form contacting surfaces in many places, limiting the angle of bend to a few degrees in the NTR model as in the TNFR-I crystal structure (Banner et al., 1993). Direct interactions between nonadjacent subdomains do not appear possible.

Effects of one subdomain on the structure of the next have been observed experimentally. Mutations in the first two subdomains can reduce or eliminate ligand crosslinking with carbodiimide (Yan & Chao, 1991; Baldwin et al., 1992; Baldwin & Shooter, 1994, 1995), whereas directly measured NGF binding affinity appears normal (Sehgal et al., 1989; Welcher et al., 1991; Baldwin & Shooter, 1995). In addition, most of these mutants are recognized by ligand-blocking monoclonal antibodies, indicating that the structure of the binding site is not damaged (Yan & Chao, 1991; Baldwin et al., 1992; Baldwin & Shooter, 1994, 1995). The crosslinking chemistry probably detects small structural perturbations propagated from the mutation site through stacked loops. As predicted, complete deletion of subdomain I (Sehgal et al., 1989; Welcher et al., 1991; Baldwin et al., 1992) has less effect than mutations within it (Yan & Chao, 1991; Baldwin & Shooter, 1994, 1995). Five of 14 mutations in subdomain II interfere with crosslinking, but none has been shown to prevent ligand binding. Carboxylate groups targeted by the carbodiimide (in crosslinking to NGF) are probably located in the second subdomain. Mutations in this unit could thus reduce crosslinking without inhibiting ligand or monoclonal antibody binding.

Another adaptation of modular architecture exploits inherent flexibility in relationships between subdomains in order to bind different ligands. The angles formed between subdomains are altered in the NTR model relative to the TNFR crystal structure, leading to a different overall bend (Fig. 2A). An analogous rotation and translation of domains allows human prolactin and growth hormone receptors to preferentially bind different ligands, despite having nearly superimposable (RMS distance =

0.85 Å/atom) domain structures (Somers et al., 1994). Likewise, modeled subdomains of NTR match their template TNFR crystal structures to an average RMS difference of 0.94 Å/atom at α carbons of structurally conserved residues. Crystal complexes of growth hormone receptor and prolactin receptor with their common ligand, growth hormone, also provide a model for understanding how NTR might interact with its other NT ligands (Somers et al., 1994). These structurally related hormone receptors bind the same ligand in distinct modes through overlapping sets of side chains; binding and specificity are controlled by general properties of the surfaces. Experimental evidence indicates that NTR ligands brain-derived neurotrophic factor (BDNF) and NT-3 likewise have different binding modes: each NT interacts with distinct on-rate and off-rate components of its equilibrium binding constant (Rodríguez-Tébar et al., 1992). Interactions of the NTR model with its other ligands should be explored when their crystal structures become available.

Modular architecture may also provide opportunities for interactions between components of a receptor complex. NTR undergoes a change in affinity from a K_d of 3 nM to approximately 30 pM upon association with members of the trk tyrosine kinase family of NTRs, presumably by formation of a physical complex (Hempstead et al., 1991; Battleman et al., 1993; Benedetti et al., 1993; Barker & Shooter, 1994; Hantzopoulos et al., 1994; Mahadeo et al., 1994; Wolf et al., 1995). The repeat unit II of NTR displays features that would indicate a functional role in intersubunit contacts. Unlike its homologue in TNFR, the modeled subdomain II of NTR does not participate in ligand binding, but bends away from ligand-binding modules III and IV. The second NTR repeat displays several structural modifications, which involve repositioning the aromatic side chain in loop α , accumulation of six negative charges, and introduction of five additional hydrophobic residues (Fig. 1D). Patches of aromatic and aliphatic side chains created by these substitutions could stabilize a protein-protein interface and could contribute to a high-affinity binding site (Clackson & Wells, 1995).

Although not accepted by all investigators, NTR interacts with gp140trk on both sides of the cell membrane to form the high-affinity binding complex (Hempstead et al., 1990; Battleman et al., 1993; Benedetti et al., 1993; Mahadeo et al., 1994; Wolf et al., 1995). Intracellular domain contacts appear to be essential for bringing the two different subunits into the complex, whereas their extracellular domains establish a high-affinity binding site (Wolf et al., 1995). Our model predicts that the extracellular NTR-gp140trk contact zone will include NTR subdomain II. In proposing subdomain II as part of the high-affinity binding site, we account for experiments showing that NGF is more readily crosslinked to NTR in its low-affinity than in its high-affinity state (Hosang & Shooter, 1985; Meakin & Shooter, 1991) and that monoclonal antibody MC192 has opposite effects on low- and high-affinity NGF binding (Chandler et al., 1984).

Implications for receptor stoichiometry and function

Two important questions about the NTR concern the kinetics of ligand binding and the mechanism of signal transmission. In its low-affinity state, NTR binds its ligand with fast kinetics of association and dissociation (Rodríguez-Tébar et al., 1992; Mahadeo et al., 1994). This kinetic behavior probably indicates a large electrostatic component of interaction, which was observed

for the model both in force-field scores and in the proportion of charged residues involved in contacts. The location of charged ligand contacts at the top of subdomain III, near the boundary with subdomain II (possibly an interface with gp140trk), suggests a structural rationale for the role of NTR in the high-affinity receptor complex. This role appears to involve contributing its rapid on-rate of NGF binding to the complex with gp140trk (Mahadeo et al., 1994).

Signaling through NTR is poorly understood, although mutagenesis has implicated transmembrane sequences in the transmission process (Yan et al., 1991). Location of the second set of ligand contacts at the bottom of subdomain IV, on the boundary between folded structures and the receptor stalk, is consistent with evidence that truncation of the stalk by proteolysis interferes with chemical crosslinking of NGF to the receptor (Vissavajhala et al., 1992). Also, NGF binding to a stalk deletion mutant is enhanced to a greater extent by MC192 monoclonal antibody than is binding to NTR with a stalk, suggesting that the binding site can be stabilized either by bound antibody or the stalk (Welcher et al., 1991). The model complex thus provides a clue in the search for a connection between the ligand-binding domain, the stalk, and the transmembrane helix.

Applications of the model

Future uses of the model include study of how ligand binding initiates signaling. Preliminary investigation of changes in model structure upon complex formation suggests that small changes may occur in loop side chains and at intersubdomain hinges. The magnitude of these changes appears consistent with CD data (Timm et al., 1994). Signal propagation through the structure can also be explored, testing the hypothesis that conformational changes cascade through stacked loops. An application of this study might be to discover small molecule agonists that mimic the ability of NTs to prevent cell death induced by the NTR (Rabizadeh et al., 1993; Barrett & Bartlett, 1994). Potential therapeutic targets include age-related neurodegeneration and Alzheimer's disease, because NTR is highly expressed in basal forebrain neurons affected in these conditions (Rabizadeh et al., 1994).

Perhaps the most interesting application will come in probing the mechanism of NTR's affinity shift. Association with gp140trk subunits has been highly controversial despite evidence of participation by NTR in high-affinity complexes (Berg et al., 1991; Hempstead et al., 1991; Battleman et al., 1993; Benedetti et al., 1993; Barker & Shooter, 1994; Hantzopoulos et al., 1994; Mahadeo et al., 1994; Wolf et al., 1995). Failure of chemical crosslinking to reveal a physical complex between NTR and gp140trk (Meakin & Shooter, 1991) and a report that gp140trk homodimers account for NGF signaling (Jing et al., 1992) have suggested that NTR does not form a heterodimer with gp140trk. The NTR model presented here would guide mutagenesis experiments to identify the subunit contact zone, if it exists. The model structure suggests that subdomain II may contain this zone, and that hydrophobic side chains might be critical for subunit association. Experimental approaches guided by modeling have successfully characterized interactions between subunits of the IL-6 receptor (Savino et al., 1994). The model presented here provides an opportunity to resolve key questions about the structure and function of the NTR.

Materials and methods

Hardware and software

Computation was carried out on a variety of Silicon Graphics workstations, including a Personal Iris (MIPS R3000/20 MHz CPU, 32 MB RAM), an Iris Indigo (MIPS R4000/50 MHz CPU, 48 MB RAM), and a Challenge (MIPS R4400/150 MHz CPU, 256 MB RAM). Computer graphic analysis (Ferrin et al., 1988) used the software packages MIDASPlus™ (Computer Graphics Laboratory, University of California, San Francisco) and SYBYL (version 6, Tripos Associates, St. Louis, Missouri). Protein-protein interactions were evaluated with the DOCK3.5 software package (Molecular Design Institute, University of California, San Francisco).

Modeling by homology

Structural coordinates for TNFR-I and the TNF- β trimer (Banner et al., 1993) were generously supplied by the authors before release to the Brookhaven Protein Data Bank (1TNR). Amino acid sequences of TNFR-I, Swiss Protein database P19438 (Loetscher et al., 1990), and NTR, SwissProt P08138 (Johnson et al., 1986), were aligned using the six Cys residues in each subdomain, the conserved aromatic residue in the first loop of each subdomain, and the conserved hydrogen bonding residue in each third loop. A single gap per loop region was placed by aligning from both ends of the loop toward the center. This procedure left the gap in the least homologous part of the loop and farthest from structurally important residues. In some cases, the gap was moved one or two positions toward the amino- or carboxyl-termini to identify a better bridging peptide in the library supplied with SYBYL. Substitutions and loop replacements were made using modules of the SYBYL package. van der Waals conflicts resulting from side chain substitutions were resolved by rotamer library and molecular mechanics functions in SYBYL. Energy minimization was carried out using the AMBER all-atom force field (Weiner et al., 1986), a distance-dependent dielectric constant of 4.0 R, a nonbonded atom cut-off of 10 Å, and either the conjugate gradient (Fletcher & Reeves, 1964) or Powell (Powell, 1964) method. The model was subjected to molecular dynamics simulations using a quenching procedure. The temperature was raised to 300 K for 10 ps in 1-fs steps at unconstrained pressure, with Boltzmann initial velocities and a random number seed. The force-field set-up was the same as that used for energy minimization. Snapshots taken at the end of independent 10-ps runs were energy minimized as described above. No solvent was included with the model.

Docking

Coordinates for the crystal structure of mouse NGF (McDonald et al., 1991) were initially supplied by Dr. J. Murray-Rust (Birkbeck College, University of London) and later obtained as a revised set from the Brookhaven Protein Data Bank (1BET). Data files containing atomic coordinates were converted to standard Brookhaven Protein Data Bank format and, where necessary, stripped of hydrogen atoms and crystallographic waters using SYBYL. Essential hydrogens and AMBER united-atom charges (Weiner et al., 1984) were added. Solvent-accessible surfaces (Connolly, 1983) were computed using a 1.4-Å probe for each

structure with the program DMS (part of the MIDASPlus™ package). These were used to prepare docking sites and scoring grids. First, docking sites were defined using biochemical data. Then matching points were calculated with SPHGEN (Kuntz et al., 1982) and selected visually in MIDASPlus™. Scoring grids were prepared with CHEMGRID (Meng et al., 1992). To permit docking within the uncertainties of the model, we set all the surface atom radii of the docking site to zero, otherwise using AMBER atomic radii. This was tested with the TNF- β /TNFR crystal complex, which was regenerated from its separated elements. Clustered matching points representing several different receptor surfaces were used to orient the NTR model in the NGF docking site. DOCK3.5 parameters were set to allow up to eight van der Waals contacts during the matching phase, and the oriented receptor was adjusted in the site using rigid-body minimization (Meng et al., 1993). Force-field scores, reflecting van der Waals and electrostatic interactions in the contact zone, were computed for the resulting orientations of the receptor in the docking site.

Acknowledgments

This work was supported by NIH grant GM-31497 (I.D.K.) and a grant from Parke-Davis Pharmaceutical Research. We are grateful to Drs. Judith Murray-Rust and Tom Blundell for providing NGF coordinates, and to Drs. David Banner, Werner Lesslauer, and Fritz Winkler for sharing the TNFR coordinates before publication. We thank A. Geoff Skillman, Connie Oshiro, and Tom Cheatham III for patience, excellent advice on software applications, and for helpful comments on the manuscript. We are indebted to Dan Gschwend, Diana Roe, and Cindy Corwin for unflagging DOCK software support.

References

- Baldwin AN, Bitler CM, Welcher AA, Shooter EM. 1992. Studies on the structure and binding properties of the cysteine-rich domain of rat low affinity nerve growth factor receptor (p75NGFR). *J Biol Chem* 267: 8352-8359.
- Baldwin AN, Shooter EM. 1994. Disulfide mutants of the binding domain of the rat low affinity nerve growth factor receptor (p75NGFR). *J Biol Chem* 269:11456-11461.
- Baldwin AN, Shooter EM. 1995. Zone mapping of the binding domain of the rat low affinity nerve growth factor receptor by the introduction of novel N-glycosylation sites. *J Biol Chem* 270:4594-4602.
- Banner DW, D'Arcy A, Janes W, Gentz R, Schoenfeld HJ, Broger C, Loetscher H, Lesslauer W. 1993. Crystal structure of the soluble human 55 kd TNF receptor-human TNF β complex: Implications for TNF receptor activation. *Cell* 73:431-445.
- Barker PA, Murphy RA. 1992. The nerve growth factor receptor - A multi-component system that mediates the actions of the neurotrophin family of proteins. *Mol Cell Biochem* 110:1-15.
- Barker PA, Shooter EM. 1994. Disruption of NGF binding to the low affinity neurotrophin receptor p75(LNTR) reduces NGF binding to trkA on PC12 cells. *Neuron* 13:203-215.
- Barrett GL, Bartlett PF. 1994. The p75 nerve growth factor receptor mediates survival or death depending on the stage of sensory neuron development. *Proc Natl Acad Sci USA* 91:6501-6505.
- Battleman DS, Geller AI, Chao MV. 1993. HSV-1 vector-mediated gene transfer of the human nerve growth factor receptor p75hNGFR defines high-affinity NGF binding. *J Neurosci* 13:941-951.
- Bazan JF. 1991. Neurotrophic cytokines in the hematopoietic fold. *Neuron* 7:197-208.
- Benedetti M, Levi A, Chao MV. 1993. Differential expression of nerve growth factor receptors leads to altered binding affinity and neurotrophin responsiveness. *Proc Natl Acad Sci USA* 90:7859-7863.
- Berg MM, Sternberg DW, Hempstead BL, Chao MV. 1991. The low-affinity p75 nerve growth factor (NGF) receptor mediates NGF-induced tyrosine phosphorylation. *Proc Natl Acad Sci USA* 88:7106-7110.
- Beutler B, Van Hufel C. 1994. Unraveling function in the TNF ligand and receptor families. *Science* 264:667-668.

- Camerini D, Walz G, Loenen WAM, Borst J, Seed B. 1991. The T-cell activation antigen CD27 is a member of the nerve growth factor tumor necrosis factor receptor gene family. *J Immunol* 147:3165-3169.
- Chandler CE, Parsons LM, Hosang M, Shooter EM. 1984. A monoclonal antibody modulates the interaction of nerve growth factor with PC12 cells. *J Biol Chem* 259:6882-6889.
- Clackson T, Wells JA. 1995. A hot spot of binding energy in a hormone-receptor interface. *Science* 267:383-386.
- Connolly ML. 1983. Solvent-accessible surfaces of proteins and nucleic acids. *Science* 221:709-713.
- Crofts A, Hacker B, Barquera B, Yun CH, Gennis R. 1992. Structure and function of the bc-complex of *Rhodospirillum rubrum*. *Biochim Biophys Acta* 1101:162-165.
- Daopin S, Cohen GH, Davies D. 1992. Structural similarity between transforming growth factor-beta2 and nerve growth factor - Response. *Science* 258:1161-1162.
- Davies AM, Horton A, Burton LE, Schmelzer C, Vandlen R, Rosenthal A. 1993. Neurotrophin-4/5 is a mammalian-specific survival factor for distinct populations of sensory neurons. *J Neurosci* 13:4961-4967.
- Drinkwater CC, Barker PA, Suter U, Shooter EM. 1993. The carboxyl terminus of nerve growth factor is required for biological activity. *J Biol Chem* 268:23202-23207.
- Drinkwater CC, Suter U, Angst C, Shooter EM. 1991. Mutation of tryptophan-21 in mouse nerve growth factor (NGF) affects binding to the fast NGF receptor but not induction of neurites on PC12-cells. *Proc R Soc Lond (Biol)* 246:307-313.
- Eck MJ, Sprang SR. 1989. The structure of tumor necrosis factor alpha at 2.6 Å resolution: Implications for receptor binding. *J Biol Chem* 264:17595-17605.
- Eck MJ, Ultsch M, Rinderknecht E, de Vos AM, Sprang SR. 1992. The structure of human lymphotoxin (tumor necrosis factor beta) at 1.9 Å resolution. *J Biol Chem* 267:2119-2122.
- Ferrin TE, Huang CC, Jarvis LE, Langridge R. 1988. The MIDAS display system. *J Mol Graphics* 6:13-27.
- Fletcher R, Reeves CM. 1964. Function minimization by conjugate gradients. *Computer J* 7:149-154.
- Hantzopoulos PA, Suri C, Glass DJ, Goldfarb MP, Yancopoulos GD. 1994. The low affinity NGF receptor, p75, can collaborate with each of the trks to potentiate functional responses to the neurotrophins. *Neuron* 13:187-201.
- Hempstead BL, Martin-Zanca D, Kaplan DR, Parada LF, Chao MV. 1991. High-affinity NGF binding requires coexpression of the *trk* proto-oncogene and the low-affinity NGF receptor. *Nature* 350:678-683.
- Hempstead BL, Patil N, Theil B, Chao M. 1990. Deletion of cytoplasmic sequences of the nerve growth factor receptor leads to loss of high affinity ligand binding. *J Biol Chem* 265:9595-9598.
- Heuer JG, Fatemie-Nainie S, Wheeler EF, Bothwell M. 1990. Structure and developmental expression of the chicken NGF receptor. *Dev Biol* 137:287-304.
- Hosang M, Shooter EM. 1985. Molecular characteristics of nerve growth factor receptors on PC12 cells. *J Biol Chem* 260:655-662.
- Ibáñez CF, Ebendal T, Barbany G, Murray-Rust J, Blundell TL, Persson H. 1992. Disruption of the low affinity receptor-binding site in NGF allows neuronal survival and differentiation by binding to the *trk* gene product. *Cell* 69:329-341.
- Ibáñez CF, Ebendal T, Persson H. 1991. Chimeric molecules with multiple neurotrophic activities reveal structural elements determining the specificities of NGF and BDNF. *EMBO J* 10:2105-2110.
- Ibáñez CF, Hallböök F, Ebendal T, Persson H. 1990. Structure-function studies of nerve growth factor: Functional importance of highly conserved amino acid residues. *EMBO J* 9:1477-1483.
- Ibáñez CF, Ilag LL, Murray-Rust J, Persson H. 1993. An extended surface of binding to *trk* tyrosine kinase receptors in NGF and BDNF allows the engineering of a multifunctional pan-neurotrophin. *EMBO J* 12:2281-2293.
- Itoh N, Yonehara S, Ishii A, Yonehara M, Mizushima SI, Sameshima M, Hase A, Seto Y, Nagata S. 1991. The polypeptide encoded by the cDNA for human cell surface antigen Fas can mediate apoptosis. *Cell* 66:233-243.
- Jing SQ, Tapley P, Barbacid M. 1992. Nerve growth factor mediates signal transduction through *trk* homodimer receptors. *Neuron* 9:1067-1079.
- Johnson D, Lanahan A, Buck CR, Sehgal A, Morgan C, Mercer E, Bothwell M, Chao M. 1986. Expression and structure of the human NGF receptor. *Cell* 47:545-554.
- Johnson MS, Srinivasan N, Sowdhamini R, Blundell TL. 1994. Knowledge-based protein modeling. *Crit Rev Biochem Mol Biol* 29:1-68.
- Jungbluth S, Bailey K, Barde YA. 1994. Purification and characterisation of a brain-derived neurotrophic factor neurotrophin-3 (BDNF/NT-3) heterodimer. *Eur J Biochem* 221:677-685.
- Korsching S. 1993. The neurotrophic factor concept - A reexamination. *J Neurosci* 13:2739-2748.
- Kuntz ID, Blaney JM, Oatley SJ, Langridge R, Ferrin TE. 1982. A geometric approach to macromolecule-ligand interactions. *J Mol Biol* 161:269-288.
- Large TH, Weskamp G, Helder JC, Radeke MJ, Misko TP, Shooter EM, Reichardt LF. 1989. Structure and developmental expression of the nerve growth factor receptor in the chicken central nervous system. *Neuron* 2:1123-1134.
- Lee KF, Bachman K, Landis S, Jaenisch R. 1994. Dependence on p75 for innervation of some sympathetic targets. *Science* 263:1447-1449.
- Lee KF, Li E, Huber LJ, Landis SC, Sharpe AH, Chao MV, Jaenisch R. 1992. Targeted mutation of the gene encoding the low affinity NGF receptor p75 leads to deficits in the peripheral sensory nervous system. *Cell* 69:737-749.
- Loetscher H, Pan YCE, Lahm HW, Gentz R, Brockhaus M, Tabuchi H, Lesslauer W. 1990. Molecular cloning and expression of the human 55 kD tumor necrosis factor receptor. *Cell* 61:351-359.
- Mahadeo D, Kaplan L, Chao MV, Hempstead BL. 1994. High affinity nerve growth factor binding displays a faster rate of association than p140(trk) binding - Implications for multi-subunit polypeptide receptors. *J Biol Chem* 269:6884-6891.
- Marano N, Dietzschold B, Earley JJ Jr, Schattman G, Thompson S, Grob P, Bothwell M, Atkinson BF, Koprowski H. 1987. Purification and amino terminal sequencing of human melanoma nerve growth factor receptor. *J Neurochem* 48:225-232.
- McDonald NQ, Hendrickson WA. 1993. A structural superfamily of growth factors containing a cystine knot motif. *Cell* 73:421-424.
- McDonald NQ, Lapatto R, Murray-Rust J, Gunning J, Wlodawer A, Blundell TL. 1991. New protein fold revealed by a 2.3-Å resolution crystal structure of nerve growth factor. *Nature* 354:411-414.
- Meakin SO, Shooter EM. 1991. Molecular investigations on the high-affinity nerve growth factor receptor. *Neuron* 6:153-163.
- Meng EC, Gschwend DA, Blaney JM, Kuntz ID. 1993. Orientational sampling and rigid-body minimization in molecular docking. *Proteins* 17:266-278.
- Meng EC, Shoichet BK, Kuntz ID. 1992. Automated docking with grid-based energy evaluation. *J Comput Chem* 13:505-524.
- Narhi LO, Rosenfeld R, Talvenheimo J, Prestrelski SJ, Arakawa T, Lary JW, Kolvenbach CG, Hecht R, Boone T, Miller JA, Yphantis DA. 1993. Comparison of the biophysical characteristics of human brain-derived neurotrophic factor, neurotrophin-3, and nerve growth factor. *J Biol Chem* 268:13309-13317.
- Powell MJD. 1964. An efficient method for finding the minimum of a function of several variables without calculating derivatives. *Comput J* 7:155-162.
- Rabizadeh S, Bitler CM, Butcher LL, Bredesen DE. 1994. Expression of the low-affinity nerve growth factor receptor enhances beta-amyloid peptide toxicity. *Proc Natl Acad Sci USA* 91:10703-10706.
- Rabizadeh S, Oh J, Zhong LT, Yang J, Bitler CM, Butcher LL, Bredesen DE. 1993. Induction of apoptosis by the low-affinity NGF receptor. *Science* 261:345-348.
- Radeke MJ, Misko TP, Hsu C, Herzenberg LA, Shooter EM. 1987. Gene transfer and molecular cloning of the rat nerve growth factor receptor. *Nature* 325:593-597.
- Raff MC, Barres BA, Burne JF, Coles HS, Ishizaki Y, Jacobson MD. 1993. Programmed cell death and the control of cell survival - Lessons from the nervous system. *Science* 262:695-700.
- Ramachandran GN, Sassiakharan V. 1968. Conformation of polypeptides and proteins. *Adv Protein Chem* 28:283-247.
- Rayment I, Holden HM, Whittaker M, Yohn CB, Lorenz M, Holmes KC, Milligan RA. 1993. Structure of the actin-myosin complex and its implications for muscle contraction. *Science* 261:58-65.
- Ring CS, Cohen FE. 1993. Modeling protein structures: Construction and their applications. *FASEB J* 7:783-790.
- Rodríguez-Tébar A, Dechant G, Götz R, Barde YA. 1992. Binding of neurotrophin-3 to its neuronal receptors and interactions with nerve growth factor and brain-derived neurotrophic factor. *EMBO J* 11:917-922.
- Rosenberg MB, Hawrot E, Breakefield XO. 1986. Receptor binding activities of biotinylated derivatives of beta-nerve growth factor. *J Neurochem* 46:641-648.
- Savino R, Lahm A, Salvati AL, Ciapponi L, Sporeno E, Altamura S, Paonessa G, Toniatti C, Ciliberto G. 1994. Generation of interleukin-6 receptor antagonists by molecular-modeling guided mutagenesis of residues important for gp130 activation. *EMBO J* 13:1357-1367.
- Sehgal A, Bothwell M, Chao MV. 1989. Gene transfer of truncated NGF receptor clones leads to cell surface expression in mouse fibroblasts. *Nucleic Acids Res* 17:5623-5632.

- Shoichet BK, Bodian DL, Kuntz ID. 1992. Molecular docking using shape descriptors. *J Comput Chem* 13:380-397.
- Shoichet BK, Kuntz ID. 1991. Protein docking and complementarity. *J Mol Biol* 221:327-346.
- Smith CA, Davis T, Anderson D, Solam E, Beckmann MP, Jerzy R, Dower SK, Cosman D, Goodwin RG. 1990. A receptor for tumor necrosis factor defines an unusual family of cellular and viral proteins. *Science* 248:1019-1023.
- Somers W, Ultsch M, De Vos AM, Kossiakoff AA. 1994. The X-ray structure of a growth hormone-prolactin receptor complex. *Nature* 372:478-481.
- Timm DE, Ross AH, Neet KE. 1994. Circular dichroism and crosslinking studies of the interaction between four neurotrophins and the extracellular domain of the low-affinity neurotrophin receptor. *Protein Sci* 3:451-458.
- Vissavajhala P, Leszyk JD, Lin-Goerke J, Ross AH. 1992. Structural domains of the extracellular domain of human nerve growth factor receptor detected by partial proteolysis. *Arch Biochem Biophys* 294:244-252.
- Weiner SJ, Kollman PA, Case DA, Singh UC, Ghio C, Alagona G, Profeta SJ, Weiner P. 1984. A new force field for molecular mechanical simulation of nucleic acids and proteins. *J Am Chem Soc* 106:765-784.
- Weiner SJ, Kollman PA, Nguyen DT, Case DA. 1986. An all atom force field for simulations of proteins and nucleic acids. *J Comput Chem* 7:230-252.
- Welcher AA, Bitler CM, Radeke MJ, Shooter EM. 1991. Nerve growth factor binding domain of the nerve growth factor receptor. *Proc Natl Acad Sci USA* 88:159-163.
- Wolf DE, Mckinnon CA, Daou MC, Stephens RM, Kaplan DR, Ross AH. 1995. Interaction with trkA immobilizes gp75 in the high affinity nerve growth factor receptor complex. *J Biol Chem* 270:2133-2138.
- Yan H, Chao MV. 1991. Disruption of cysteine-rich repeats of the p75 nerve growth factor receptor leads to loss of ligand binding. *J Biol Chem* 266:12099-12104.
- Yan H, Schlessinger J, Chao MV. 1991. Chimeric NGF-EGF receptors define domains responsible for neuronal differentiation. *Science* 252:561-563.