C A S E
OF A
WOMAN WITH THREE HANDS;
ILLUSTRATED BY
ANALOGOUS MALFORMATIONS IN THE LOWER ANIMALS.

BY
J. JARDINE MURRAY, F.R.C.S.E., BRIGHTON.

COMMUNICATED BY
C. H. MOORE, Esq., HON. SEC.

Received Nov. 10th.—Read Dec. 9th, 1862.

THE following case derives additional interest from the circumstance that no similar malformation appears to have been met with in the human species.

Its nature is in some measure explained and illustrated by referring to comparative teratology. I am permitted by the President of the Royal College of Surgeons of England to exhibit drawings of two preparations of double fore-foot in the pig,¹ in both of which instances the abnormal anatomy of the limb seems to be quite analogous to that of this woman's arm. It is to be regretted that the preparations have not been carefully dissected, for the arrangement of the soft parts might have assisted our specu-

¹ These preparations, which are both dried, are thus described in the Catalogue of the Museum of the Royal College of Surgeons:

"No. 285. A double foot of a pig (*Sus domesticus*). HUNTER."

"No. 286. The right fore-foot of a pig with a supernumerary foot attached to the carpus. W. BLIZARD."

lations on the abnormal anatomy of the case now under consideration. In both instances of double fore-foot in the pig, as also, I believe, in this double-hand, the duplicity of bony structure begins at the carpus.

By the kindness of my friend, Dr. Cobbold, I am enabled to exhibit drawings of a somewhat similar malformation occurring in birds.

Anne S—, æt. 38, residing in Brighton, is a well-developed, healthy, active, and intelligent woman. She is married and has had one child, now seven years of age, who is in every respect normal. She is not aware that any relative of hers has been, or is, the subject of malformation.

The left upper extremity is the only part of her body which is abnormal in development. The limb is very muscular. The shoulder-joint is natural. The external condyloid ridge of the humerus is strongly defined. The elbow-joint may be fully extended, but flexion cannot be carried further than is represented in fig. 1, and the impediment seems to be the unusually large coronoid process of the ulna. The muscles and tendons of the fore-arm are so prominent that it is not at first easy to decide whether there be a second radius or ulna; but after numerous examinations made under favorable circumstances, I feel sure that the duplicity of bony structure begins at the carpus. Pronation and supination are freely performed.

The following measurements show the relative proportions of the right and left fore-arms :

	Right.	Left (abnormal).
1. Round olecranon and bend of elbow .	9 $\frac{1}{4}$ inches.	11 $\frac{3}{4}$ inches.
2. Round middle of fore-arm .	9 "	9 "
3. Round wrist .	6 "	7 "

The photographs give a better idea of the position and relations of the double-hand than could be conveyed in words, and it is therefore only necessary to supply such information as they do not convey to the eye.

The supernumerary hand is somewhat smaller than that which it so strangely accompanies. The fingers of both

hands are clubbed at their tips. The middle and ring fingers of the supernumerary hand are webbed¹ as far as the proximal joints, and the movements of the fingers of this hand are somewhat stiff and imperfect. The thumbs seem to be represented only by the prominence on the dorsal surface seen in fig. 3. The bursa on the back of the wrist, and also the smaller one on the knuckle of the index finger of the larger hand, show that, while at work as a charwoman, the patient is in the habit of leaning on the flexed carpus; and, from the rigidly arched conformation of the metacarpal portion of the double-hand, it is evident that she could not lean on the palm. The usual position of the hands, when closed, is seen in fig. 1. The power of independent action of the fingers is very limited. No single finger can be retained fully extended while the other seven fingers are flexed. But if both index fingers be extended, then the remaining six fingers may be flexed; or the four fingers of either hand, together with the index finger of the other, may be extended, while the remaining three fingers are flexed. The index fingers cannot be flexed while the other fingers are extended, nor can the little fingers be extended while the others are flexed. The limited character of these movements is probably due partly to want of training and partly to peculiarity in the arrangement of the extensor tendons. The double-hand can grasp firmly, although the maximum of power is not equal to that of the right hand. Sensation appears to be equally acute in all the three hands.

The case affords interesting material for the speculations of the anatomist; and the teratologist will remember that, in the progressive development of the upper extremity in the foetus, the fore-arm is formed subsequently to the hand, and at an earlier period than the arm and shoulder.

¹ The web is due to absence of fission, and is properly an arrest of development; for, in the embryo, the rudiments of hands and feet do not at first present any division of the fingers and toes.

DESCRIPTION OF PLATE II.

Fig. 1.—Exhibits the usual position of the hands when closed, and the greatest possible extent of flexion of the fore-arm.

Fig. 2 —The palmar aspect of the double-hand. All the fingers are extended, and the imperfections of the second, third, and fourth fingers of the supernumerary hand are well shown.

Fig. 3.—The dorsal aspect. In a line with the cleft between the index fingers is seen a prominence representing the thumbs. The bursæ on the wrist and knuckle of the first finger of the larger hand are also noticeable.

The figures are taken from large photographs by Messrs. Hennah and Kent, copies of which are in the possession of the Royal Medical and Chirurgical Society. A plaster cast of the arm may be seen in the Museum of the Royal College of Surgeons of England.

Fig. 1.

Fig. 2.

Fig. 3.