ON A CASE
OF
HYDATID DISEASE OF THE LIVER,
AND
REMARKS ON THE TREATMENT OF SIMILAR TUMOURS.

BY
JOHN HARLEY, M.D. LOND., F.L.S.,
ASSISTANT-PHYSICIAN TO KING'S COLLEGE HOSPITAL, ETC.

Received Jan. 11th.—Read April 24th, 1866.

THE following case of hydatid disease of the liver, in which cure has been effected, presents, from the enormous size of the tumour, its communication with the gall-bladder, and the occurrence of hæmorrhage from the liver, so many points of interest that I have thought it worthy of bringing before the notice of the Society.

In June, 1863, Mr. B. G—, æt. 29, a gentleman of good development and vigorous constitution, applied for my advice on account of a large tumour which had arisen in the epigastric region about four years previously, and slowly increased to such an extent as to cause, when dressed, very conspicuous deformity of the chest and abdomen.

At this time the patient was moderately muscular, defi-

cient of fat, pale and sallowish, with dark areolæ around the eyes. All the functions were naturally performed. He had been, and continued to be, constantly engaged in an office in the City. Long standing induced a feeling of faintness; but he was able to go about as usual, and not unfrequently walked four miles without inconvenience. The lower half of the chest and the whole of the abdomen were enormously but uniformly distended by a dull, tense, fluctuating tumour. The lower ribs, as well as those bounding the lower part of the chest-wall, were pushed widely outwards. The distance between the point of the ensiform cartilage and the umbilicus was much increased, and the most prominent part of the tumour corresponded to the centre of this elongated space. As the tumour ascended beyond the margin of the chest-wall in front, and at the sides where the parts were very much stretched, no limit could be felt in the upward direction. Below, the tumour gradually subsided towards the pubic region, and its lower rounded border could be distinctly felt two fingers' breadth above the pubis and the inner ends of Poupart's ligaments. From this lower boundary, and excepting a narrow space on either side of the abdomen, uninterrupted dulness extended upwards to all parts of a horizontal line surrounding the chest on a level with the nipples. Percussion upon either of the distended regions produced distinct fluctuation throughout the whole of the tumour. Neither the liver, the spleen, nor any other solid body could be felt. Two little veins emerged at the right edge of the ensiform cartilage, and formed swollen ramifications as they passed upwards and disappeared.

Over the most prominent part of the tumour—a point midway between the ensiform cartilage and the umbilicus—the body was $40\frac{1}{2}$ inches in circumference.

Above the line of dulness, the lung-sounds were natural; but there was an absence of vesicular murmur, and distinct bronchophony was heard at the bases of both lungs behind. On taking a deep inspiration, a rubbing sound, like that caused by the motion of two pieces of wet leather the one

upon the other, was heard in the left scapular region. The heart was displaced upwards, and the cardiac dulness corresponded to a space of about two inches in diameter, the centre of which was half an inch above and to the sternal side of the left nipple. The heart-sounds were distinctly heard everywhere over the front of the chest on the left; they were most distinct a little above and to the left of the nipple, and thence towards the axilla. Over the natural position of the heart, and thence to the margin of the left ribs, the sounds were peculiarly naked and tense, as if transmitted from a distance by a good conducting medium. A moderately loud systolic bruit was heard at the base near the left edge of the sternum, on a level with the nipple; and, a little to the left of this, a fainter diastolic bruit. The pulse was natural, and the volume moderate.

Previous history.—The patient has lived temperately, is free from constitutional disease, and has always enjoyed good health; is liable to boils, but was never laid up with any illness except in connection with the disease under consideration; has never had rheumatism, nor any cardiac or pulmonary affection; never had jaundice or tapeworm; has lived seven years in Australia, and been exposed to the hardships incidental to a country life there; was often obliged to drink stagnant, and sometimes muddy water, and always preferred imperfectly cooked meat. He returned from Australia in the year 1859. In March of this year, and while still in Australia, he was attacked with severe stabbing pains in the left side of the abdomen, a little towards the front. For twenty-four hours the pain was continuous and agonising, and he could neither lie down nor suffer himself to be touched. The pain did not produce vomiting, but it was accompanied by a dry cough and sharp pain on inspiration. The pain recurred at intervals for a fortnight, and then gradually decreased. Since this time he has occasionally had a slight return of pain in this situation; and as pressure upon the left induces it, he constantly lies down upon the right side. About this time an abundant herpetic (?) eruption appeared upon the trunk and

limbs, and continued for some months. Some finely pitted scars, attesting its pustular nature, still remain upon the legs.

Three or four months after this severe attack of pain, in July 1859, while on his passage home, he discovered a little fulness of the epigastrium below the margin of the ribs ; and since this time he has lost a little flesh, and his strength has gradually diminished.

In 1860 he had another attack of severe pain in the same part. He was laid up for a fortnight, and in the second week there was constipation for five days. Since this time the tumour has gradually increased to its present size. His health has continued fair, and his occupations have only been interrupted twice or thrice, and for only a day or two at a time, by the recurrence of the old pain and a feeling of weakness.

From the history of the case, and the nature and situation of the tumour, I concluded that the patient was the subject of hydatid tumour of the liver. As he enjoyed good health and his occupation required no great physical exertion, and as the tumour had already attained so great a size, I thought it better to recommend no present interference. I explained the nature of the disease to the patient, and directed him to avoid straining, jolting, and all violent motion, and to be slow and careful in all his movements. As his condition was somewhat anæmic I prescribed chalybeate medicine, and requested him to carefully notice if there should be any increase in the size of the tumour, and to see me from time to time.

October 8th, 1863.—Has enjoyed fair health and pursued his usual occupations continuously since June, and he is now in the same condition as at that date. Complains that he occasionally has deep-seated pain about the upper angle of the left scapula. It is sometimes very troublesome in bed. At times he experiences a deeply-seated pain in a single spot behind the left edge of the sternum, in the situation of the heart. Measurement around the body, at the same level as before (midway between ensiform cartilage and umbilicus),

shows an increase of half an inch, viz., 41 inches. The lower border of the tumour is rounded and distinct in the right iliac region; towards the left iliac region, a harder and more solid, thick, and slightly uneven boundary is felt. I take it to be the anterior margin of the left lobe of the liver.

July 7th, 1864.—Measurement over the same part $41\frac{1}{4}$ inches. No increased dulness of the chest, and the heart- and lung-sounds have not undergone any alteration. The patient complains of pain transmitted from below to a very tender little swelling over the junction of the manubrium and second piece of the sternum, whence it extends to the left shoulder and down the inside of the left arm.

In all other respects he has continued well, and there has been an increase in girth of only three fourths of an inch during the last year.

June 15th, 1865.—Measures around the same part $42\frac{5}{8}$ inches. The tumour has descended a little, so as to leave another inch of horizontal resonance of the chest. The abdomen is fuller and the tumour more tense; otherwise the condition of the patient is the same as it has been during the last two or three years.

The tumour was at this time so tense, and the cyst-wall appeared so thin, that I now began to feel great anxiety lest it should be ruptured by some accidental strain. I cautioned the patient to avoid everything that might lead to such an issue, and determined to adopt at an early period some operative means for his relief.

July 17th, 1865.—On getting out of bed this morning, the patient was seized with a severe darting pain in the right iliac region; the pain increased at intervals to paroxysms of intense agony. The paroxysms were announced by a severe stabbing pain, which soon assumed an excruciating twisting and bursting character. While the pain continued the part was very tender to the touch; but in the intervals there was no particular tenderness, but only a fear lest palpation should excite the pain afresh.

I was summoned to him on the 21st of July, and saw him

in consultation with Dr. John Meaburn Bright, of Forest Hill, who was in attendance upon him. I found that the tumour had increased considerably in size in the right iliac region; but as yet there was no indication of the rupture of the sac, and the peritoneal inflammation was of very limited extent. Believing that a slight increase of fluid within the sac would determine its rupture and cause the death of the patient, I resolved to puncture the tumour. Dr. Bright concurred with me as to the necessity of this step, but we deferred the operation, hoping that the inflammatory action would result in adhesion of a portion of the sac to the abdominal wall, on account of which we might puncture the tumour with greater safety. For the present, therefore, we watched carefully and employed only palliative treatment.

July 26th.—The intervals of freedom from pain have been longer, and yesterday he was altogether free from pain for six hours. Since the 17th inst. the abdominal portion of the tumour has considerably increased in size. It is everywhere rounded; smooth, very tense, and everywhere fluctuant, and the fluid is evidently included within a single cyst. The pulse has lately risen to 120 during the paroxysm; and to-day there is superficial tenderness over the right hypochondriac and iliac regions, where the distension has lately increased very much. The patient is free from feverishness; the bowels have been well relieved by castor oil and enemata; opiates have given relief, but the patient has little or no continuous sleep.

The following are the measurements taken this day:—Round the body on a level with the tip of the ensiform cartilage, $40\frac{3}{8}$ inches; midway between the tip of the ensiform cartilage and the umbilicus, where the former measurements were made, $44\frac{1}{2}$ inches, which is nearly 2 inches more than the measurement taken only a month previously. Over the umbilicus, 40 inches; midway between the umbilicus and the pubes, $37\frac{3}{4}$ inches; from the tip of the ensiform cartilage to the pubes, $13\frac{1}{2}$ inches. As the tumour was undergoing such rapid enlargement, I resolved to delay no longer in evacuating the fluid. The patient willingly acceded. On further

consideration, I concluded that it would not be safe to tap in the right hypochondriac region, for I did not feel sure of the adhesion of the sac in this situation; I therefore selected a point in the median line midway between the ensiform cartilage and the umbilicus, where the tumour was most prominent. The patient kept the recumbent posture, and bandages having been previously applied around the margin of the ribs and the abdomen, Dr. Bright introduced a No. 11 canula, which on withdrawal of the trocar readily slipped backwards to its shoulder; the trocar having been introduced, with the point directed very slightly upwards, for the distance of only an inch.

Dr. Ray, of Dulwich, and Mr. Hill, of Sydenham, were present at the operation, and kindly gave us their assistance.

Immediately that the trocar was removed, fluid as limpid, clear, and colourless as water, was ejected with great force. The complete evacuation of the fluid occupied two hours. I occasionally interrupted the flow for a minute or two at a time, in order to secure the more effectual contraction of the cyst. The last two or three pints oozed slowly away, and the discharge was facilitated by pressure on the abdomen and sides of the chest, and ultimately by causing the patient to take deep inspirations. Nineteen pints and a half (390 ounces) were ultimately withdrawn. The greater portion of the fluid was colourless; but towards the end of the operation it gradually assumed a light yellowish colour, which increased in intensity. The last pint was grumous and of a bright yellow colour from admixture with bile. Not a trace of blood appeared throughout. Some ruptured hydatid cysts of gelatinous appearance, about the size of walnuts, and a number of minute unruptured ones, the size of peas, came away with the fluid. The evacuation of the fluid was unattended by pain, and gave the patient immediate relief. There was no tendency to syncope; the abdomen became quite flaccid and free from pain on pressure, and the rounded outline of the shrunken cyst could be felt distinctly just below the umbilicus, and thence extending upwards on the sides

towards either hypochondrium, where the outlines were indistinct. At the end of the operation, a little fluid continued to run from the canula on taking a deep inspiration; and as I had determined not to remove the canula so long as there should be any oozing of fluid, a pad of lint and a folded towel were placed over the orifice, and the whole retained by moderate pressure with the abdominal bandage.

Flannel rollers were tightly applied around the lower margin of the chest. The pulse after the operation was 95, soft and compressible. To take 10 grains of compound soap-pill to-night.

The morning after the operation, the patient was in a very comfortable condition. He had passed a good night, and was now free from pain. The abdomen soft, flaccid, and free from tenderness. Pulse 124; respirations 30. A severe eruption of urticaria covers the neck and arms. The bandages and compresses are saturated with discharge, and stained bright yellow. The canula was worn without inconvenience, and a yellow bilious fluid containing a little mucus constantly oozed from its orifice.

Three days after the operation, a large rounded tumour about the size of a child's head could be felt in the right hypochondriac and epigastric regions. In common with the rest of the abdomen, these parts were free from tenderness.

The patient continued free from febrile disturbance and pain, and progressed most favorably, living upon a liberal allowance of mixed animal and vegetable diet and stimulants, until the *eighth* day, when, the canula having partially slipped out of the wound, there was complete retention of the discharge for about twelve hours. The distended sac was tender, and formed a rounded prominence in either hypochondrium; there was pain along the margin of the left ribs; the skin was hot; the countenance very sallow. Pulse 120; respirations 17. On replacing the canula, eight ounces of deep yellow turbid fluid of offensive odour were immediately evacuated, and free discharge of the contents of the sac was re-established. The febrile symptoms disappeared, and all went on well until the membranous walls

of disintegrated hydatid cysts began to obstruct the discharge. Fragments of laminated membranes first appeared on the *tenth* day ; but they did not cause any obstruction until the *fifteenth* day, when there was complete arrest of the discharge. The sac was distended, and the pain along the margin of the left ribs and febrile action were renewed. An elastic catheter was introduced through the canula to a distance of nine inches within the sac, and with some difficulty twenty ounces of light ochre-coloured, slightly viscid fluid, and much broken-down cyst-wall, were evacuated. This was followed by immediate relief.

For the first fifteen days, about twenty ounces of fluid were discharged every twenty-four hours, but on several occasions it was much more abundant ; and this temporary increase was apparently due to rupture of the larger secondary cysts.

From the *fifteenth* to the *seventy-eighth* day, disintegrated cyst-wall was constantly discharged, causing impediment, and occasionally, for a few hours at a time, complete obstruction to the discharge of the fluid contents of the sac. During the whole of this period great difficulty was experienced in emptying the sac.

On the *forty-third* day, the canula was removed, and a No. 9 elastic catheter inserted and retained in place of it. From this date to the *fiftieth* day the discharge diminished to four or six ounces daily, and the sac meanwhile became distended. On the last-mentioned day, twenty ounces of foetid fluid were withdrawn, and the discomfort and feverishness attendant upon the distension of the sac abated.

The day after, a new and dangerous symptom appeared. Hæmorrhage to a considerable extent took place within the sac. The pulse went up from 96 to 140, and in the evening to 160. Vomiting had occurred several times ; the skin was hot, dry, and jaundiced ; the epigastric and hypochondriac regions were occupied by the hard and distended sac, but the rest of the abdomen was soft and natural. A pint of thick, foetid, bloody fluid, of the same rich chocolate-red colour as the blood which exudes from the cut surface of

fresh liver, was removed from the sac. Fragments of cyst-wall continually obstructed the eye of the catheter, so that very little discharge now oozed from its outer extremity. By patiently withdrawing and introducing the catheter time after time, and injecting creasote water through it to wash away the débris of the cysts, half a pint more of the bloody fluid was removed.

For many days the fæces had been completely destitute of bile, and were of a glistening aspect and putty colour. This evening the patient passed a very copious pultaceous shiny stool, having the same rich reddish-brown colour as the discharge from the sac. It was clear that the distended sac had relieved itself by the passage of a portion of its contents into the intestinal canal.

During the next week, much disintegrated blood-clot came away, and oozing of blood within the sac continued. The patient meanwhile was much weakened and depressed. The jaundice disappeared, and the appetite had returned on the *fifty-third* day. The alvine secretions, however, continued to be entirely destitute of bile, and they were very offensive. The discharge from the sac had contained a large quantity of bile from the first, and the proportion of this secretion appeared now to be on the increase. After the sac had been washed out, a stream of pure, unmixed ropy bile flowed from the orifice of the catheter as it lay inserted eight or nine inches within the sac. I collected half an ounce of this pure bile in the course of a few minutes, on several occasions. It was evident that we were in direct communication with the gall-bladder, and that the discharge from the sac, which some days previously had found its way into the intestine, must have been conveyed by the bile-duct. The discharge of pure bile at the end of dressing continued from the *fifty-seventh* to the *sixty-fifth* day. The discharge from the sac then resumed its usual bilious character, and decreased in quantity. The fragments of the cysts continued to offer a great obstruction to the flow of the discharge, although the passage leading to the sac was now sufficiently dilated to admit three large elastic

catheters. But on the *seventy-fifth* day we succeeded in evacuating a teaspoonful of thick laminated membranes, some of which were six inches square. This was effected by moderately distending the sac with creasote water, and suddenly withdrawing the three catheters, the patient giving at the same time a series of little coughs. Excepting a stray fragment or two, the whole of the laminated membranes that had been retained in the sac were removed on this occasion, and henceforward there was no impediment to the flow of the discharge, nor difficulty in washing out the sac. On two occasions, the sixty-second and the seventy-fifth days, large thick fragments of dense blood-stained fibrous tissue came away. These were, no doubt, portions of the original cyst which had become detached from the liver. The hæmorrhage was clearly due to the separation of these fragments, and the exposure of the parenchymatous substance of a portion of the liver.

On the *eightieth* day, the patient had recovered his lost ground, and the fæces had returned to a natural condition.

On the *eighty-first*, a severe attack of dyspepsia, accompanied by great flatulent distension of the abdomen, and violent eructations and vomitings, occurred. The attack lasted with decreasing severity for a week; meanwhile the discharge from the sac increased. Of late the average had been eight ounces in the twenty-four hours, and it was a dirty yellowish-white muco-purulent fluid; but at this day, and for the following week, from sixteen to twenty ounces of a grass-green mucous fluid—diluted bile—were daily discharged; and this was associated with absence of bile from the alimentary canal. The sac, however, was daily contracting, and I began to fear that the common bile-duct would be so implicated in the contraction as to become obliterated. It was with extreme satisfaction that Dr. Bright and myself witnessed the reappearance of bile in the stools on the *one hundred and second* day.

During the next month, the bilious discharge gradually decreased.

On the *one hundred and twenty-third* day, the discharge was totally free from bile, and consisted of dirty-white inodorous pus, amounting to three ounces only in the twenty-four hours. The sac meanwhile contracted, and the catheter, which formerly passed nine or ten inches into its interior, would now pass only four inches.

On the *one hundred and thirty-eighth* day, a tinge of bile appeared for the last time. The sac had almost entirely contracted. The discharge consisted of a dirty-white inodorous pus, and did not exceed a table-spoonful in the twenty-four hours: the catheter merely occupied the retracting passage, and not more than two drachms of fluid could be injected.

On the *one hundred and forty-eighth* day, the discharge had ceased, and the catheter was finally removed. During the last month the patient had rapidly regained flesh and strength, and he was able to leave his room and walk out.

On the *one hundred and fifty-eighth* day (January 1st, 1866), the patient resumed his ordinary occupations in the City, and has continued them without interruption up to the present time, with the enjoyment of increasing health and strength.

The following was his condition on the 1st of January, 1866, rather more than five months after the puncture of the tumour:—A little stouter than he was before the operation, but pale and anæmic. Measurement around the body at the level above indicated—viz., midway between the ensiform cartilage and the umbilicus— $31\frac{1}{2}$ inches, being 13 inches less than the measurement on the day when the tumour was punctured, and 9 inches less than when the patient first came under my notice, in June, 1863. The cicatrix retracted upwards, and the parts around dull and hard. No trace of the tumour remains. The hepatic dulness normal. The spleen feels hard, and it is enlarged, projecting below the margin of the left ribs. The appetite is good; the bowels act regularly, and the fæces are quite healthy. The compressed lungs have expanded, and

the extent of resonance and vesicular breathing is normal. The heart has returned to its proper position; the first sound is now healthy, and the second is accompanied by only a faint diastolic bruit.

I made my last examination of my former patient on the 7th September, 1866, when these sheets were in the printer's hands. The following are the particulars:—Body fat and muscular, and complexion healthy; weighed before the operation from eleven stone to eleven stone three pounds; weighs now eleven stone, showing an increase, *minus* the tumour, of about two stone; figure perfectly natural. The cicatrix looks exactly like a second navel, situated midway between the tip of the ensiform cartilage and the umbilicus. The hypochondriac regions and the epigastrium, including the parts immediately around the cicatrix, are perfectly soft, and the fingers deeply pressed into these regions fail to detect any tumour or thickening whatever, or to excite pain or uneasiness. The epigastric and left hypochondriac regions are resonant. The liver dulness is of normal extent, commencing four fingers' breadth below the right nipple and terminating at the margin of the ribs. The abdomen is flat and natural, and the spleen has returned to its normal size and position—indeed it can no longer be distinguished. Measurements:—From the tip of ensiform cartilage to the umbilicus, $5\frac{1}{4}$ inches (the cicatrix is equidistant from these parts, inclining a little to the left of the median line, and is slightly retracted upwards); around the body, on a level with the cicatrix, $33\frac{1}{2}$ inches; and on a level with the ensiform cartilage, 34 inches. Resonance of the chest of normal extent; lung-sounds healthy. Cardiac dulness normal in extent and situation, apex-beat an inch below, and internal to the left nipple. The systolic bruit is no longer heard, the first sound being strong and healthy; a distinct sawing sound accompanies the diastole. Its seat is immediately beneath the junction of the first and second bones of the sternum. The pulse is natural; all the functions are healthily performed. Although his work has been unusually heavy since he resumed it

in January, he has progressively improved in health and strength.

Treatment.—As there was no evidence of the adhesion of the walls of the cyst to the abdominal parietes, I feared to do less than withdraw at once the chief bulk of the fluid; and when I found that the elasticity of the sac was such as to expel the fluid with extreme force, I had reason to be satisfied that I had not adopted partial measures, such as capillary tapping or the removal of only a portion of the fluid. In this case either of these modes of treatment would most probably have been followed by the escape of hydatid fluid into the peritoneal cavity. Anticipating that the sac would rapidly refill—in which case the introduction of the trocar and withdrawal of the canula would have been almost equivalent to rupturing the sac and allowing its contents to escape into the abdomen—I resolved to retain the canula within the wound as long as there should be any discharge.

Dr. Bright used a medium-sized instrument, and the treatment consisted in placing a flannel bandage around the abdomen, below the canula, with a compress upon the right hypochondrium, and another bandage around the lower part of the chest above the canula, which latter was retained in its position by placing folded diapers over its orifice and retaining them by another light bandage. The canula was retained within the wound until the 7th September, when having become loose and inclined to slip out of the wound, it was removed, and an elastic catheter (No. 9) introduced into the sac in its stead. Prior to this, the discharge of the cyst-wall of the disintegrated hydatids began to obstruct the discharge of the fluid which constantly accumulated within the sac. Fearing that an elastic catheter would not be effectual in maintaining a direct passage into the sac, and that there might be some difficulty in reintroducing it, I wished to preserve the communication with the sac by means of a curved silver tube. A No. 12 double-canuled silver catheter was readily introduced nine inches within the sac, but it proved useless. I therefore procured a

silver tube, curved like an ordinary catheter, of No. 16 gauge, a little contracted at the point to facilitate its introduction, and terminating in a round orifice of No. 11 gauge. This tube passed without much difficulty into the sac, and it was worn with comfort for three days but not an ounce of discharge flowed away through it during the whole of this time, and the accumulated discharge was evacuated by other means at the times of dressing. On visiting the patient on the evening of September 22nd, we found that no fluid had been discharged through the silver tube, although the sac was distended, and the wide tube moved freely in every direction in the accumulated fluid. It was removed; the discharge was evacuated with difficulty by means of an elastic catheter. The silver tube was again introduced, and about a pint of creasote water injected; none returned until the sac was painfully distended, when about half a pint was rejected with a sudden gush; more could not be obtained, although the tube was moved to every part of the sac, and inserted now a short way, and now seven or eight inches within it. It was evident that the orifice of the tube was entangled amongst the detached laminæ of the hydatid cysts, which being forced towards it in every direction by the distension of the sac, occluded it like so many valves. The same thing occurred when a single catheter was introduced, and we now began to experience the greatest difficulty in evacuating the contents of the sac. The mode in which it was effected for two or three weeks about this time was as follows:—A No. 9 elastic catheter was inserted into the sac; a little discharge would then flow for a second or two, and then an arrest occurred; a fragment of cyst-wall was removed from the orifice of the catheter, or a little creasote water injected to wash it out of the internal orifice, and the discharge would perhaps again flow. By this means, and by frequently removing the catheter and washing cyst-wall out of its eye, and by teasing away fragments which had followed it to the orifice of the wound, the sac was emptied and washed out with cold creasote water (℞ Kreasoti m℥i, Aquæ ℥xxxvj) every night

and morning. Dr. Bright attended the patient night and morning throughout the progress of the case, and to his assiduous and patient attentions the safety of the patient was due. The time required to evacuate the contents of the sac and the difficulty in effecting it, now required the adoption of some means to effect the dilatation of the passage for the discharge of the larger coriaceous fragments of cyst-wall. For this purpose I employed elastic catheters : we left three small ones (Nos. 9, 7, and 4) in the sac, and by daily increasing their size to No. 12 our end was attained. When the catheters were introduced within the sac so that their extremities reached to slightly unequal distances, the cyst-wall was generally kept away from the orifice of one at least, and while creasote water was being injected along one catheter, discharge accompanied by cyst-wall was ejected from the others. By this means thickish laminated fragments six square inches in extent were discharged. On October 11th, the passage was sufficiently dilated to allow of the discharge of a teacupful of large leathery fragments of cyst-wall—all, in fact, that now remained within the sac. The ejection of these was secured by injecting the sac with creasote water and preventing its escape, and then suddenly withdrawing the catheters while the patient gave a succession of coughs.

While we were endeavouring to secure free discharge of the fluid formed within the sac, I had two other objects in view : viz., 1st, to prevent decomposition of the contents of the sac ; and, 2nd, to excite inflammatory action within its interior. These were attained by the injection of iodine and creasote water : $\mathfrak{zj}$ of the compound tincture of iodine of the London Pharmacopœia was injected into the cyst three days after it was tapped (July 28th) ; and this was repeated morning and evening for about a week, when the quantity was increased to $\mathfrak{zij}$ twice a day. From the 14th to the 23rd of August, $\mathfrak{3j}$ of the tincture was injected through the canula daily. The evening injection was now discontinued on account of the profuse night-sweats which it appeared to produce, and $\mathfrak{zvj}$ were injected every morning only, until the

17th of September, when the injection of iodine was left off altogether. During these seven weeks, upwards of ℥xxx of the tincture of iodine, diluted with an equal quantity of creasote water, were thrown into the sac. No pain was ever produced; but when the sac became a little tender, a feeling of warmth in the epigastrium followed the injection.

The iodine injection was continued until hæmorrhage occurred within the sac, when gr. v—gr. x of nitrate of silver, dissolved in a few ounces of water, was substituted, the sac being previously washed out as before with creasote water. At the end of a week I discontinued the nitrate of silver, and directed the sac to be first washed out with plain creasote water, and, subsequently, with ℥j of sulphate of zinc dissolved in ℥x of creasote water, every morning and evening.

Throughout the whole treatment I found it of the utmost importance to give support to the lower part of the chest, and had laced bandages made for this purpose. The ribs had been so widely distended, that if left to themselves they were laxly divergent. It was equally necessary to keep the abdomen tightly bandaged, with a compress upon the right side so as to press the contracting sac towards the right hypochondrium. As often as the compress was set aside, a slight distension took place in that most dependent part of the sac which lay in the right iliac region. Another difficulty that we had to contend with was the deficiency, and, for a very considerable portion of the time, a total absence of bile from the intestines. Fortunately, the patient's appetite did not suffer in consequence, and I attempted to supply the place of the absent secretion by the introduction of inspissated ox bile, in doses of 20 grains, every night, with occasional intervals. For the greater part of the time he took ℥xv—xx of perchloride or perntrate of iron, alone or in combination with quinine.

At one period (Oct. 16—20) the patient was greatly troubled with flatulent dyspepsia, and this was associated with fæcal accumulation in the ascending colon, and it was necessary to employ free purging. The accumulation was,

doubtless, to be attributed to adhesions, which, having formed between the sac and the right bend of the colon, impaired the contraction of this part of the alimentary canal. Pressure upon the right hypochondrium at these times readily induced eructations.

General observations and rules for treatment.—It is remarkable how little disturbance of the sympathetic system occurred throughout the treatment of the case, and how readily the displaced and compressed viscera resumed their position and recovered from the effects of pressure so prolonged. The tumour was seated in the midst of the sympathetic plexuses, and caused distension of all parts about the solar plexus. The sac evidently lay upon the aorta, for whenever a catheter was passed backwards until its point touched the posterior wall of the sac, strong pulsations were felt; at one time it was easy to feel that the sac had contracted extensive adhesions with the diaphragm along the margin of the ribs upon the left side, and whenever it became distended pain was felt in this region. The absence of bile from the intestinal canal was always associated with a disgust for fat of all kinds; and the putty-coloured fæces contained undigested, unemulsified oil, which gave to them a soft glistening appearance, like that of frosted silver.

It would appear that the cyst was in an actively growing state; no cretaceous matter was ever observed in the discharges. The first ten or twelve pints of fluid withdrawn were of the usual character—a clear limpid, slightly salty fluid, destitute of albumen, and of specific gravity 1007. The last portions were turbid, from the presence of yellow grumous matter, altered and inspissated bile. Each drop of the turbid deposit contained multitudes of hooklets, free scolices, and minute cysts, the $\frac{1}{80}$ th of an inch in diameter, containing numbers of attached scolices in all stages of development. The parasite was evidently derived from Australia, but the scolex is identical in appearance with that found in hydatid cysts of European origin.

The pure bile which was occasionally discharged un-

doubtedly came from the gall-bladder, for it contained so much mucus and was so viscid that it could be drawn out into strings more than a foot long. It was equally clear that the great cyst did not directly communicate with the gall-bladder, else the whole of the fluid would have been tinged with bile. I can only explain the fact by supposing that a small and probably collapsed hydatid cyst existed beside the large one, and communicated, on the one hand, with the gall-bladder, and, on the other, with the larger cyst, and that during the growth of the tumour the aperture of communication was closed either by the secondary cysts or by the compressed fluid in some such way as the orifices of the ureters are closed by a distended bladder.

Having observed this complicated case with much interest, and given it close attention, and subsequently studied the histories of those cases in which cure by operative measures has been attempted, I am convinced of the necessity of observing the following rules in the treatment of hydatid tumours of the liver :

1. They should be punctured *above* the umbilicus, because the sac, however large, possesses great elasticity, and ultimately contracts into the epigastric or hypochondriac regions.

2. As soon as operative measures are determined upon, the sac should at once be punctured with a large trocar, and the canula retained so as to keep up free communication with the interior of the sac. No care need be taken to prevent the admission of air; there is much less danger in exposing the interior of a large cyst to the air for months than in closing up the external orifice for a few days. Nor ought we to be content with merely keeping a canula inserted; but by passing a small catheter from time to time within it, should anticipate obstruction of the discharge and distension of the sac by pushing away, breaking down, or teasing-out the secondary cysts or their fragments, which, so long as they remain within the sac, will continue to impede or prevent the flow of fluid, however large the orifice of the canula may be.

3. The canula may be retained until it is loosened by suppuration and tends to slip out. Its place should then be supplied by two or three elastic catheters, and their size gradually increased until three or more of No. 12 size can be readily introduced. By advancing the ends of two beyond the others, and causing the eyes to look inwards towards each other, we have an effectual contrivance for preventing the closure of some, at least, of the eyes by the pliable fragments of cyst-wall. A single catheter, or a single wide tube, is ineffectual in emptying the sac.

4. In order to facilitate the disintegration of the cyst-wall, and to excite inflammation and adhesion of the sac, iodine should be freely injected. As soon as a glow of heat is felt on injecting it within the sac, or this becomes tender under pressure, the iodine injection should be discontinued.

5. To prevent decomposition of the fluid within the sac, it should be washed out morning and evening with creasote water (creasote 40 drops, water 36 ounces). I found this most effectual in preserving a healthy condition of the sac. On one occasion more than a gallon of cold creasote water was passed through the sac. The injection should be passed down one catheter, and allowed to flow out with equal rapidity through one or two others. Distension of the sac should be most carefully avoided, especially when it is healing. Pain, and, on more than one occasion, bleeding, was produced in the case above related by injecting the fluid faster than it flowed away.

6. After the separation and discharge of the cyst-wall, the sac may be washed out with a solution of sulphate of zinc in creasote water. I employed it in this case chiefly with the view of arresting and preventing hæmorrhage, but there can be no doubt that it facilitated the contraction of the sac, and I should be inclined to recommend its use in all cases during the healing of the sac.

7. For treating tumours of great size, the strictest attention must be given to bandaging, in order to preserve the contraction of the sac. In the case above recorded the

sac had evidently contracted such firm adhesions to the diaphragm and inferior margin of the chest-wall that, being firmly retained in the epigastric and hypochondriac regions, it extended itself chiefly in the lateral directions, pushing the inferior parts of the chest and lower ribs much outwards. Long after the softer parts had regained their resiliency I found that the bony chest-wall remained flaccid and spread outwards, and that by compressing the hepatic and splenic regions together the capacity of the sac was diminished, as evidenced by the escape of air or fluid. A laced bandage should be placed around the lower part of the chest, embracing the hepatic, splenic, and epigastric regions, before the sac is punctured, and it should be constantly worn and tightened during the contraction of the sac, and for some months after the patient has recovered. Support should equally be given to the abdomen, and compresses placed upon any depending portion of the sac. By these means the attenuated diaphragm is restrained in its movements until it has recovered.

8. Bleeding and the discharge of bile into the sac can only be effectually prevented by promoting its contraction. The external bandages should be tightened, and injections of nitrate of silver and sulphate of zinc freely used. If there be no impediment to the discharge from the sac, it may be frequently washed out with cold water.

As far as my search has extended I find that the tumour in the above case is the largest for which operative proceedings have been undertaken with success; and since the treatment here advocated has been successful in the case of so large a tumour, it may, *à fortiori*, be employed for the destruction of those of smaller size. The withdrawal of a pint or two of fluid from smaller tumours has not unseldom proved fatal, and we cannot be sure that such treatment will result in the radical cure of the disease (see Table I).

Appended are synoptical tables of all those cases of hydatid tumours of the liver in which operative measures have been adopted that I have been able to glean from

various sources. In my remarks I have endeavoured to point out the advantages of securing the free discharge of the contents of the sac, and the dangers which follow its suppression. In the compilation of these tables I am chiefly indebted to four sources :

1. Mr. Cæsar Hawkins's papers "On Eycysted Tumours of the Liver" in the 18th vol. of the 'Medico-Chirurgical Transactions;'

2. M. Cruveilhier's Art. "Acéphalocystes," in vol. i, 'Dictionnaire de Médecine et de Chirurgie pratiques;'

3. M. C. Davaine's 'Traité des Entozoaires;'

4. Dr. Cobbold's 'Entozoa.'

From an analysis of these 100 cases it appears—1st, that of the 34 cases which were treated by a single puncture, evacuation of a portion or of the whole of the fluid, and immediate closure of the wound (see Table I), there were 11 *cures*, 13 *recoveries*—*i. e.* cases which were relieved by the operation, but which, since the tumour was not wholly removed or the result sufficiently certified, cannot be regarded as radical cures—and 10 *deaths*. Irritating injections were employed in Cases 27, 31, 33, and 34. One death (Case 13) was probably due to premature labour two or three weeks after the operation, the cyst being found collapsed; another (Case 15) was caused by rupture of another cyst into the lungs.

In the great majority of cases a single puncture is wholly ineffectual (see Tables II, III, IV, and V), and those who most strongly advocate this mode of treatment admit that it is only effectual where the tumour is of small capacity. Unless decided inflammatory action ensue within the cyst, which in a closed sac must always be attended with great danger, destruction of the parasite and radical cure of the disease can hardly be expected. In Cases 96 and 99 the parasite continued to maintain its existence for many years, notwithstanding the spontaneous rupture of the cyst and (in one) the operation performed upon it three years afterwards. The immediate dangers of puncture and closure of the wound are due—1st, to re-accumulation of fluid and

distension of the sac, resulting, when there are no adhesions with the abdominal wall, in overflow into the peritoneal cavity or elsewhere (*e. g.* Cases 32, 35, &c.); and 2nd, to excessive inflammatory action and the formation of putrid fluid within the sac. In Cases 48, 51, 54, 63, &c., clear watery fluid followed the first puncture, and fœtid pus the second. This second danger might probably be avoided by the plan adopted by Dr. Budd (see Case 1). In five, at least, of the fatal cases (Table I) death is attributable to re-accumulation of the fluid and its alteration and retention within the sac. In several cases of recovery re-accumulation of fluid took place, and in some this was accompanied by considerable pyrexia. But even when the death of the parasite follows capillary puncture, a simple cyst may remain, or the sac and its contents may undergo steatomatous or calcareous degeneration—results which cannot be considered as altogether satisfactory, when it may be proved that radical cure—*i. e.* the complete evacuation of the contents of the cyst, and the contraction and healing together of its walls—can be effected with less risk of a fatal termination. Case 56 shows how safely the means for the radical cure of the disease advocated in this paper may be employed against a small tumour.

Of 13 cases treated by successive puncture, with or without the injection of iodine (see Table II), there were 8 recoveries, 3 deaths, and 2 cases in which the operation was ineffectual. In two of the cases of recovery iodine injection was employed. In one case (47) ten punctures were required, and the last one followed by a strong iodine injection, and still the cure was incomplete. In two cases (35 and 38) five tapplings proved ineffectual, and death was doubtless due to the retention of the accumulated fluid, thus showing the necessity of maintaining a free communication externally.

Of the 30 cases in which the tumour was punctured once or oftener, and an external communication sooner or later preserved (see Table III), there were 23 cures, 18 at least of which may fairly be considered as radical, and

7 deaths, a proportion, notwithstanding the larger size of the tumours, less than that which followed simple puncture. The fatal result in five of these cases is directly attributable to re-accumulation of the discharge or to its retention and decomposition within the sac. The necessity for preserving an opening for the free discharge of fluid from the sac is well illustrated by Case 63.

Of 13 cases in which the tumour was directly incised (see Table IV), there were 6 cures and 7 deaths, a much larger proportion than resulted when the other modes of treatment were adopted; but it will be observed that some of these cases were taken for ordinary abscess, and no means employed to secure the flow of discharge from the sac. Even when a free incision has been made the discharge may most effectually be obstructed by one or more secondary cysts, floated against the outlet. Case 80 shows the advantage of making and maintaining a free opening.

Of 10 cases in which the tumour was opened by caustic potash, or by spontaneous absorption and rupture of the integuments and cyst-wall, there were 3 cures, 3 recoveries, in which the disease continued, and 4 deaths.

Attentive consideration of all the cases in which the tumour has been opened, whether by trocar, by the knife, or by caustic, will still further assure us, on the one hand, that the chances of recovery are in exact proportion to the care taken in the evacuation of the contents of the sac; and, on the other, that the fatal result is due, not to exhausting discharges from a large suppurating cavity, but to a want of care in evacuating the cyst membranes, and thus securing the discharge of fluid which, sooner or later, becomes putrid and poisons the blood. In only one of these 100 cases did the patient die from the exhaustion induced by prolonged discharge from the sac, and in this case the great cyst in the liver was found "full of hydatids and a purulent fluid." (See Case 82.) The advantages of a free opening and the speedy evacuation of the contents of the sac are well seen in Case 56.

In the treatment of hydatid tumours of the liver there

are two other points which deserve consideration, viz., the use of caustic potash and the injection of bile. Careful review of the cases in which the former has been employed, for the purpose of causing adhesion between the sac and the abdominal wall, or with the further object of opening the tumour, will serve to show that it is ineffectual for the first of these purposes (see Case 73), and that it possesses no advantage in effecting the second. Such painful treatment, therefore, may very well be altogether laid aside. As to the injection of bile into the sac, in order to cause inflammatory action within its interior and destruction of the entozoon, it is only necessary to observe that the cyst is so frequently found in direct communication either with one of the branches of the bile-ducts or with the gall-bladder itself as to lead one to the conclusion that the animal enters the liver by the common bile-duct, and that the hydatid tumour is sometimes, at least, formed as a diverticular growth from the walls of some one of its branches within the liver, or, perhaps, within the obstructed portion of the duct itself. In Vol. XVIII, p. 148, of these 'Transactions,' a case is recorded in which the common bile-duct was obstructed near its duodenal end by a hydatid cyst. Dr. Billing records a case ('Med. Gaz.,' vol. vii, p. 542) in which hydatids were found free in the gall-bladder. Jaundice and death has ensued from plugging of the common bile-duct by hydatids. (Dr. Nicholson, 'Med. Times and Gaz.,' 1861, vol. ii, p. 515.) See also Dr. Barlow's case, 'Guy's Hosp. Reports,' vol. vi, ser. 2, p. 185; a case by Dr. Bristowe, 'Path. Soc. Trans.,' 1858; and Case 4, Table I, of this paper.

SYNOPTICAL TABLES OF THOSE CASES OF HYDATID TUMOUR OF THE LIVER IN WHICH CURE WAS ATTEMPTED BY OPERATIVE PROCEEDINGS, INCLUDING THOSE IN WHICH SPONTANEOUS RUPTURE TOOK PLACE EXTERNALLY.

TABLE I.—*Cases in which the cyst was opened by a single puncture, the canula withdrawn, and the opening allowed to close by primary adhesion or otherwise.*

No.	Sex and age.	Situation of the tumour and treatment.	Nature and quantity of fluid withdrawn at the operation.	Progress.	Result and remarks.	Reference to author.
1	M. adult	Large tumour of liver. Tapped with ordinary sized trocar, and fluid withdrawn by a double acting stomach-pump syringe, fitted to the canula so as to prevent the entrance of air into the sac	7½ pints of hydatid fluid	Immediate relief. Liver still large from presence of hydatid sac, which, although emptied of its contents, was still of considerable size. No increase of girth two months after, and recovery of flesh and strength	Tumour reduced in size Recovery	G. Budd, M.D., 'Med. Times and Gazette,' 1860, p. 494.
2	M. 56	Large fluctuating tumour occupying the right hypochondriac, the epigastrium, and part of the left hypochondriac regions. Tapped with medium-sized trocar, and issue of—	5 pints of thick tapeworm-like fluid and broken down hydatids	When seen a fortnight after, there was no recumulation, and the patient was apparently quite well	The patient did not remain enough under treatment to assure us of its ultimate results Recovery	Mr. Holthouse, 'Med. Times and Gazette,' 1860, p. 344.

3	F. 30	Large tumour occupying the right hypochondrium. Puncture with small trocar. The cyst appeared only half emptied when the fluid ceased to run	1 to 2 pints	Inflammation of cyst with presence of air, then slow diminution of tumour, which could not be detected eight months after the operation	Radical cure	Mr. Jonathan Hutchinson, 'Lancet,' 1862, vol. ii, p. 389.
4	F. 15	The whole abdomen distended with a fluctuating tumour. Paracentesis abdominis, and evacuation of—	12½ pints of brownish and intensely bilious fluid	Temporary relief; recumulation of fluid, and death on the 23rd day. The cyst was found to be in communication with the gall-bladder	Death	Dr. Headlam Greenhow, 'Lancet,' 1862, vol. ii, p. 476; Dr. Murchison, 'Ed. Med. Journ.,' 1865, case xiv, p. 513.
5	M. 12	Fluctuating tumour in the right hypochondrium. Tapped with a flat trocar	1½ pint of colourless hydatid fluid	No constitutional symptoms. Recovered and left the hospital in a short time	Recovery	Sir B. Brodie; Mr. Caesar Hawkins, 'Med. Chir. Trans.,' vol. xviii, p. 118; Davaine, 'Traité des Entozoaires,' obs. cclviii, p. 569; Budd, G., 'Disease of Liver,' 3rd ed., p. 460.
6	F. 20	Fluctuating tumour of the abdomen connected with the liver. Tapped with medium-sized trocar	3 pints of clear fluid	No bad symptoms. The patient was seen after six years, and there was no return	Radical cure	Sir B. Brodie; Mr. Caesar Hawkins, 'Med. Chir. Trans.,' vol. xviii, p. 119; 'Med. Gaz.,' vol. i, p. 334; Davaine, obs. cclix, p. 569; Budd, 'Disease of Liver,' 3rd ed., p. 460.

No.	Sex and age.	Situation of the tumour and treatment.	Nature and quantity of fluid withdrawn at the operation.	Progress.	Result and remarks.	Reference to author.
7	M. 14	Tense elastic tumour of the right hypochondrium. Tapped with a medium-sized trocar—	Upwards of 3 pints of clear fluid	"The wound healed, and the disease probably did not return"	Recovery	Dr. Thompson; Mr. Caesar Hawkins, 'Med.-Chir. Trans.,' vol. xviii, p. 121; 'Med. Gaz.,' vol. i, p. 468.
8	F. adult	Tumour of liver. Tapping with medium-sized trocar, and removal of—	16 pints of watery fluid	"The result is not mentioned, so that it may probably be concluded that the case ended fatally"	Death (?)	Mr. Caesar Hawkins, 'Med.-Chir. Trans.,' vol. xviii, p. 121.
9	—	Tapping of the anterior of two distinct cysts in liver, and withdrawal of about—	8 pints of watery fluid	Death a few days afterwards. Another cyst containing about 12 pints of clear fluid was found between the liver and diaphragm	Death	Abercrombie, 'Dis. of Stom.,' p. 356; Mr. Caesar Hawkins, in 'Med.-Chir. Trans.,' vol. xviii, p. 123.
10	F. adult	Fluctuating tumour of the liver. Puncture with common trocar, and removal of—	A washhand-basinful of broken-down hydatids and thick yellowish fluid	Temporary relief, and death shortly afterwards	Death	Dr. Hewett; Mr. Caesar Hawkins, 'Med.-Chir. Trans.,' vol. xviii, p. 157.
11	Boy	Hydatid tumour of the liver. Tapped with common trocar, and—	12 pints of greenish water evacuated	Death two days after. Gall-bladder contained 8 pounds of bile, the duct was full of gall-stones. A large hydatid cyst of the spleen also existed	Death	'Edin. Essays and Obs.,' vol. ii, p. 352; 'Med.-Chir. Trans.,' vol. xviii, p. 152.

12	F. adult	A small tumour of right hypochondrium. Punctured with a very fine trocar, and evacuation of—	A small quantity of limpid, salty, non-albuminous fluid.} hospital perfectly cured	Cure	Récamier, 'Rev. Méd., 1825, t. i, p. 28; Cruveilhier, 'Diction. de Méd. et Chir.,' t. i, Art. "Acéphalocystes," p. 230; Barrier, Thèse, Paris, 1840, p. 57; Davaine, obs. cxlvii, p. 568.
13	F. adult, pregnant	Large hydatid cyst on right side of abdomen. Tapped with common trocar, and issue of—	A large quantity of purulent fluid and broken hydatids recovered somewhat from the operation, but after 2 or 3 weeks miscarried, sank, and died	Death	Dr. Cholmely; Dr. Bright, 'Abd. Tumours,' Syd. Soc. ed., p. 41; 'Guy's Hosp. Rep.,' vol. ii.
14	F. adult	Fluctuating tumour of epigastrium. Punctured with small trocar, and issue of—	4 pints of limpid colourless fluid	Recovery	Mr. Key; Dr. Bright, 'Abd. Tumours,' Syd. Soc. ed., p. 42; 'Guy's Hosp. Rep.,' vol. ii.
15	F. 30	Exploratory puncture below left false ribs, and issue of— 7 days after, a fresh puncture 2 months afterwards, a third puncture, to the right of the umbilicus	About 3 pints of clear fluid Both the second and the last puncture were "without result"	It is not stated whether any communication between the two tumours existed	Gonpil, in a Thesis by Cadet de Gassicourt, 'Sur la Rupture des Kystes hydatiques du Foie,' Paris, 1856, No. 50, p. 46; Davaine, obs. lxxvi, p. 447.

No.	Sex and age.	Situation of the tumour and treatment.	Nature and quantity of fluid withdrawn at the operation.	Progress.	Result and remarks.	Reference to author.
16	M. 42	Large tumour of right hypochondrium. Punctured with a very small exploratory trocar, and removal of about—	12 ounces of clear fluid	Syncope, rigors, bilious vomiting, collapse, and death 18 hours afterwards. The right lobe of the liver contained a cyst capable of holding about 9 pints	Death	J. Moissenet, 'Arch. Gén. de Méd.,' Fév., 1859, p. 144; Davaine, obs. cclvi, p. 567.
17	F. 19	Tumour of epigastrium. Punctured with capillary trocar	About a pint of clear water	No bad effects followed; the tumour disappeared, and had not returned 3 years after the operation	Radical cure	A. A. Boinet, 'Traitement des tumeurs hydatiques du Foie,' obs. v, p. 13, Paris, 1859; Davaine, obs. cclxii, p. 570.
18	F. 35	Tumour of right hypochondrium. Punctured with capillary trocar	About 4 ounces of clear fluid	Disappearance of tumour; no return after three years	Radical cure	Boinet, op. cit., obs. vi, p. 14; Davaine, obs. cclxiii, p. 570.
19	M. 34	Tumour in region of liver. Punctured by a very fine trocar; 7 days after, application of caustic potash	Limped hydatid fluid, quantity not stated	The tumour was reduplicated with fluid 3 days after the puncture. The caustic potash did not open the tumour. Nervous symptoms supervened	Death 25 days after the puncture	Récamier; Cruveilhier, 'Dict. de Méd. et de Chir.,' Art. "Acéphalocystes," t. i, p. 235; Davaine, obs. cclxxxiv, p. 590.
20	F. 33	Tumour occupying epigastrium and both hypochondria. Punctured in the epigastrium with a small	More than two pints of clear fluid	Progressed well; the remaining fluid was absorbed; there was no recumulation	Scarcely any remains of the tumour 3 months after recovery	Mr. Jonathan Hutchinson, 'Brit. Med. Jour.,' Feb. 20, 1864.

21	F. 36	exploratory trocar, and then, on the issue of a little clear fluid, with one the size of a No. 4 catheter	3 pints of clear fluid	At first febrile symptoms from peritonitis or inflammation of the cyst, which became distended and tympanic. Recovery in 6 weeks, and when seen, some months after, she was wholly free from the disease	Radical cure	Mr. Jonathan Hutchinson, 'Brit. Med. Jour.,' Feb. 20, 1864.
22	M. 46	'Tumour distending right hypochondrium and epigastrium. Punctured with trocar the size of a crow-quill at the outer margin of the rectus muscle, 2 inches below the margin of the ribs	6 pints of clear watery fluid	After 3 days, pain in region of the liver and repeated rigors. The febrile symptoms subsided, and a permanent cure resulted in 3 months' time	Cure	Frerichs, 'Dis. Liver,' Syd. Soc. ed., vol. ii, p. 268.
23 and 24	—	Professor Langenbeck communicated two cases of echinococi of the liver treated by simple puncture	—	No history, but both cases are said to have been cured	Cure Cure	Frerichs, loc. cit., p. 269.

No.	Sex and age.	Situation of the tumour and treatment.	Nature and quantity of fluid withdrawn at the operation.	Progress.	Result and remarks.	Reference to author.
25	F. 31	Tense fluid tumour of right hypochondrium. Tapped with trocar not quite so large as No. 1 catheter	About 12 ounces clear fluid	Three days after, fever and tympanic distension of abdomen. Left bed on the 10th day, and the hospital on the 32nd, when the tumour was soft, free from tension, and gradually diminishing. Six months afterwards had almost regained her ordinary health. There was still a fulness in the region of the tumour, but the lower margin of the liver was quite an inch above the umbilicus	Recovery, with a portion of the tumour remaining 6 months after	Charles Murchison, M.D., 'Edin. Med. Jour.,' Dec., 1865, Case 1, p. 503.
26	M. 35	Large tumour of right hypochondrium. Punctured with a grooved needle, and issue of— About 6 weeks after, tapped with ordinary trocar, and issue of—	1 ounce of limpid Punc-fluid 23 ounces of limpid fluid	No bad effects followed either operation. After the second, the cyst immediately collapsed and did not refill. 3 months after the operation he had recovered health and strength, and there was no external indication of the former cyst	Cure	Dr. William Budd, 'Brit. Med. Journ.,' 1859, p. 273.

27	M. 37	Fluctuating tumour of epigastrium and right hypochondrium. Puncture of right hypochondrium with capillary trocar and injection of a mixture composed of 3iss each tinct. of iodine and water, and 3ss of tinct. of potassium. The injection left in the cyst and the canula withdrawn	About 1½ pint of clear fluid	No pain followed the injection, but an hour afterwards symptoms of iodism and considerable general reaction, which continued for 5 days. Improvement from this time. A month after the operation the prominence of the hypochondrium was very little marked, and the general state satisfactory	Recovery, with a portion of the tumour remaining	Dr. Aran, 'Bull. Thérap.,' Sep., 1854, t. xlvii, p. 218; 'Arch. gén. de Méd.,' 5e série, t. iv, p. 477; Boinet, 'Iodothérapie,' Paris, 1855, p. 396; Davaine, obs. cclxxviii, p. 595.
28	M. adult?	Hydatid tumour of liver. Exploratory puncture, which gave issue to—	A transparent hydatid fluid	The tumour never reappeared, "and after several years the cure is not to be denied"	Recovery	M. Robert, 'Soc. de Chirurg.,' 18 Mars, 1857, p. 381; 'Gaz. des Hôp.,' 1857, p. 147; Davaine, op. cit., obs. cclxi, p. 570.
29	F. —	A similar case to No. 28, similarly treated	A transparent hydatid fluid	Progress similar	Recovery	M. Robert, 'Soc. de Chir.,' Mars, 1857, p. 381.
30	F. 18	Fluctuating tumour of the epigastric and right hypochondriac regions. Puncture with ordinary trocar, and removal of—	About 1½ pint of faintly greenish watery fluid tracted to half its original capacity. 7 weeks after, the contraction was still more evident, and she was discharged cured in about 3 months after the operation. The patient was seen three months afterwards, and a still further contraction was observable; she remained quite well and able-bodied	For some days prostration and tendency to sickness. In a week substantial recovery. On the 21st day the sac had contracted	Recovery, with some remains of the tumour	W. Brinton, M.D., 'Lancet,' 1862, vol. ii, p. 689.

No.	Sex and age.	Situation of the tumour and treatment.	Nature and quantity of fluid withdrawn at the operation.	Progress.	Result and remarks.	Reference to author.
31	F. 40	Growing tumour in left lobe of the liver the size of a young child's head. Puncture with an exploratory trocar two fingers' breadth below ribs in left hypochondrium, removal of nearly— Injection of 5ij of alcohol within the sac, and rapid withdrawal of the canula	2 pints of fluid	The tumour began to reappear 2 days after the operation. On the 3rd and 5th days vomiting; 6th, improvement, but appearance of jaundice; 8th, operation upon marked improvement; a tumour in the jaundice disappeared in 6 days, when the tumour had attained its original size. From the 18th day gradual diminution, and 3 months after, no appreciable traces of the tumour	Recovery This patient had previously undergone an operation upon	Adolphe Richard, 'Bull. gén. de Thérap.', 1855, t. xlviii, p. 414; Davaine, op. cit. obs. cxcvii, p. 606; 'Med.-Chir. Rev.', 1856.
32	F. 28	Fluctuating tumour of the liver. Application of caustic potash to the integuments covering its most prominent part; subsequent tapping of the cyst, and issue of—	2 pints of serous fluid	The day after the operation, jaundice, rigors, and symptoms of inflammation of the cyst. Peritonitis for 6 weeks, slight amelioration, afterwards persistent vomiting, and first death. A large hydatid cyst was found in the liver, the intestines were matted together, and the peritoneal	Death. It appears that no attempt was made to relieve the sac of its contents after the puncture, and that the hydatid fluid escaped into the peritoneal cavity.	Dr. Rogers, 'Brit. Med. Journ.', 1862, vol. i, p. 71.

33	M. 37	Fluctuating tumour in the liver. Puncture with a capillary trocar, and injection of a mixture of ʒiss each of tinct. iodine and water, with ʒss iodide of potassium	1½ pint of clear watery fluid	peritoneum was studded with cysts	vity, causing peritonitis, and extension of the disease.	Dr. Aran, 'Des injections iodées dans les Kystes hydatiques du foie Bull. Thérap.,' Sept., 1854, t. xlvii, p. 218; Davaine, obs. cclxxxviii, p. 595.
34	F. 21	A large deep-seated tumour in the right hypochondrium. Punctured with a fine trocar, and a mixture composed of ʒss each of extract of male fern, and solution of caustic potash, and ʒvj of water injected into the sac	ʒiv limpid colourless fluid, containing hooklets	At first febrile action, vomiting and purging. The tumour diminished, and she left the hospital remaining on the 23rd day.	Recovery, with a portion of the tumour remaining	Dr. Pavy, 'Med. and Chir. Trans.,' vol. xlix.

TABLE II.—Cases in which the cyst was punctured twice or oftener in succession, the opening being allowed to heal up directly after each operation.

A. Without injection. B. With injection of irritating fluid into the cyst.

No.	Sex and age.	Situation of the tumour and treatment.	Nature and quantity of fluid withdrawn at the operation.	Progress.	Result and remarks.	Reference to author.
35 A	M. 31	Tumour occupying right hypochondriac and epigastric regions, and extending to the left hypochondriac, umbilical, and right lateral regions, the edge being level with the umbilicus. Tapped 5 times within six weeks	About 8 pints of fluid removed at each operation	Relief after the tapping; reaccumulation of fluid. About two months after the first tapping and cessity of a fortnight after the last serving an external opening. The patient began to expectorate hydatid fluid in large quantities, and died the 3 days after. The hydatid cyst in the liver had tended to make an opening in the diaphragm and burst into the lungs	Death. No case could better illustrate the necessity of pre-tapping after the last serving an external opening. The patient began to expectorate hydatid fluid in large quantities, and died the 3 days after. The hydatid cyst in the liver had tended to make an opening in the diaphragm and burst into the lungs	Dr. S. O. Habershon, 'Guy's Hosp. Rep.,' 3rd series, vol. vi, p. 174.

36 A	M. 45	Fluctuating tumour of liver, the size of a child's head. Puncture with an exploratory trocar, and removal of the greater portion of the fluid. After 6 days a second puncture with a larger trocar	About a pint of limpid fluid with the punctures, and the first patient left the hospital cured. No fluid cured operation. The second removal followed the first	No bad effects followed the punctures, and the patient left the hospital cured	Recovery	Denarquay, A.A. Boinet, op. cit., p. 30; Davaine, obs. cclxiv, p. 571.
37 A	M. 20	Elastic fluctuating tumour of the epigastrium punctured twice, at an interval of 4 months, with a capillary trocar	About a pint the first operation, and the first tapping the cure appeared radical, but 4 months afterwards there was a distinct swelling. The second operation appeared to effect a complete cure	For some time after the first tapping the cure appeared radical, but 4 months afterwards there was a distinct swelling. The second operation appeared to effect a complete cure	Recovery	A. A. Boinet, op. cit., obs. viii, p. 18; Davaine, obs. cclxviii, p. 576.
38 A	M. 26	Swelling on the left side below the ribs, extending across the epigastrium to the right hypochondrium. Tapped 5 times within a month	Quantity not stated; the fluid was at first clear, but on the four subsequent occasions it was purulent	Reaccumulation of the fluid after each tapping. After the last "the liver rapidly began to swell," and the patient died 34 days after admission. Large hydatid cyst of the liver containing 4 secondary cysts, the sac were holding a pint of perfectly clear fluid	Death. This case shows how rapidly the fluid may reaccumulate, and it is to be observed that the contents of the sac were purulent when the second puncture was made.	Dr. Willshire and Mr. Canton, 'Lancet,' Sept. 1, 1860.

No.	Sex and age.	Situation of the tumour and treatment.	Nature and quantity of fluid withdrawn at the operation.	Progress.	Result and remarks.	Reference to author.
39, 40, and 41 A	—	Three cases of small hydatid tumours of the liver treated by successive puncture with an exploratory trocar	2 ounces of fluid withdrawn each time	In two cases a gradual diminution in the size of the cyst was to be traced for a long time afterwards. The third case was lost sight of	Recovery, with some portions of the tumour remaining	Frerichs, op. cit., vol. ii, p. 256.
42 and 43 A	—	Two cases of large hydatid tumour of the liver treated by successive puncture with an exploratory trocar	2 ounces of fluid withdrawn each time	Both cases quite unsuccessful; the cysts went on growing without interruption	Operation ineffectual	Frerichs, op. cit., vol. ii, p. 256.
44 A	M. 58	Fluctuating tumour of the epigastrium tapped with a small trocar in median line 2 inches above the umbilicus. 6 days after, tapped a second time	1½ pint of a turbid yellowish fluid the first time; a still larger quantity the second	No relief, and death 10 days after the first operation	Death	Mr. Moreton, of Tarvin, and Dr. Dobie, of Chester; Murchison, 'Edinb. Med. Journ.,' Dec., 1865, case vi, p. 509.
45 A	F. 34	Tumour the size of a child's head below the false ribs on the right side. Puncture with	After first puncture, 6 ounces of clear limpid fluid. After febrile disturbance, pain at the second, 6 to 8 in the epigastric region,	2 days after the second puncture, shivering, great	Recovery	M. Jobert, Barrier, 'Thèse cit.,' p. 83; Davaine, obs. cclvi, p. 573.

46	M. adult	very fine trocar, and re-ounces of a similar tenderness of the abdomen of the canula until fluid. After the third, domen. Recovery in 3 the morrow. 9 weeks 8 to 10 ounces of a days. 4 days after the afterwards, a second ex-turbid, brownish, pu-third puncture the wound ploratory puncture. 3 rulent fluid weeks later, a third was made across an eschar produced by caustic potash. A medium-sized trocar was used this time, and the canula retained until the morrow. An elastic catheter was introduced on two occasions, the same and the following day, but was not retained	A clear limpid fluid followed the first second puncture; a turbid one the second symptoms were set up and continued for several months. Ultimately, slow amelioration and cure	Some days after the second puncture the tumour reappeared, febrile symptoms were set up and continued for several months. Ultimately, slow amelioration and cure	Recovery	Chassignac, 'Gaz. des Hôpitaux,' 1857, p. 147, et 'Leçon cliniq.," p. 366, same journal; Davaine, obs. cxcx, p. 597; 'Bull. de la Soc. de Chirurg. de Paris,' 1856-7, p. 380.

No.	Sex and age.	Situation of the tumour and treatment.	Nature and quantity of fluid withdrawn at the operation.	Progress.	Result and remarks.	Reference to author.
47 B	M. 31	Tumour of the hypochondria, punctured with a capillary trocar, and issue of— 18 days after, a second puncture, and issue of— 32nd day a third puncture, and discharge of— 60th day a fourth puncture, and issue of about— 70th day a fifth puncture, and issue of— 84th day a sixth puncture, and discharge of— 95th day a seventh puncture, and issue of about— 115th day an eighth puncture, and— 123rd day a ninth puncture, and issue of about— 140th day a tenth and last puncture; evacuation of— Injection of cyst with a mixture of $\frac{3}{16}$ ss each tinct. iodine and water, and $\frac{3}{j}$ of iodide of potassium	From $\frac{3}{xij}$ to $\frac{3}{xiv}$ of clear fluid about $\frac{3}{x}$ of turbid liquid from $\frac{3}{iij}$ to $\frac{3}{iv}$ of slightly turbid fluid $\frac{3}{iv}$ fluid about $1\frac{1}{4}$ pint of red, dish-yellow turbid fluid $\frac{3}{j}$ of turbid fluid $1\frac{1}{4}$ pint of turbid fluid no result $\frac{3}{xij}$ of turbid sanguinolent fluid $\frac{3}{x}$ of similar fluid	No bad effects appeared to have followed the punctures, all of which were made with a capillary trocar. The first puncture was followed by a great diminution of the swelling and hepatic dulness, but the others had no apparent effect in diminishing the size of the tumour. No pain was caused by the injection, and 2 months afterwards the patient was discharged cured. At the time of his discharge the liver projected from $2\frac{1}{4}$ to 3 fingers' breadth beyond the margin of the ribs	Recovery, with a portion of the tumour remaining	Dr. Aran, 'Bull. thérapeut.', 1854, t. xlviii; Davaine, obs. cxcii, p. 599.

TABLE III.—Cases in which the tumour was punctured, once or oftener, and an external communication established at once or subsequently.

A. No injections employed. B. Injection of irritating fluids into the cyst.

No.	Sex and age.	Situation of the tumour and treatment.	Nature and quantity of fluid withdrawn at the operation.	Progress.	Result and remarks.	Reference to author.
48 A	M. 31	Tumour of epigastrium and right hypochondrium, encroaching 2 inches to the left of the median line, and below into the umbilical and right iliac regions. Tapped with exploratory trocar midway between umbilicus and cartilage of ninth rib. A month after, tapped again with a small trocar, and two days after with a full-sized trocar; an elastic catheter introduced, and the canula withdrawn	On first tapping, 38 ounces of clear and liquid. On the second, 10 ounces of fetid pus. On the third, 39 ounces of fetid pus and broken-down hydatids	Discharge of fetid pus and membranous flakes of hydatids for 3 months. Gradual contraction and disappearance of the sac; healing of the fistulous opening, and cure	Radical cure. It is to be observed that fetid pus served on Med. Gaz., vol. xiii, p. 1101, 1849; 'Bull. de Thérap.', 1848, t. xxxv, p. 331; 'Arch. gén. de Méd. de Paris,' Juillet, 1849, p. 346; Davaine, obs. cclxvii, p. 576.	Dr. Owen Rees, 'Guy's Hosp. Rep., ser. ii, vol. vi, p. 17, 1848-9; 'Lond. Med. Gaz., vol. xiii, p. 1101, 1849; 'Bull. de Thérap.', 1848, t. xxxv, p. 331; 'Arch. gén. de Méd. de Paris,' Juillet, 1849, p. 346; Davaine, obs. cclxvii, p. 576.

No.	Sex and age.	Situation of the tumour and treatment.	Nature and quantity of fluid withdrawn at the operation.	Progress.	Result and remarks.	Reference to author.
49	M. 36	Tumour commencing an inch above the umbilicus, and extending over the abdomen. Punctured with a small trocar several times within 11 weeks. Afterwards a small quantity of fluid was drawn off every third day. Ultimately the opening was enlarged with a bistoury, and a large canula introduced	$\frac{1}{2}$ a pint of clear fluid followed the first puncture. 4 pints of similar fluid followed the second. About the same quantities of turbid brownish fluid followed the subsequent punctures	The patient progressed favorably, the discharge ultimately became quite purulent and fetid, but it was only after enlarging the opening that the evacuation of the broken-down cysts was effected, a fistulous opening was established, and the sac slowly contracted	Radical cure	Dr. Babington and Mr. Cock; Dr. Habershon, 'Guy's Hosp. Rep.,' vol. vi, 3rd series, p. 179; 'Med. Times and Gaz.,' 1856.
50	M. 15	Prominent fluctuating tumour, the size of a fist, on the right side of the abdomen. Puncture through eschar formed the previous day by caustic potash. An elastic catheter placed in the wound until the morrow	$\frac{1}{2}$ a pint of limpid fluid followed the puncture, and the catheter gave issue to a glassful more	Escape of fragments of an hydatid cyst through the wound for a few days. Slight erysipelas of the wound. Subsequently bilious vomiting and fever for some hours. Gradual subsidence of the tumour, cicatrization of the wound, and cure in about three months	No particular care appears to have been taken to keep the wound open, and it closed again several times before it finally healed. The constitutional disturbance was doubtless due to the tem-	MM. Richerand and Jobert; Barrier, 'De la Tumeur hydatique du Foie,' Thèse, Paris, 1840, p. 80; Davaine, op. cit., obs. cclxv, p. 572.

51 A	F. 19	Large tumour of the epigastrium, punctured in 4 ounces of clear fluid. The centre with a capillary. On the second, 6 to 8 trocars. 23 days after tapping pus and attempt to close the wound	On first puncture, 4 ounces of clear fluid. On the second, 6 to 8 ounces of thick fetid pus. 23 days after tapping pus and attempt to close the wound	4 days after the first operation, rigors and distension of the tumour, necessity for a slight tympanites. After free external the second tapping, the opening is well tumour again became dis- seen. It will be tended, and caused the further observed wound to give way on the that the effect of 5th day, when there was the first puncture a discharge of air and there was to continue fetid pus. The discharge vert a clear fluid continued for the next 24 within the sac months, at the end of into a thick which time the patient fetid pus was gaining flesh and strength, and the opening was discharging still	Radical cure? In this case the Sep. 1, 1860, p. 210.	Dr. Garrod, 'Lancet,' Sep. 1, 1860, p. 210.
52 A	F. 22	Inferior part of right side of the chest distended of glutinous fluid escaped on puncture contained in the pleuritic cavity. Tapped with ordinary trocar	A small quantity of glutinous fluid escaped on puncture	Discharge of similar fluid and hydatid membranes continued; and on the 33rd day after tapping, a large quantity of pus and hydatid membranes came away. She was discharged cured on the 43rd day?	Radical cure? Observation of the case not sufficient to propose	Dr. Goolden, 'Lancet,' Sep. 1, 1860, p. 210.

No.	Sex and age.	Situation of the tumour and treatment.	Nature and quantity of fluid withdrawn at the operation.	Progress.	Result and remarks.	Reference to author.
53 A	M. 59	Tumour in epigastric and hypochondriac regions. Punctured with a small trocar; 7 days after, punctured again with a much larger trocar; the canula withdrawn, and a flexible tube the size of the trocar retained in the wound. The tube became obstructed, and was removed 5 days after, and the wound kept open by coiled lint	9 drachms of slightly turbid fluid followed the first puncture, and a little clear fluid and hydatids and the second	After the insertion of the tube, only a little discharge oozed from the sac by the side of the catheter, and there was some constitutional disturbance. The tube was blocked up by a collapsed hydatid. After the removal of the tube, broken hydatids and fetid pus continued to be discharged for about 5 weeks, gradually decreasing in quantity. On the 79th day from the first puncture, he was in perfect health, and a cord-like cicatrix occupied the place of the former opening	Radical cure	Dr. Owen Rees, 'Lond. Med. Gaz.,' vol. xliii, p. 1101, 1849.
54 A	M. 42	Enormous tumour occupying the epigastric and hypochondriac regions, and reaching as low as the umbilicus. Tapping glass, with a trocar; and 10 days	At the first operation, only a little pus was obtained by means of a cupping glass. At the second operation, offensive	In the interval between the first and second operations, the tumour increased in size, and the presence of air was manifest. Inflammatory symptoms, trid fever ap-	The cyst in this case was not permanently relieved of its contents. Putrid fever ap-	Mr. Hilton, 'Lancet,' 1851, vol. i, p. 353.

55	A	M. 12	after, an incision, $\frac{3}{4}$ inch gas and a number of toms supervened. After the long, was made down to hydatid cysts were the second puncture, hy- been set up, re- the cyst, and a trocar and removed; and 5 days, datids were freely dis- sulting in death canula then plunged into afterwards, $1\frac{1}{2}$ pint of charged with purulent it. A plug of lint was purulent fluid was discharged	8 pounds of clear fluid	Fluctuating tumour in the hepatic region. In- cision and evacuation of— The lining membrane of the cyst was withdrawn by gentle traction	Inflammation and sup- puration of the cyst fol- lowed the complete evis- ceration of its contents, rapid and com- plete evacuation of the cyst	Dr. A. Monro, 'Morbidity of the Gullet,' Prov. Med. and Surg. Journ., vol. xiv, 1850, p. 394.	Radical cure promptly effected by the 'Prov. Med. and Surg. Journ.,' vol. xiv, 1850, p. 394.
56	A	M. 56	Small hydatid tumour of the liver, occupying the epigastric and right hypochondriac regions. Exploratory puncture with grooved needle. The following day, a crucial incision over the most prominent part of the tumour; and after the flaps were dissected back, caus-	A little viscid serum containing book-lets was obtained by the grooved needle. After the opening was dilated, from ʒiv to ʒviii cure in the case of sanguinolent viscous matter, crowded with hydatid cysts, were discharged daily for 10 weeks, during which time the patient remained in a very of the cyst as	Considerable pain and restlessness followed the introduction of the trocar. After the opening was dilated, from ʒiv to ʒviii cure in the case of sanguinolent viscous matter, crowded with hydatid cysts, were discharged daily for 10 weeks, during which time the patient remained in a very of the cyst as	This is a capital illustration of the progress of the disease, and shows the necessity of a small tu- mour. It also shows the necessity of a small tu- mour. It also shows the necessity of a small tu- mour.	J. Hughes Bennett, M.D., 'Princ. and Prac. of Medicine,' 4th Edit., p. 522.	Radical cure promptly effected by the 'Prov. Med. and Surg. Journ.,' vol. xiv, 1850, p. 394.

No.	Sex and age.	Situation of the tumour and treatment.	Nature and quantity of fluid withdrawn at the operation.	Progress.	Result and remarks.	Reference to author.
57 A	M. 10	tic potash was applied to the subjacent tissue. On the 9th day, a second application of the caustic. On the 11th, a trocar and canula, $\frac{3}{4}$rd inch in diameter, were passed through the slough into the tumour, and a scooped director used to evacuate its contents. 16th day, dilatation of the wound, and the insertion of tents of lint to preserve the opening Hydatid tumour of the liver tapped with a large trocar, and the opening preserved	Discharge of a large quantity of turbid fluid and cyst-wall	weak condition. A tough membrane was subsequently evacuated, and henceforward the discharge rapidly decreased, and in about 14 weeks from the time of puncture with the trocar the wound was quite healed. A month afterwards there was no trace of any hepatic tumour, and the patient was in perfect health Portions of cyst were subsequently discharged in larger quantities, and the patient made a good recovery	soon as possible. After the discharge of the parent cyst, there was a speedy recovery. Radical cure Cure, probably radical	Dr. Hjalalin; Dr. Leared, 'Path. Soc. Trans., vol. xiv, p. 176.
58 A	F. 31	Fluctuating tumour commencing in the right hypochondrium, and extending over the whole of	On first tapping, only a few ounces of pus; a few hours after, 16 pints of purulent	Immediate relief, rapid recovery of health and strength. The discharge gradually decreased to a	If we accept the patient's statement that the tumour com-	A. Copeland Hutchison, 'Practical Observations in Surgery,' London, 1816, p. 145.

59	M. adult	the abdomen; most pro- fluid and hundreds of few drachms daily. Moremened in the right of the ensiform car- subsequently the daily operation, nearly 2 quarts, drum, and take tilage. Puncture with discharge of large of healthy pus, free from into account the large-sized trocar, and re- quantities of fluid Intention of the canula, through the canula to push away or rupture the secondary cysts. Canula stopped with a cork during daily intervals. Removal of the canula after five days, and substitution of a sponge-tent	6 pints of sero- purulent fluid and small hydatids	Pain and febrile symp- toms followed the 1st ope- ration, and on the 7th day the tumour had regained its previous size. The from obstruction same occurred after the of the discharge, second puncture, and, al- and the advan- though one puncture at tages of making least remained open, after free openings the 18th day it was neces- into the sac sary to make the other openings in order to give vent to accumulating fluid, and it was only after the	This case illus- trates very com- pletely the dan- 1823, vol. xix, p. 228. Radical cure	H. C. Sherwin, 'Edin- Med. and Surg. Journal,' 1823, vol. xix, p. 228.

No.	Sex and age.	Situation of the tumour and treatment.	Nature and quantity of fluid withdrawn at the operation.	Progress.	Result and remarks.	Reference to author.
		On the 7th day, second puncture an inch higher with a very large round trocar, and the withdrawal of— This opening was also healed On the 13th day, trocar introduced a third time, and the canula retained and plugged On the 24th day a fourth tapping, and on the 32nd day a fifth tapping, were resorted to, and— On the 39th day, an incision an inch long was made between the 9th and 10th ribs, behind their angles, and a large round trocar introduced in a direction pointing to the scrobiculus cordis. The canula was retained for 10 days, a thick piece of wax bougie was then substituted	10 pints of sero-purulent fluid and hydatids 8 pints of purulent fluid and hydatids were obtained by the third operation Several pints of purulent fluid removed 6 pints of very fetid pus and some larger hydatids, and towards the end a little bile, were obtained by this last puncture	6th operation that this was sufficiently secured. The discharge then rapidly decreased, the sac contracted, and the patient recovered his health and strength and returned to his work 3 months after the operation. 7 weeks after this date, the sac was entirely obliterated, and nothing remained but a slight fistulous opening		

60 A	M. 38	A large circumscribed fluctuating tumour in the right hypochondrium, extending to the umbilical region. Previously to the operation, jaundice, great pain in the tumour, shivering, and increase of the tumour. Puncture, and the application of poultices to the punctured part	A gallon of thin, purulent, offensive matter and hydatids followed the drawal of the instrument	Hydatid membranes and a yellowish green tenacious fluid continued to be discharged. Ten-secure the evacuation of the tumour increased, and the contents of the patient died on the 26th cyst, and the day after the operation, fatal result is An immense cyst, full of attributable to numerous large hydatids, retention of the was found in the right discharge Death.	No means appear to have been adopted to 1832, p. 756.	John Elliotson, M.D., F.R.S., 'Lancet,' vol. i, 1832, p. 756.
61 A	F. 20	Painful tumour in the right hypochondrium, indistinctly fluctuant. Puncture, and withdrawal of—	10 ounces of bloody pus and hydatids	Relief. The puncture was healed on the 3rd day, and the tumour re-filled. On the 11th day the integuments gave way, and a considerable quantity of matter was discharged. On the 17th day the opening was healed and the cyst contracted	In this case the accumulated re-fluid produced sufficient distension to effect spontaneous rupture, and the safety of the patient was thus secured. Cure	Dr. Cholmeley, 'Lancet,' 1826.
62 A	F. 62	Tumour of the liver extending from the right lumbar region across the abdomen to 2 inches left of the umbilicus, and upwards as far as the 7th rib. Puncture, and the opening preserved by lint tents and simple dressing	4 pints and a few ounces of offensive pus and hydatids followed the withdrawal of the instrument	At first moderate febrile action, but on the 4th day it had ceased, and the patient was in a very favorable condition, and there was great discharge from the wound	The history stops short on the 4th day; but from the plan of treatment adopted and the freedom of the discharge, it is probable that the patient recovered	Dr. Roberts, 'Lancet,' 1834, vol. i, p. 189.

No.	Sex and age.	Situation of the tumour and treatment.	Nature and quantity of fluid withdrawn at the operation.	Progress.	Result and remarks.	Reference to author.
63 A	M. 25	Fluctuating tumour of the right hypochondrium. Two applications of Vienna paste; puncture through the eschar with an ordinary trocar. A second puncture 3 months afterwards; closure and cicatrization of the opening. Subsequent rupture through the cicatrix, and establishment of a fistulous opening	5½ pints of clear fluid followed the first puncture. Pus and broken-down hydatids followed the second	Reaccumulation took place after both punctures. The spontaneous rupture was followed by the discharge of fetid fluid and hydatids. Exhaustion and death 5 months from the time of the first operation. A hydatid cyst, the size of an adult's head, and a second smaller one, were found in the liver	Even the ruptured opening was insufficient to secure the evacuation of the contents of the cyst, and no means appear to have been used to evacuate the secondary cysts. Death	Cruveilhier, 'Gaz. des Hôpit.,' 1842, 2e série, t. iv, p. 317.
64 A	M. 38	Tumour of epigastrium, exploratory puncture. Two applications of caustic potash, and eight days after the last one incision through the eschar. A month afterwards a fresh puncture, and the opening preserved by means of a sponge-tent	A little clear fluid followed the exploratory puncture, and half a pint of turbid serosity the incision. On making the last opening, a most fetid fluid and hydatids escaped	On the 22nd day the tumour had nearly disappeared, but the patient continued in an unsatisfactory state. On the 45th day the belly was still a little painful and distended. The last opening produced immediate relief; a free outlet was maintained, and the patient progressed satisfactorily, and the cyst was diminishing every day	Radical cure	Récamier, 'Lond. Med. Gaz.,' vol. ii, p. 573, 1828, 'La Clinique,' May 6, 1828 (?); 'Lancet,' 1827-8, vol. ii, p. 327 (?).

65 A	F. adult	A case of supposed hydatrothorax. Puncture between the lower false ribs of the right side, and a tent placed in the wound	Fluid and hydatids	The patient died very quickly. The thoracic organs were healthy, but the liver was in great part destroyed by a hydatid cyst	Death	Ruysch, 'Obs. Anat. Chirurg.,' obs. lkv, p. 61, cit. par Lassus. Mén., cit., obs. v; Cruveilhier, op. cit., p. 229; Davaine, obs. cclxxv, p. 581.
66 B	M. 30	Circumscribed tumour of the right hypochondrium. Exploratory puncture, and— 6 applications of caustic potash; incision of the eschar, and evacuation of— A tent placed in the wound. Occasional introduction of an elastic catheter, and washing-out of the cyst. Injection of tincture of iodine	Issue of about 4 ounces of watery fluid. 1½ pint of turbid, fetid, yellowish serosity	After the puncture, there were vomiting, fever and exquisite sensibility of the belly. These symptoms abated, but returned with great severity after the 6th application of caustic potash, and the tumour increased in size. The patient remained in a precarious state after the incision, but 3 months subsequently he had regained his flesh and strength. The fistula gradually closed, and the patient was cured in about 4 months	Radical cure	Robert, 'Bull. gén. de Thérap.,' Paris, 1843, t. xxv, p. 379; Davaine, obs. cccxiv, p. 501.
67 B	F. 8½	Tumour of epigastrium, capillary puncture, and removal of about— 16th day, a second puncture, with medium-sized trocar; issue of— Injection of iodine; soon	2 pints of clear hydatid fluid 2 pints of greenish purulent matter 2 pints of fetid relief, but as the sac filled it was necessary	No bad effects followed until the 10th day, when, the sac having refilled, severe febrile symptoms supervened. The second operation gave complete relief, but as the sac filled it was necessary	Radical cure.	Dr. Boinet, 'Gaz. hebdomad. de Méd. et de Chir.,' sér. 2, 1864, tom. i., p. 86. The author records another successful case similarly treated in the 'Gaz. Méd.,' 1860.

No.	Sex and age.	Situation of the tumour and treatment.	Nature and quantity of fluid withdrawn at the operation.	Progress.	Result and remarks.	Reference to author.
68	M. 33	after, application of caustic potash and chloride of zinc. 42nd day, puncture with a very large trocar, and evacuation of— Insertion and retention of a large elastic catheter. 49th day, another large puncture made into the cyst and drainage tubes kept in both openings. 64th day, incision into the cyst, and withdrawal of its contents	greenish pus and broken-down hydatids	again the dangerous symptoms reappeared. As soon as the external opening was freely established all went on well, and after the removal of the cyst membranes by incision the sac speedily contracted, and was completely obliterated on the 110th day	to establish a free external opening	Mesnet and Boinet, 'Revue méd.,' 15 Février, 1853; 'Bull. Soc. Chir.,' 1852; Boinet, op. cit., p. 387; Cadet de Gassicourt, Thèse cit., p. 76; Davaine, obs. cxcxi, p. 597.
B		Large tumour of right hypochondrium raising the walls of the chest, punctured with a common trocar, and issue of— The flow being arrested, the sac was opened at once with a bistoury and a dilator introduced. The sac was repeatedly injected with iodine	Several spoonfuls of pus and a hydatid. On making the incision, about 2 pints of pus and hydatid membranes were evacuated with another cyst which occupied nearly the whole of the right lobe of the liver and extended through the diaphragm to the right lung	The patient went on satisfactorily for nearly a month after the operation. Subsequently he had shivering, fever and fetid stools, and died on the 42nd day. The punctured cyst contained yellowish brown liquid and two hydatids. It communicated with another cyst which occupied nearly the whole of the right lobe of the liver and extended through the diaphragm to the right lung	No mention made of any continuance of the operation; but a month after the operation an hydatid was withdrawn from the cyst. The punctured datid was with-obs. cxcxi, p. 597.	

69 B	F. 27	Hard tumour on right side of abdomen. Exploratory puncture and evacuation of— 2 days after, application of Vienna paste nearly midway between false ribs and crest of ileum. Repeated until the cyst burst spontaneously. This occurred on the 22nd day Injection of tinct. iodine through an elastic catheter. On the 24th day, removal of catheter and enlargement of the opening by a bistoury, and repeated injection of iodine	4 pints of clear fluid. Discharge of 4 pints of a reddish fluid	Rigors and rather severe fever, with bilious vomiting, came on after the first injection of iodine, but soon ceased. The contents of the cyst were gradually evacuated. The patient's state continued to improve until rather more than 3 months from the first puncture, and she was discharged cured	Radical cure	M. Demarquay, 'Gaz. Méd. de Paris,' Sept. 16, 1865; 'Brit. Med. Journ.,' Dec. 16, 1865, p. 642.
70 B	F. 16	Fluctuating tumour of right hypochondrium. Several applications of caustic potash; exploratory puncture, and 12 days after, puncture with a large trocar. Next day dilatation of the wound with prepared sponge. Injection of iodine on the 20th day. On the 28th, solution of nitrate of lead; and subsequently chlorinated water twice a day	2 spoonfuls of clear fluid followed the first of puncture, 4½ pints of clear limpid fluid the second. Next day, opening discharged only a few drops of serosity, and at the end of 4½ months she was perfectly cured and the fistula had healed	Free discharge, at first of bilious fluid. On the 12th day it was purulent. In 2½ months the fistulous opening discharged only a few drops of serosity, and at the end of 4½ months she was perfectly cured and the fistula had healed	Radical cure	Laboulbène; Davaine, obs. cclxxxiii, p. 588.

No.	Sex and age.	Situation of the tumour and treatment	Nature and quantity of fluid withdrawn at the operation.	Progress.	Result and remarks.	Reference to author.
71 B	M. 36	A tumour of considerable size in the hepatic region. Application of caustic potash until it was assumed that adhesions had taken place, then tapping with ordinary trocar. The sac washed out with water, injected with iodine, and an elastic catheter inserted and kept in the wound. Iodine and chlorine injections frequently used	1½ pint of pus	The patient progressed well. Pus and débris of hydatids continued to be discharged. On the 11th day, bile was mixed with the pus. From the 13th to the 19th day, the discharge consisted of pure bile. Left the hospital on the 60th day, the fistulous orifice discharging a little pus. 5 months afterwards, the fistula was closed, the wound cicatrized, and the general health excellent	Radical cure	M. Nélaton; Cadet de Gassicourt, These cit., obs. iii, p. 18; Davaine, obs. cclxxxix, p. 596.
72 B	F. 34	Fluctuating tumour of right hypochondrium. Three exploratory punctures at intervals of 20 days, and evacuation of about— Afterwards, 3 applications of Vienna paste; and on the 38th day after the last puncture, an opening	8 pints altogether of fluid; clear and albuminous on two occasions, purulent on the last. 3½ pints of pus followed the fourth puncture	No bad symptoms followed the first two punctures; the third induced shivering and fever. Ca-pillary punctures proved insufficient to produce contraction of the tumour. Some days before the wound was enlarged, febrile symptoms appear.	Radical cure. This case shows the insufficiency of a tumour of the removal of a tumour of moderate size	Demarquay, 'Gaz. des Hôpitaux,' 19 Février, 1859, p. 82; Davaine, obs. cxcv, p. 603.

73 B	F. 27	was made with a bistoury, an elastic catheter introduced, and injections of iodine and perchloride of iron employed from time to time. On the 62nd day the opening was enlarged by a bistoury; withdrawal of the catheter on the 86th day	5½ pints of very fetid pus followed the second puncture; but no fluid followed the last application, puncture with cataract needle. 3 days after, puncture with ordinary trocar through the eschar. The canula was withdrawn, but replaced again on the 5th day; but being too fine to allow a free flow, another puncture was made on the 8th day. Iodine injections were used on the 5th and subsequent days	ied. This operation was followed by the ejection of pus and débris of hydatids, and an abundant flow of blood. 6 months after the first puncture, the patient was in a satisfactory state, and may be considered as cured	Severe diaphragmatic peritonitis followed the exploratory puncture. No effect followed the intervals of 325, Paris, 1856; Davaine, obs. i, p. 604. Death. 3 ap- plications of Vienna paste at the intervals of 3 days failed to produce adhesion of the sac	F. Dolbeau, 'Sur les Vi- grands Kysts du Foie,' Thèse, No. 113, obs. i, p. 325, Paris, 1856; Davaine, obs. cxcvi, p. 604.

No.	Sex and age.	Situation of the tumour and treatment.	Nature and quantity of fluid withdrawn at the operation.	Progress.	Result and remarks.	Reference to author.
74 B	F. 53	Fluctuating tumour distending the lower ribs of the right side. Application of Vienna paste between the 8th and 9th ribs; 2 days afterwards, puncture through the slough with an exploratory trocar; 3 days after this, puncture with a large trocar, injection of bile morning and evening, and retention of the canula	After exploratory puncture, issue of clear water and minute hydatids. After day there was diarrhoea between the 8th and 9th ribs; 2 days afterwards, puncture through the slough with an exploratory trocar; 3 days after this, puncture with a large trocar, injection of bile morning and evening, and retention of the canula	The patient did well up to the 8th day, when the discharge failed; the next day there was diarrhoea and some fetid discharge from the sac; fever and pneumonia followed, and the patient died on the 36th day. The cyst contained fluid and hydatids; there was some pneumonia of the left lung, calculi and distension of the gall-bladder, and ulcerations of the duodenum and small intestine	Death	Auguste Voisin, 'Bull. Soc. Anat.,' Paris, 1857, ann. xxvii, p. 132; Davaine, obs. cccxviii, p. 608.
75 B	M. 29	Very large fluctuating tumour extending from the epigastrium over the whole of the abdomen and into the lower part of the chest, displacing the lungs and heart. Puncture with No. 11 trocar; maintenance of the opening by the	Discharge of 19½ pints of fluid, the greater part bright and clear as water, and the latter portions tinged yellow and turbid from the pre-down cyst-wall. Except absence of bile: and also a number of minute discharge was very profuse	Great relief and favorable progress, occasionally interrupted by distress and febrile symptoms due to retention of the discharge and charge by the broken-down cyst-wall. Excepting at these times, the	Radical cure	Synopsis of the case related in this paper. 'Med. Times and Gaz.,' 1866, vol. i, p. 482; 'Lancet,' 1866, vol. i, p. 538; 'Gaz. Méd. de Lyon,' 1866.

76	F. 26	canula and by elastic catheters. Gradual dilatation of the passage by means of elastic catheters. Injection of a large quantity of iodine, and a daily cleansing of the cyst with a solution of zinc and creasote water	and amounted to many gallons. For 21 weeks from the day of tapping, the discharge contained a considerable quantity of bile. On the 51st day there was hæmorrhage from the liver. On the 75th day the whole of the contents of the cyst were evacuated, and from this time the discharge decreased and cyst contracted, and ultimately disappeared	Radical cure	M. Leudet, 'L'Union Médicale,' 1859, No. 90, p. 202.
		Fluctuating tumour of the upper part of the abdomen. Destruction of the integuments by caustic potash, and puncture of cyst with a capillary trocar; closure of the orifice and reopening by means of a bistoury, and insertion of an elastic catheter; frequent injection of iodine and alcohol. Catheter retained for four months	Intense general symptoms after the second operation. Subsequent evacuation of a great number of fragments of hydatid cysts. Closure of the fistula, and recovery of health in about 5 months from the first operation		

No.	Sex and age.	Situation of the tumour and treatment.	Nature and quantity of fluid withdrawn at the operation.	Progress.	Result and remarks.	Reference to author.
77	F. 17	Hydatid tumour of the right hypochondrium. Incision down to the peritoneum, 7 days after, 3iv puncture with a flat trocar, and the wound closed. second A second puncture on the 10th day, and the cannula retained	2 pints of watery colourless fluid by the first puncture. 3iv yellowish-brown turbid fluid by the second	Severe febrile symptoms followed the first puncture. Purulent fluid and hydatid membrane continued to be discharged. The tumour gradually subsided, and the fistula closed	Cure. The patient died 5 months after of phthisis	Ried and Brehme, 'Deutsche Klinik,' No. 59, p. 377 and 386.

TABLE IV.—Cases in which the tumour was directly incised without previous puncture, or (excepting Case 89) the application of caustic potash, and an external communication preserved.

No.	Sex and age.	Situation of the tumour and treatment.	Nature and quantity of fluid withdrawn at the operation.	Progress.	Result and remarks.	Reference to author.
78	M. 28	Fluctuating tumour of the posterior and inferior part of the right side of the chest. Incision about 1½ inch long between the third and fourth false ribs, in the centre of the tumour	At first limpid fluid; afterwards a great quantity of yellow pus, and for the next 12 days several hydatids with a great quantity of pus	15 days after the operation, precordial anxiety, painful cough and expectation of hydatids. This continued to the 43rd day. On the 72nd day, the wound was reduced to a small fistulous opening, with only a slight suppuration of a healthy character, and the condition of the patient was improving from day to day	Radical cure	Fréteau, 'Journ. gén. de Sédillot,' t. xlii, p. 121; Cruveilhier, 'Dictionnaire de Méd. et de Chir. pratique,' t. i, art. 'Acépl.,' p. 249 (extract); Davaine, <i>op. cit.</i> , obs. xxxiv, p. 415.
79	M. 37	Very large fluctuating tumour of the epigastrium and lower part of abdomen. Opened by incision with a scalpel 3 inches below the ensiform cartilage. The discharge being obstructed two days afterwards, the canula was again introduced into the wound	At first about 9 pints of fluid were evacuated. On reintroduction of the canula, 4 pints more	At first, relief; but, two days after the operation, return of the dyspnoea. After the reintroduction of the canula, there was a constant oozing until death, which took place 9 or 10 days afterwards. The cyst was found to contain nearly 14 pints of viscid bilious fluid in the peritoneal cavity	Death. The result was clearly due to the reintroduction of the fluid from the distended sac into the peritoneal cavity	Dr. Hastings, 'Midland Medical Reporter,' Aug. 1829; 'Trans. Med. and Chir. Soc.,' vol. xxiii.

No.	Sex and age.	Situation of the tumour and treatment.	Nature and quantity of fluid withdrawn at the operation.	Progress.	Result and remarks.	Reference to author.
80	M. 36	Fluid tumour of the epigastrium opened by an incision 2 inches long between the umbilicus and ensiform cartilage. A pledget of lint was placed in the wound	At first a quantity of hydatids and fluid equal to 12 pints	Progressed well; fluid and cyst-wall continually escaping. The fluid became extremely fetid, and a month after the operation the patient was very weak. During the second month there was about a pint and a half of discharge each day. Diminished to 2 ounces during the next month. At the end of 5 months a slight discharge. Three years after the operation, the patient was in fair health	Radical cure	J. Russel, 'Duh. Journ. Méd.,' Nov. 1837; 'Archives gén. de Méd.,' 1838, t. i, p. 106; Davaine, obs. cclxxi, p. 578.
81	M. adult	Supposed abscess on the right side of the abdomen. Incision	Discharge of a great number of vesicles, it is assumed, with a yellowish fluid	The discharge of hydatid cysts continued for several days. The patient was perfectly cured	Cure	Rivière; Cruveilhier, 'Diction. de Méd. et de Chir. Prat.,' t. i, Art. "Acéphalocystes," p. 223.
82	M. adult	Tumour, right hypochondrium, increasing for 6 years, and causing much distress. It was taken for an abscess and incised	A great quantity of hydatids, accompanied by a thick viscid matter	More than 300 hydatids came away at different times. The patient was worn out, and died after a year. The great cyst in the liver was full of hydatids and a purulent fluid	Death	Camerarius; Cruveilhier, op. cit., p. 224; 'Boneti Sepulchretum,' p. 1532.

83	M. adult	Tumour in the hepatic region mistaken for an abscess, opened by incision	Discharge of hydatids and fluid immediately, and for 5 days after	The patient grew weaker and weaker, and died on the 5th day	Death	Panaroli, obs. xvi, cit. par Lassus, 'Rech. et Obs. sur l'Hydropisie enkystées du Foie,' Davaine, obs. cclxxiv, p. 581; Cruveilhier, op. cit., p. 225. Sue; Lassus, Mém. cit., obs. ix; Davaine, obs. cclxxvi, p. 581; Cruveilhier, op. cit., p. 225.
84	M. adult	Tumour of the epigastrium taken for an abscess. Incision	About 2 pints of limpid serosity	Watery fluid continued to be discharged, and there was constant hiccup and vomiting, and death on the 3rd day	Death	Sue; Lassus, Mém. cit., obs. ix; Davaine, obs. cclxxvi, p. 581; Cruveilhier, op. cit., p. 225.
85	M. 62	An enormous hydatid cyst of the liver distending the right hypochondrium, opened by a bistoury	A great number of hydatids and much yellow purulent fluid discharged	The patient died three days after the operation	Death	Récamier; Briançon, 'Essai sur le Diagnostic et le Traitement des Acéphalocystes,' Thèse de Paris, 1828, p. 16; Davaine, obs. cclxxvii, p. 582.
86	F. 47	Fluid tumour of right hypochondrium opened by two incisions; the first down to the peritoneum; and 4 days after this, the second was carried through this membrane into the cyst. The wound was kept open by a tent, and detergent injections were thrown into the cyst	Much yellow fluid was evacuated	A sero-purulent fluid continued to be discharged; but fever ensued, and the patient died soon afterwards	Death	Rayer and Velpeau, 'Bull. gén. de Thérap.,' 1844, t. xxvi, p. 58; Davaine, obs. cclxxviii, p. 583.

No.	Sex and age.	Situation of the tumour and treatment.	Nature and quantity of fluid withdrawn at the operation.	Progress.	Result and remarks.	Reference to author.
87	M. 35	Large tumour of the liver pointing in the right hypochondrium. Incision, a tent placed in the wound, hydatids and removed each day	A sero-purulent, inodorous liquid, and fetid, on the 5th day, a great number of hydatids	The discharge became fetid, and the cyst disappeared, on the 5th day, the patient died on the 6th. There were several cysts in the liver, and the omentum was strewn with cysts from the size of a nut to that of a fist	Death. Long before the tumour was opened the patient had had several attacks of severe sharp pain	M. Pinault; Cruveilhier, op. cit., p. 226.
88	F. 29	A small tumour of the liver opened by incision, clear citrine-coloured fluid at first, and discharge of brownish serum, and three days after about 6 ounces each fluid; afterwards, evacuation of hydatid cysts and into the cyst. The wound day for the next 3 weeks kept open by a plug. The discharge of hydatids facilitated by traction with a forceps. Injection of marshmallow water from the 8th day	About a pint of clear citrine-coloured fluid at first, and discharge of brownish serum, and three days after about 6 ounces each fluid; afterwards, evacuation of hydatid cysts and into the cyst. The wound day for the next 3 weeks kept open by a plug. The discharge of hydatids facilitated by traction with a forceps. Injection of marshmallow water from the 8th day	Day after the operation, fever, abdominal pain, and discharge of brownish serum, and three days after about 6 ounces each fluid; afterwards, evacuation of hydatid cysts and into the cyst. The wound day for the next 3 weeks kept open by a plug. The discharge of hydatids facilitated by traction with a forceps. Injection of marshmallow water from the 8th day	Cure	Jarjavay, 'Gaz. des Hôpitaux,' 1850, No. 89, p. 353, and No. 100, p. 397; Davane, op. cit., obs. cclxxix, p. 583.
89	M. 33	Tense tumour of epigastrium and right hypochondrium. Application of caustic potash below	A pint of limpid fluid	Progressed well for a month; the sac discharging fluid and cyst-wall, and contracting. Then	Radical cure	M. Récamier; Debouis, Thèse de Paris, 1828, No. 263.; Barrier, Thèse cit., p. 81; Cruveilhier,

90	M. 18	the ensiform cartilage; incision through the eschar, and injection of emollient fluid. After a month, enlargement of the wound	arrest of discharge, great abdominal pain, tympanitis, increase of the tumour, and dyspnoea. After the wound was enlarged, an issue of a great quantity of fetid pus, and a little sero-pus. Great relief; and two days after, discharge of cyst-wall, and increase in the flow of yellow-coloured fluid. Uninterrupted progress, and at the end of 11 months the cyst was contracted to the capacity of an ounce. Discharged cured 5 days afterwards	Cure. The At the end of 2 months, 60 to 80 minute hydatids liver was incised to the depth of 388; Davaine, obs. cdxlxxii, p. 587.	Art. "Acéphalocystes," p. 236 (?); Davaine, obs. cdxlxxii, p. 586.
		Considerable tumour of the epigastrium and right hypochondrium. After an exploratory puncture, an incision. Injection of water and alcohol into the cyst, and retention of an elastic catheter	Discharge of serous fluid, and a considerable quantity of hydatids		
		Another successful case of incision is recorded by—			
				E. Alexander, 'Boston Med. and Surg. Journ.,' vol. xviii, 1838.	

TABLE V.—Cases in which the tumour was opened wholly by caustic potash, or by spontaneous rupture.

No.	Sex and age.	Situation of the tumour and treatment.	Nature and quantity of fluid withdrawn at the operation.	Progress.	Result and remarks.	Reference to author.
91	F. 40	Enormous hydatid cyst of the right lobe of the liver. After seven applications of Vienna paste upon the centre of the right hypochondrium, the cyst gave way on the 7th day	Issue of 5 pints of puriform fluid and hydatids	Rapid amelioration; discharge through the fistulous opening for 5 months, and completion of the cure	Radical cure	Adolphe Richard, 'Bull. gén. de Thérap.', 1856, t. xlviii, p. 414; Davaine, op. cit., obs. cccxvii, p. 606; 'Med.-Chr. Rev.', 1856.
92	M. 20	Painful tumour of the right hypochondrium. Two applications of caustic potash to the most prominent part of tumour, and spontaneous opening of the cyst. Emollient injection to prevent entrance of air	Issue of 10½ pints of yellowish limpid charge of hydatids and fetid fluid continued, and in 3 weeks' time the sac had contracted to the capacity of 4 ounces; but it subsequently became distended with matter resembling that contained in the end of the small intestine, and fragments	No bad symptoms; discharge of hydatids and fetid fluid continued, and in 3 weeks' time the sac had contracted to the capacity of 4 ounces; but it subsequently became distended with matter resembling that contained in the end of the small intestine, and fragments	Radical cure	Récamier, 'Clin. Méd. de l'Hôtel-Dieu de Paris,' 1827; 'Lancet,' 1833, vol. i, p. 789; L. Martinet, "Obs. d'un Kyste hydat. du Foie," 'Revue Médicale,' t. iii, p. 436 1827; Barriér, Thèse cit., p. 58; Cruveilhier, op. cit., Art. "Acéphalocystes," p. 231; Duputyrén, 'Journ. Hebd.

93	F. 20	Large tumour of right hypochondrium, causing tension and ulceration of the abdominal walls	Discharge of clear fluid and hydatids	Recovery	of the food were observed in the discharge. These soon ceased to pass, and the patient went out of hospital in about 10 weeks from the opening of the cyst, with only a slight fistulous orifice	de Méd. et Chir., t. iii, p. 390; Davaine, op. cit., cclxxx, p. 585.
94	M. 34	Tumour of right hypochondrium, diagnosis confirmed by exploratory puncture. Application of caustic potash, after which Canquoin's paste (chloride of zinc) for 9 days successively, until the cyst was opened	9 pints of yellow fluid	Recovery	After rupture of the cyst and evacuation of its contents, the discharge was arrested to prevent hemorrhage into the cyst. This, however, occurred on the 3rd day, and the patient sank on the 4th day, after the cyst was opened	F. Plater, 'Observat. selectæ,' obs. xviii, p. 44; cité par Cruveilhier, Art. "Acéphalocystes," p. 223; Davaine, obs. cxlv, p. 508. M. Gayet, 'Gaz. Méd. de Lyon,' 15 Mai, 1865; in the 'Brit. Med. Journ.,' vol. i, 1865, p. 650.
95	F. 50	Fluctuating tumour in the hepatic region; redness of the integuments, and spontaneous rupture at the umbilicus	Discharge of liquid pus and a great number of hydatids	Death	The discharge continued a month, and death then took place from exhaustion.	Frericha, 'Dis. Liver,' Syd. Soc. Ed., vol. ii, p. 237.

No.	Sex and age.	Situation of the tumour and treatment.	Nature and quantity of fluid withdrawn at the operation.	Progress.	Result and remarks.	Reference to author.
96	F. 42	An hydatid tumour burst through the umbilicus and discharged— 3 years afterwards, the epigastric and right hypochondriac regions again swelled and became painful. The tumour was incised at a red prominent point in the hypochondrium	A pailful of a peculiar gelatinous matter Issue of blood and pus	Sanious matter and cyst-wall continued to be discharged, and a fistulous opening remained for nearly three years. 17 years after the spontaneous rupture of the cyst animal. The patient came under notice, when both apertures were found open and the surrounding parts slightly retracted to them. The whole of the proper substance of the right lobe of the liver was converted into a thick cartilaginous mass partially ossified, and imbedded therein was a large cyst containing a pint and a half of fluid and hydatids of various sizes	Spontaneous rupture of the cyst, and subsequent puncture ineffectual for destroying the animal. The patient died of albuminuria and uræmia	J. W. Griffiths, M.D., F.L.S., 'Lond. Med. Gaz.', vol. xxxiv, p. 585, 1844.
97	F. adult	Large hydatid tumour of liver bursting through the abdominal walls immediately below the umbilicus	Enormous discharge of fluid, and subsequent evacuation of large cyst was situated on the under surface of the liver, where was found a	Exhaustion and death in about 4 days. The large cyst was situated on the under surface of the liver, where was found a	Death	Dr. Bright, 'Mem. Abd. Tumours,' Syd. Soc., p. 49; 'Gay's Hosp. Rep.,' vol. ii.

98	M. 40	Tense, circumscribed tumour of right hypochondrium. Nine months after, fluctuation, reddening of skin, and spontaneous rupture	pints; then serous fluid and blood	cavity lined with false membrane and filled with grumous blood derived from the mouth of an hepatic vein	Cure	Guattani, 'De Ext. Aneur.,' p. 109, Romæ, 1772; Cruveilhier, 'Dic. de Méd. et de Chir. prat.,' t. i, Art. "Acephl.," p. 223.
99	F. 53	Hydatid tumour of the liver for 30 years. Spontaneous rupture near the umbilicus 29 years before death	"A particular liquid which occasionally presented a purulent character"	At the time of rupture a great number of hydatids were discharged through the opening, and failed to arrest the growth of this recurrent from time to time. Two tumours were found near the umbilicus which communicated with a passage full of a chalky matter, contained calcaeous tinuous with the upper part of the liver	The evacuation of the contents of the tumour Gascourt, Thèse de Paris, 1856, No. 50; Davaigne, obs. v, p. 384.	Thompson; 'Gaz. Méd. Paris,' 1844; Cadet de Gascourt, Thèse de Paris, 1856, No. 50; Davaigne, obs. v, p. 384.
100	—	Hydatid tumour of liver, rupture through abdominal parietes	—	Infiltration of hydatid fluid under the skin and abdominal muscles, causing sloughing and—	Death	G. Budd, M.D., and Sir W. Fergusson; Budd on 'Dis. Liver,' 3rd edition, p. 437.