CONTRIBUTIONS
TO THE
PATHOLOGY OF ANEURISMS AND
TUMOURS,
INVOLVING THE UPPER PORTION OF THE CHEST
AND ROOT OF THE NECK.

BY
JOHN COCKLE, M.D.

COMMUNICATED BY
F. SIBSON, M.D., F.R.S.

Received May 14th.—Read June 25th, 1867.

THE following examples of aneurismal and other tumours involving the upper part of the chest and root of the neck, are submitted to the notice of the Society as exhibiting points of interest in the pathology and diagnosis of these affections.

The first case is one of traumatic aneurism of the innominate artery, with eventually either sudden yielding or consecutive rupture of the sac. (See Plate IV, fig. 1).

John V—, æt. 35, a cabinet-maker by occupation, was admitted into the Royal Free Hospital under the author's care, June 12th, 1865, with the following history :

He had enjoyed tolerably good health until eighteen

months ago, when he fell with great violence upon the right shoulder. This injury compelled him to keep his arm in a sling for a period of three weeks. No further complaint was made until five or six weeks ago, when he felt pain in both shoulders, but more severe in the right than in the left shoulder. A fortnight previous to his admission he observed, on rising in the morning, that his throat was swollen. When admitted, the sternal end of the right clavicle was observed to be displaced forwards by a pulsating tumour extending from between the heads of the sterno-mastoid muscle into the episternal notch, and forming in this situation a considerable globular prominence (the size of a hen's egg). The impulse of the tumour was circumscribed, forcible, expansile, and highly liquid over the entire surface during the diastole of the artery, with a faint recoil during its systole. Slight silvery fremitus was felt over the swelling during each dilatation of the artery. Marked concussion impulse, and a loud and somewhat harsh arterial diastolic murmur were observed over the swelling and at the sterno-clavicular articulation. At this region the percussion sound was dull, but both dulness and murmur diminished sensibly below these limits. The murmur was transmitted into the right carotid and subclavian arteries. The contents of the tumour could be partially displaced by pressure. The trachea appeared to be pushed somewhat to the left side; the voice was harsh and metallic; the right radial pulse was weaker than the left; the right carotid and temporal arteries were also weakened in pulsation, but not to the same extent; pressure on the right carotid artery sensibly diminished the pulsation in the tumour; the superficial thoracic veins were large and turgid. The right conjunctiva was suffused, but there was no inequality in respect of size of the pupils; heart and lungs apparently healthy. June 17th.—Impulse in tumour much diminished; the tumour appears larger but flatter upon the surface; the lateral expansion increased; the superficial veins are smaller; pulse 88; respirations 20 per minute.

20th.—Has suffered from troublesome cough the last two nights with slight expectoration of frothy mucus. The axillary artery was compressed for an hour and a quarter, and the carotid artery for fifteen minutes, which was as long as the compression could be borne.

21st.—Pulsation in tumour slightly diminished; cough less frequent during the night; pulse 80; respirations 20; harsh voice absent; trachea in normal situation; compression of the axillary artery for upwards of an hour; carotid artery three quarters of an hour.

22nd.—Passed a good night; cough better; right radial pulse weaker; 76 pulsations per minute; respirations 24; impulse less over sac and slightly less liquid; pressure applied to both vessels for one hour and ten minutes.

23rd.—Bad night; complains of pain in the right cheek and in the infra-scapular region; larynx displaced to the left side; voice harsh and laryngeal; sharp, ringing cough; no difference in the size of the pupils; impulse over sac faint both during diastole and systole of the artery.

24th.—Impulse in tumour less; pressure applied to the arteries for an hour.

25th.—Tumour appears much smaller with still diminished impulse; pulse and respiration normal; pressure on the arteries reapplied.

27th.—Headache; impulse less; pressure on the arteries daily for one hour.

30th.—Impulse less, scarcely visible; the tumour feels firmer to the touch.

The case seemed to improve up to—

July 7th.—At this date pain at the back of the head was complained of, particularly after pressure upon the vessels.

9th.—Headache; a sulcus can now be seen bisecting the tumour vertically, which latter feels more solid to the touch.

11th.—Severe headache; cough troublesome at night; the tumour has become more prominent. From this date to—

21st.—The prominent symptoms were severe pains

in the head, shoulders, chest, and back; on this day additionally, pain in the sternal region immediately below the margin of the tumour, which latter has now extended to the left sterno-mastoid muscle.

23rd.—Tumour larger; impulse increased; right radial pulse absent; pain in the right arm extending from the tumour down to the elbow.

August 1st.—Severe pain in the shoulder preventing rest; the tumour presents the appearance of a cricket-ball.

4th.—Pain in the right arm as far as the wrist with occasional numbness; right pupil slightly contracted.

6th.—Bad night: feeling of pressure on the windpipe, causing a sense of suffocation; headache and cramp in the arm, to—

11th.—No change beyond enlargement of the superficial veins of the forearm; puffiness of the shoulder; loss of feeling in the little, and half of the ring finger.

15th.—Physical phenomena the same; slight clubbing of the fingers; right pupil still smaller than the left; some difficulty in swallowing.

16th.—Superficial veins of the right arm more developed; right palpebra turgid and inflamed.

19th.—Temperature of right hand and arm considerably above that of the left.

21st.—Temperature of right arm higher by 10° than that of the left.

25th.—Pulsation in sac nearly synchronous with left radial pulse; feeble pulsation in right brachial and ulnar arteries; radial pulsation absent; right conjunctiva red and lachrymose; palpebra swollen; ulceration at right outer canthus; pupil contracted.

27th, at 10 a.m.—Sudden and intense dyspnœa; pain in neck and harsh cough; said "he felt as if the tumour had burst;" pulsation entirely absent at the right wrist and also in the sac itself; tumour greatly diffused towards both right shoulder and left sterno-mastoid muscle. (See Plate IV, fig. 2.)

28th.—Tumour larger and more diffused, extending to

the left side of neck and right shoulder ; three attacks of suffocative dyspnœa.

29th.—Pulsation returned in sac ; general engorgement of the lower lip.

September 1st.—Left radial pulse small ; great extension of swelling round the neck ; face swollen and pallid ; considerable dysphagia ; food obliged to be cut very small. (See Plate IV, fig. 3.)

5th.—Tumour extending daily and becoming more diffused ; ecchymosis of skin anteriorly ; redness, heat, and tension at the lower portion of the tumour. (See Plate IV, fig 4.)

8th.—Died suddenly. Within five minutes after death the immense tumour had completely collapsed, leaving a slight depression, with an elevated ridge at its lower portion. (See Plate IV, fig. 5.)

Post-mortem examination two hours after death :—The sac collapsed as described, the sternal end of the right clavicle displaced forwards, larynx and trachea displaced an inch and a half to the left side ; congestion of the external mammary veins. The right pupil more dilated than the left, whereas, before death, the right pupil was most contracted. An incision was carried from the chin to the symphysis pubis, dividing the skin and adipose tissue of the chest, and laying open the abdominal cavity ; a second incision extended from the chin to the mastoid process of the temporal bone, and a third incision from the sternum to the acromion process of the scapula ; the integument was separated from the muscles on either side of the neck and chest in order to expose the relations of the aneurism. The connective tissue between the integument and pectoral muscles exhibited a brawny condition. Upon dissecting away the pectoral muscle on the right side, the tissue was dense and firm, resisting the edge of the scalpel ; the muscles were pale and atrophied compared with those of the opposite side ; the external and anterior jugular veins on this side were pervious ; the sterno-mastoid muscle was bulging and prominent, stretched over the sac, but pale and wasted

compared with that of the opposite side. The ribs on the left side were now cut through, the right clavicle, with first and second ribs, were divided on a line drawn from junction of the middle with the outer third of the clavicle, and turned forwards. The sternum was next cut through; the pericardium and lungs having been exposed, the subclavian muscle, vein, and artery were now seen with the upper part of the pleura: these parts formed the anterior and inferior coverings of the sac. The sterno-mastoid muscle was next divided and found intimately adherent to the sac, and, with the cervical fascia, formed its anterior and superior coverings. Tracing the sac to the left, it extended an inch and a half beyond the mesian line, displacing the œsophagus; inferiorly, it was bounded by the apex of the right pleura; anteriorly, stretching the sterno-mastoid muscle and eroding the sternal end of the clavicle which was pushed forward one inch more than that of the left side; superiorly, it extended upwards on a level with the thyroid cartilage; externally, it extended as far as the trapezius muscle, having the cervical fascia, considerably thickened, for its chief covering; posteriorly and internally the sac was bounded by the common carotid artery, internal jugular vein, pneumogastric and sympathetic nerves, trachea, larynx, and œsophagus, all of which organs were compressed and displaced towards the left side. An aperture of somewhat quadrangular shape exists in the anterior wall of the innominate artery, with this exception, the artery is perfectly healthy throughout its entire extent. Neither does the slightest disease exist in any other part of the large vessels or in the heart itself. No appearance of any former injury of the right clavicle is discernible.

CASE 2.—This case (see Plate V, fig. 1) most strikingly resembles that last described, so far as relates to the exact position of the pulsating tumour in its earliest stage, but differs from it signally in its spontaneous origin, and in being most probably one of true aneurism or dilatation of

the ascending and origin of the transverse aorta, innominate and root of the right common carotid arteries.

Mary Ann E—, æt. 52, a married woman, was admitted into the Royal Free Hospital under the author's care, September 12th, 1866, with the following history :—She had always enjoyed good health until twelve months previous to her admission, when she first felt a pulsating swelling in the throat (episternal notch). This swelling came on gradually, without any obvious cause, and was soon followed by dimness of vision in the right eye; the only remaining symptoms complained of have been pain, at times, near the top of the sternum, and an occasional sense of choking; the pain complained of seemed, in part, external, and the tissues, some time since, felt somewhat thickened over the region of the right sterno-clavicular articulation and upper part of the sternum, and still remain tender even under the gentlest percussion. The tumour appears to arise near the former region, encroaching upon the outer head of the right sternomastoid muscle and passing into and filling the hollow above the episternal notch. The origin of the right common carotid can be felt dilated. The tumour has varied considerably in point of visible size; some weeks ago, about the period of admission, it was much more prominent than at the present time, bulging out as large as a hen's egg from the surface. The impulse has invariably been forcible, markedly expansile, and very liquid, with slight thrill (diastolic arterial) limited to the space between the heads of the right sterno-mastoid muscle. Perfect collapse of swelling during systole. A loud and somewhat harsh murmur is audible over the tumour during the arterial diastole, and, though much weaker, during the systole; it passes loudly during arterial diastole into the right carotid, subclavian, and brachial arteries, and is even faintly heard in the arteries of the left side. The double murmur, well marked, passes down the aorta, but is most decidedly weakened near the aortic mouth and at the apex of the heart. The pulsation in the carotid arteries has been, as a rule, equal, but a strongly marked difference has existed in the right radial pulse, which

is at times full and strong, almost jerking in character, at other times very sensibly weakened, as is well shown in the sphygmographic tracings taken at various times by Dr. Anstie, Dr. Sibson, and the author. There is some dulness on percussion over the upper portion of the sternum, but no tactile or stethoscopic impulse can be detected. The position of the heart, its sounds, and impulse, are normal; the lungs also appear in a physiological condition. This patient has just quitted the wards of the hospital (June 23rd), her condition being unchanged.

CASE 3.—This case is an interesting complement to the last, being one, probably, of more general dilatation of the aorta, and also involving the innominate trunk and origin of right carotid artery.

Emma R—, æt. 46, married, has been an out-patient of the Royal Free Hospital, under the author's care, for nearly five years. She first complained of swelling and pulsation in the hollow of the throat, speedily followed by dyspnœa, pain in the chest, and very troublesome cough. An intermitting prominence is visible at the lower part of the episternal notch, the swelling becomes very marked during the arterial diastole, but quickly emptying itself during systole; the swelling extends more definitely from behind the right sterno-clavicular articulation along the tracheal border of the sterno-mastoid muscle, where the arterial enlargement is readily traceable; abnormal pulsation existed, though in a less degree, in both carotid and subclavian arteries. In the earlier stage neither thrill nor murmur could be detected; the pulsation of both radial arteries was comparatively weak: the second cardiac sound was unusually sharp and clear. At a later period, in the progress of the case, the pulsation in both radial arteries became somewhat jerking in character, and a double murmur, louder and harsher during arterial diastole became audible down the aorta, extending to the apex of the heart; the impulse of this organ being increased, and signs of hypertrophy of the left ventricle being superadded. Marked percussion dulness

exists at the right sterno-clavicular junction and down the entire course of the ascending aorta. The tracings attached were taken some weeks ago by Dr. Anstie. The left jugular vein is nearly always in a state of permanent distension, and presents, at times, visible pulsations. Dyspnoea and constant cough are permanent.

CASE 4.—*Vast aneurism of the aorta, with secondary sacculation ascending above the sternal end of the left clavicle over the site of the carotid artery.* (See Plate V, figs. 2 and 3).

Deborah W—, æt. 36, sempstress by occupation, married, has had four children, and one miscarriage. She was admitted an out-patient of the Royal Free Hospital, October 19th, 1866. Ten years ago she contracted syphilis from her husband, and since that time has been so maltreated by him that she thinks her chest may have sustained some injury from his violence. During the last two years she has lost flesh very considerably, otherwise she had previously enjoyed tolerably good health with the exception of cough and cold during the winter months. Nine months prior to her admission to this hospital she began to feel pain in the region of the left sterno-clavicular articulation, which pain she considered to be rheumatic, and for which she thinks she was treated at University College Hospital, where she became a patient. This pain she described as being, at times, intolerable. About this period a gradually increasing hoarseness occurred, uninfluenced by either local or constitutional treatment. Three months later she observed a swelling in the situation of the articulation named. When first seen here, a firm, unyielding tumour was detected in the neighbourhood of the left sterno-clavicular articulation and inner third of the clavicle, pushing upwards and forwards the clavicle and clavicular attachment of the sterno-cleido-mastoid muscle, the inner third of the clavicle being lost, as it were, in the tumour, the whole mass conveying the impression of an osteoid growth. No impulse existed over the swelling, but a double sound could be heard through the tumour as

if communicated by the common carotid artery over which the tumour seemed to be immediately lying. At this time, iodide of potassium was administered internally, and an opiate lotion applied over the swelling.

October 26th.—Seven days afterwards the tumour was smaller in size and less painful; the patient expresses herself relieved, but there is no apparent change in respect of density of the tumour to the touch.

November 15th.—About three weeks subsequently she was transferred to the author's care. She was now greatly emaciated, with a marked leaden tint of complexion and livid lips. The breathing was quickened, although she made no complaint of dyspnœa. Indeed, her only complaint was of pain in the old spot, weakness of her voice, and occasionally double vision. A tumour was to be seen, about the size of a small hen's egg, displacing the left sterno-clavicular articulation, and passing up behind the inner third of the clavicle beneath the origin of the sterno-mastoid muscle, encroaching somewhat upon the hollow of the throat, and extending about an inch and a half up the neck in the track of the carotid artery, slightly displacing the larynx to the right. A very slight visible impulse existed in the tumour; the skin covering it was ecchymosed with a congested state of the capillaries surrounding it. The veins below the clavicle, more particularly upon the left side, were distended and tortuous. No bulging of the general surface of the chest was perceptible, beyond a fulness at and below the left sterno-clavicular articulation corresponding to about the second intercostal space. At this stage of the examination the impression was conveyed of the case being aneurism of the carotid artery. Upon further examination, however, the tumour seemed to fuse into the substance of the clavicle itself, whilst the faint impulse, almost limited to the apex of the tumour, felt very liquid; the apex itself was readily depressed with the point of the finger; no murmur whatever could be detected over the swelling. The upper third of the entire chest, anteriorly and posteriorly, was uniformly dull upon percussion, with a marked sense of tactile resistance.

No unnatural impulse whatever could be detected by the hand firmly placed over the area of dulness. Below this area, the percussion sound was clearer, but yielded by no means a natural resonance. The area of the heart's impulse was increased; the impulse was forcible and jerking, and the cardiac sounds muffled and indistinct, but without murmur or thrill. The circulation was very feeble, entirely absent in the left radial and ulnar arteries, and exceedingly feeble in the right radial artery. All respiratory murmur was extinct over the upper portion of the chest, but over the remaining part slight dry, and liquid râles were audible. At the root of the neck, on either side, an impulse existed, one of simple elevation, without either thrill or murmur, as if some very solid body were heaved up at each contraction of the heart. At this stage of the case great doubt was entertained whether it was one of intra-thoracic tumour, or a large aortic aneurism filled with solid clot; the latter conclusion was eventually drawn, based principally upon the peculiar elevation on either side of the root of the neck during each contraction of the heart, and the general condition of the circulation.

After some short interval (two or three weeks) the following changes occurred:—The tumour in the neck subsided, completely, permanently, and with comparative quickness. In proportion to its disappearance, the external chest veins increased in size and became more tortuous and, now, pulsating. (See Plate V, fig. 3.) Pulsation could no longer be felt in the right radial, carotid, and temporal arteries. When the patient was placed recumbent, faint pulsation was to be felt in the abdominal aorta and crural arteries. The only change in the symptoms was an increase in the frequency of double vision, contraction of the left pupil, transient loss of power in the limbs, and occasional momentary unconsciousness on awaking in the night. The patient (refusing constantly to be admitted into the wards) managed, in this condition, to walk to the hospital twice in the week, for a few weeks. No pulsation could ever be detected at the wrists, although frequently examined by

different observers. She fell suddenly dead while in the act of preparing some vegetables for dinner—a not frequent example of aortic aneurism terminating suddenly, without rupture of the sac or more than ordinary exhaustion. The death was, probably, syncopal. The author is indebted to Mr. J. D. Hill for the account of the post-mortem examination. The entire specimen is preserved, but the sac has not been laid open.

Post-mortem examination twenty-four hours after death.—

The body was greatly emaciated, and exhibited bruises upon the face and nose. No tumour was visible externally. When the chest and abdomen were laid open, an immense aneurismal tumour occupied the upper half of the left side of the thorax, encroaching upon the upper and inner half of the right side. The sac was adherent, in front, to the under surface of the left clavicle, first rib, and intercostal muscles, and, posteriorly, to the contents of the posterior mediastinum by dense areolar tissue. Upon the right side the sac encroaches upon the right lung, being adherent to the pleura, and drawing the pericardial layer of pleura, resting upon the root of the lung, and compressing the bronchial vessels and tubes. Upon the left side the sac crowds the lung into the outer half of the chest; it is adherent to both layers of pleura, and presses upon the bronchial tubes and vessels. The lung is quite solid, and adherent at its apex to the upper part of ribs and sac. Superiorly, the sac is still seen above the clavicle; the contents of the posterior mediastinum are compressed, and pushed out of their normal position. The trachea is dislocated two inches to the right side: the œsophagus is also displaced, though to a less extent. Inferiorly the sac is attached to the pericardium, and displaces the heart downwards, the apex of this organ corresponding to the seventh intercostal space two inches beyond the costal attachment of the cartilages. The right lung was condensed at its base, and adherent to the sac. The left subclavian artery is very much smaller than the right. Atheromatous disease, in an early stage, exists in the right carotid

artery. The left innominate vein is obliterated, and its place probably supplied by a branch of the thyroid vein coursing across the front of the sac between the sterno-mastoid and sterno-hyoid muscles of the left side, and finally taking a direction towards the right internal jugular vein. This vein communicates with or takes its origin from the left internal jugular or subclavian vein. The spleen was large, as also the liver; its right lobe was dislocated below the umbilicus. The gall-bladder was filled with gall-stones. No other organs were much changed. The brain, however, was not examined.

CASE 5.—Large cervical tumour of the left side of the neck, accompanied with obscure pulsation around its base.

Ann B—, widow, æt. 49, was admitted into the Royal Free Hospital under the author's care, February 12th, 1867, with a large tumour occupying the left side of the neck. She had suffered for six months from pain in the neck and left infra-clavicular region darting through the chest; at the expiration of this time, severe cough came on, and shortly after, she observed a fulness just above the left collar-bone. Upon her admission there existed a hard, bulging, and very slightly elastic tumour with some diffused impulse at its base. At the root of the neck the swelling corresponded to the inner two thirds of the left clavicle, extending upwards and inwards; it dislocated the larynx and trachea to the right side; externally, the swelling pushed the trapezius muscles upwards and backwards. The skin was red and livid in patches, over the surface. The face, forehead, and nose were of deep leaden tint. Very great dyspnœa existed, with violent paroxysmal cough. No abnormal signs detected within the chest, but the left radial pulse was weakened.

19th.—Dyspnœa greatly increased, amounting to veritable orthopnœa; cough almost incessant and exhaustive, accompanied by scanty mucous expectoration; severe pain in the head. The tumour presents the appearance of a large carbuncle. As no intra-thoracic change is obvious beyond

consonating râle, the great dyspnœa and cough are probably caused, for the most part, by the mechanical pressure of the swelling. Still, the existence of the pain in the chest for six months, leaves the diagnosis doubtful.

March 1st.—The swelling presents some points of indistinct fluctuation. Black, gangrenous patches are seen here and there over the surface, dyspnœa and cough still urgent; great constitutional distress.

20th.—A large slough separated, followed by copious purulent discharge. Immediate relief to the cough and dyspnœa; return of larynx and trachea to their normal position.

April 1st.—Nearly convalescent, though still complaining of pain below the left clavicle; the large wound filled up by granulation.

Two cases of subperiosteal abscess have been noticed; one occurred in a young woman, and was situate just above the left mamma, and connected with disease of the rib. The second case occurred in a male, and was situate on the centre of the sternum. These examples of abscess are interesting from their occupying the site of intrathoracic aneurisms, and occasionally receiving concussion impulse from the heart. No difficulty, however, ordinarily attends their diagnosis.

Remarks.—Whenever a pulsating tumour which involves the root of the neck to any extent is first submitted to observation, and is unaccompanied by a sufficiently accurate previous history, there is, generally, a felt difficulty of diagnosis, whatever the actual position of the tumour may be. Such difficulty may occasionally arise from an inability to determine the real nature of the tumour, but most frequently (an aneurismal character being assumed) it consists in isolating the vessel engaged.

The general rules laid down for guidance in this latter particular are based almost mainly upon the position of the tumour, and the relative state of the circulation in the

corresponding arteries of the arms and neck ; the exceptions, however, to such rules are sufficiently frequent to lessen their value to a very considerable extent.

It is not within the scope of this communication to attempt more than a brief illustration of some of these exceptions, principally from cases of aortic aneurism giving rise to secondary sacculations ascending the neck, at the ordinary site of innominate or carotid aneurism, and causing modifications in the circulation and sensibility of the right upper extremity.

Plate VI, fig. 1, represents the case of a man, an Irish drover, admitted into the Hospital under the writer's care, for (supposed) subclavian aneurism.

He had suffered for some time from pain and stiffness in the neck, right shoulder, and clavicle. Eventually a tumour *first appeared above the right clavicle near its sternal end*, and gradually ascended the neck, as seen in the drawing. There was an entire absence of pulse in the right radial, ulnar, and brachial arteries. The carotid artery was apparently felt beating above the sac. The left radial pulse was of normal volume, but quickened in pulsation, ranging at 90. The motility of the right arm was decidedly lessened. This was a case of aneurism of the ascending aorta, and although the innominate artery was involved, it was a part of the aortic sac which mounted up the neck.

Plate VI, figs. 2, 3, represents a second example of aortic aneurism sending loculi up the neck in the site of innominate aneurism. One of these tumours mounts at the episternal notch, and a second, just at the outer border of the right sterno-mastoid muscle. In this case, James P—, the man, suffered from violent pain in the shoulder and right arm, with subsequent impairment of sensation and motion in this extremity ; marked atrophy of the arm followed. There was considerable weakening of the circulation at the wrist.

The difficulties that attend the differential diagnosis of innominate aneurism proper are in tolerably direct pro-

portion to the part of the trunk affected. When the origin of the vessel is alone involved (so far as the author's experience extends), it is *invariably* combined with aneurism, either true or false, of the ascending aorta. This form of innominate aneurism thus loses all individuality (so to speak), and merges into the general pathology of aortic aneurism. It has absolutely no clinical history of its own. If, as at times (though with extraordinary rarity), an aneurism should limit itself to the mid-portion of the trunk and anterior wall, it may remain for a time, or even throughout its entire course, intra-thoracic, descending and moulding itself upon the aorta, evoking all the physical phenomena and symptoms of aneurism of the ascending and transverse aorta, or it may additionally, by one or more loculi, ascend the neck either on the tracheal or acromial side of the sterno-cleido-mastoid muscle. So that should a pulsating tumour, with or without cervical extensions, occupy the space corresponding to the second and third costal cartilages of the right side, and upper portion of the sternum, and present the ordinary symptoms of aneurism, though we may be practically safe in locating the disease in the aorta, it is not from any inherent difference of signs of symptoms, but from the rarity alluded to, of intra-thoracic innominate aneurism.

It is only when the distal end of the trunk becomes the seat of aneurism that accuracy of diagnosis is attainable, though, in such a case, the origins of the subclavian and carotid arteries often share the dilatation. Certain signs and symptoms have been ascribed to innominate aneurism. Their value is high, directly as intra-thoracic disease can be eliminated; it becomes, as has been just shown, extremely small as the evidence for such disease increases. This is by far the most important point in diagnosis. The only possible source of error is the rare fact of an aortic sac arising from the convex wall of the arch, and mounting up the neck. Valuable aid is afforded by the knowledge of some antecedent local injury, or violent and unusual effort (for very many of such cases are traumatic). These points

determined, should severe pain of the right shoulder, clavicle, right side of the heart, neck, and arm, with local venous congestion or œdema precede the appearance of a pulsating tumour which, emerging from beneath the right sterno-clavicular articulation, and often displacing the head of the clavicle during its ascent, distends the space between the heads of the sterno-mastoid muscle, and fills the *episternal notch*—should there be a marked weakening of pulsation in the right radial and carotid arteries, pressure upon one or other artery lessening the impulse on the tumour—should a murmur, loudest over the tumour and at the sterno-clavicular articulation, diminish downwards, yet ascend the arteries on the right side, if pervious, such murmur being absent upon the left side, we may with tolerable safety diagnose an aneurism of the terminal portion of the innominate artery; at least such were the rules which led to a successful diagnosis in Case 1.

DESCRIPTION OF PLATES IV, V, AND VI.

PLATE IV.

Fig. 1, Case 1.—John V—. Case of traumatic aneurism of the innominate artery, June 12, 1865.

Fig. 2, ditto, August 27.—Tumour greatly diffused towards right shoulder and left sterno-mastoid muscle.

Fig. 3, ditto, September 1.—Great extension of swelling round the neck (outline).

Fig. 4, ditto, September 5.—Tumour more diffused; ecchymosis of skin, &c. (outline).

Fig. 5, ditto, September 8.—Condition at moment of death.

PLATE V.

Fig. 1, Case 2.—M. A. E—. Spontaneous true aneurism of the ascending and transverse aorta.

Fig. 2, Case 4.—Deborah W—. Vast aneurism of the aorta, with secondary sacculaton above sternal end of left clavicle. November 20th, 1866.

Fig. 3, ditto.—Ditto, February 2nd, 1866.

PLATE VI.

Fig. 1.—Case of aneurism of the ascending aorta, involving the innominate artery in an Irish drover.

Fig. 2.—Second case of aortic aneurism sending loculi up the neck in the site of innominate aneurism. Case of James P—, July 3rd, 1866.

Fig. 3.—Ditto, ditto, August 4th, 1866.

Fig. 1.

Fig. 2.

Fig. 3.