Published in final edited form as:

Semin Cancer Biol. 2007 December ; 17(6): 436–443.

Clinical and biological significance of HLA-G expression in ovarian cancer

Jim Jinn-Chyuan Sheu^{a,b,c} and Ie-Ming Shih^{a,*}

^a Departments of Pathology, Johns Hopkins Medical Institutions, Baltimore, Maryland, USA

^b Human Genetic Center, China Medical University Hospital, Taiwan

^c Graduate Institute of Chinese Medical Science, China Medical University, Taiwan

Abstract

Ovarian cancer is the most lethal gynecologic neoplastic disease in which the molecular etiology remains largely unclear. Like other cancer types, evolution of ovarian tumor cell species is accompanied by acquisition of novel gene products and these new tumor-associated antigens elicit a host immune response that creates selection pressure upon the emerging tumor clones. One of the mechanisms that ovarian cancer cells evade immune surveillance is by up-regulating human leukocyte antigen-G (HLA-G) expression. HLA-G is a non-classical MHC class I molecule and accumulated evidence has suggested its biological role in inactivating immune response. It has been well known that HLA-G expression is frequently detected in the most aggressive type of ovarian cancer, i.e., high-grade serous carcinoma, and measurement of HLA-G protein levels has shown promise for detection and prognosis prediction in ovarian cancer. This review summarizes those recent studies on HLA-G expression in ovarian cancer with special focus on its clinical and biological significance which is fundamental to elucidate the molecular mechanisms in ovarian cancer development and paves the foundation for future HLA-G based diagnostics and therapeutics.

Introduction

Ovarian cancer represents one of the most lethal neoplastic diseases in women. In 2006, it has been estimated that 20,180 women will be diagnosed with ovarian cancer and 15,310 will die from the disease in the United States alone (<http://www.cancer.org/docroot/home/index.asp>). Currently, the incidence of ovarian cancer in the United States ranks eighth among cancers (excluding skin cancer), but fifth in terms of mortality in women. The above figures are similar to those in other developed countries (<http://www-dep.iarc.fr/>). The high mortality of ovarian cancer is mainly due to a lack of success in diagnosing ovarian cancer at an early stage and treating it effectively. Until the basic biology of ovarian cancer is more clearly understood, attempts at developing new diagnostic methods and therapy will remain empirical.

The increased vulnerability to major human diseases including ovarian cancer can be viewed as a result of mal-adaptation, i.e., compromise and trade-offs, of the evolutionary processes [1]. For example, the feto-placental unit which represents a semi-allograft has developed molecular strategies that effectively abort maternal immune surveillance in placentas to ensure the success of pregnancy [2]. On the other hand, ovarian tumor cells take advantages of similar

* Correspondence author: Tel: +1-410-502-7774, Fax: +1-410-502-7943, e-mail address: ishih@jhmi.edu (Ie-Ming Shih).

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

molecular mechanisms to escape from immune recognition and attack in response to the expression of panoply of tumor-associated antigens. The mechanisms shared by feto-placental and cancerous tissues to counteract immune responses are complex and involve at least the expression of non-classical major histocompatibility (MHC) antigens. Numerous studies have provided cogent evidence that human leukocyte antigen-G (HLA-G), a non-classical MHC molecule, participates in immune regulation, especially in immune tolerance in physiological process such as pregnancy. Shortly after the identification of HLA-G in trophoblastic cells [3], there has been increased interest in studying HLA-G expression in neoplastic diseases including ovarian cancer (Fig. 1). These studies provide *in vivo* evidence demonstrating HLA-G expression in human ovarian tumor tissues and support the view that HLA-G may participate in tumor development by suppressing immune regulation within tumor microenvironment. Besides, several translational studies suggest the potential roles of HLA-G as a novel biomarker in clinical applications such as cancer detection, diagnosis and prediction of clinical outcome in ovarian cancer patients [4]. This review summarizes those recent studies of HLA-G expression in ovarian cancer with special emphasis on its potential clinical applications and possible molecular mechanisms of how HLA-G molecules contribute to ovarian cancer development.

Molecular pathways and model in the development of ovarian cancer

Ovarian neoplasm is composed of more than 100 histological subtypes based on the World Health Organization (WHO) classification [5]. Among them, epithelial tumors (i.e., carcinomas) are by far the most common malignant tumors and are what generally referred as “ovarian cancer”. Despite considerable efforts aimed at elucidating the molecular mechanisms of ovarian carcinoma, its pathogenesis is still unknown due to the lack of an established model for its development [6,7]. Recently, a new model for the pathogenesis of ovarian cancer based on clinical, pathological, and molecular genetic studies has been proposed [8,9]. In this model, ovarian carcinomas are divided into two broad groups designated Type I and Type II. Type I tumors are slow growing, indolent tumors that are generally confined to the ovary at diagnosis and develop from well established precursor lesions that are termed “borderline” tumors. Type I tumors included low-grade serous carcinoma, mucinous carcinoma, endometrioid carcinoma, and malignant Brenner tumors. From a molecular genetic standpoint, they are characterized by mutations in a number of different genes including KRAS, BRAF, ERBB2, PTEN, and beta-catenin. In contrast, Type II tumors are rapidly growing, highly aggressive neoplasms for which well defined precursor lesions have not been well recognized. The vast majority of ovarian carcinomas belong to the Type II category. These tumors include high-grade serous carcinoma (most common), malignant mixed mesodermal tumors (carcinosarcomas) and undifferentiated carcinomas. Type II carcinomas are characterized by a very high frequency (~80%) of p53 mutations and a high level of genetic instability [8,10]. The model not only has important clinical implications but also suggests the fundamental difference in biological aspects between Type I and Type II ovarian neoplasms. It is likely that Type II carcinoma cells are equipped with unique molecular machinery that contributes to their highly aggressive clinical behavior. For example, as will be discussed below, HLA-G is frequently expressed in Type II ovarian carcinoma but only rarely in Type I carcinoma.

Expression of HLA-G in different types of ovarian carcinoma

In normal human tissues, constitutive HLA-G expression is restricted to a few tissue types including trophoblastic cells and thymic epithelium [3,11]. In contrast, analysis of HLA-G expression in human cancers demonstrated that HLA-G gene was upregulated in a wide variety of human neoplasms. These tumors include ovarian carcinoma, cutaneous melanoma, lung carcinoma, endometrial carcinoma, gastric carcinoma, hematopoietic tumors, renal cell carcinoma, mesothelioma, breast carcinoma, trophoblastic tumors, glioma tumors, bladder

carcinoma, colorectal carcinoma and lymphoproliferative disorders (Fig. 2) [12-22]. In ovarian cancer, immunohistochemistry study has demonstrated that HLA-G immunoreactivity was detected in 45 (61%) of 74 ovarian high-grade serous (Type II) carcinomas but was rarely detected in Type I carcinomas [16]. This finding is of great interest and raises the possibility that HLA-G expression is not only associated with Type II ovarian cancer but may confer tumor cells the “aggressiveness” through its role in modulating tumor immune response in tumor microenvironment. It should be noted that the percentage of HLA-G expressing tumor cells varies widely among ovarian cancer specimens, and the HLA-G positive cells range from 2 to 100% within a specimen. Furthermore, secreted form of HLA-G (sHLA-G) is detected in the supernatant of malignant effusions from ovarian cancer patients [16,23-25] and more importantly, increased plasma levels of sHLA-G were detected in patients suffering from a variety of neoplastic diseases including ovarian cancer [25,26]. These studies provide important biological and clinical implications for HLA-G expression in human ovarian cancer, supporting the view that HLA-G in its membrane bound and secreted forms may participate in tumor development by suppressing the local immune response to cancer cells. For example, it is likely that the HLA-G non-expressing cells in adjacent to expressing cells can also be protected from immune attack because sHLA-G that is synthesized by the HLA-G expressing cells diffuses and acts remotely to inactivate immune response within the tumor.

Possible mechanisms underlying HLA-G expression in ovarian cancer

As previously described, HLA-G is highly expressed in Type II ovarian carcinomas but the molecular mechanisms that upregulate its expression remain unclear. Since the DNA copy number of the HLA-G gene locus does not increase in ovarian cancer genome [27], elevated HLA-G promoter activity may be important for its upregulation. Furthermore, a variety of environmental cues such as hypoxia [28,29], stress [30], hormones [31,32], and certain cytokines [33-37] may also be involved to modulate the HLA-G promoter activity. Sequence analysis of the HLA-G gene promoter reveals that almost all regulatory motifs described for classical MHC class I genes are not conserved in the HLA-G promoter [38]. This finding suggests that the mechanisms involved in the transcriptional regulation of HLA-G are unique. Recent studies have demonstrated that HLA-G gene transcription activity could be controlled by *cis*-acting epigenetic mechanisms including DNA methylation and histone acetylation [39-41]. For example, primary tumor cells gradually downregulate HLA-G expression in culture [29,42] and cells treated with histone deacetylase inhibitors or DNA demethylating agent restore their HLA-G expression [38,39]. The epigenetic regulation is even more pronounced under certain microenvironmental conditions, such as hypoxia [42] which occurs very frequently in ovarian carcinoma, especially in tumor ascites. Stabilization/activation of hypoxia-inducible factor I (HIF-1) is the key cellular response upon hypoxia conditions. HIF-1 acts as a transcription regulator that controls a wide variety of gene expression including HLA-G in response to hypoxia [28].

Tumor microenvironment such as ascites in ovarian cancer is enriched by a variety of cytokines released by tumor-infiltrating lymphocytes. Interaction between cytokines and tumor cells regulates HLA-G expression. For example, GM-CSF and IFN- γ secreted by the infiltrating cytotoxic T cells have been shown to enhance HLA-G expression in tumor cells [17,43,44]. On the other hand, secretion of IL-10 by tumor cells can also upregulate HLA-G expression level in tumor tissues through an autocrine or paracrine fashion [18,33,34]. HLA-G, in turn, modulates the cytokine expression in immune cells to produce a profile of the Th2 type cytokines including IL-10, IL-4 and IL-3. Clinical studies have shown that this shift towards the Th2 cytokine profile, especially the enhanced IL-10 secretion, is frequently associated with an impairment of anti-tumor immunity [45,46]. HLA-G synergizes with IL-10 in suppressing the local immune response in tumor microenvironment and contributes to tumor progression of lymphoma from a low-grade to a high-grade disease [12]. Indeed, co-expression of HLA-

G and IL-10 has been shown to correlate with cancer at advanced clinical stages [18,34,47]. For example, HLA-G expression was found in 51% primary cutaneous T-cell lymphoma, whereas 73% of HLA-G positive cells also expressed IL-10. The tumors co-expressing HLA-G and IL-10 tend to be clinically more aggressive than those without [34,47-49]. Although these studies were performed primarily in other cancer types, it is plausible that similar mechanisms may also operate in upregulating HLA-G expression in ovarian carcinomas.

The biological significance of HLA-G expression in ovarian cancer

It has been long thought that the biological significance of HLA-G expression in cancer is mainly to abort immune surveillance and compromise the attack of tumor cells by immune effector cells. The evidence supporting the role of HLA-G in immune suppression in ovarian cancer primarily comes from the clinical studies in correlating HLA-G expression levels and clinical outcome in pregnancy and organ transplantation which represent two main conditions for host to respond to non-self tissues. In pregnancy, it has been well accepted that maternal immune-tolerance to the “semi-allograft” fetus and placenta is attributed to the presence of HLA-G. The fact that trophoblastic cells in human placentas express very high level of HLA-G suggests its involvement in regulating immune reaction in the fetal-maternal interface [50]. Recent studies have confirmed the predictive value of HLA-G expression in pregnancy outcome. The presence of sHLA-G by early conceptus appears to be essential for a successful implantation and sHLA-G has been used as a reliable marker of increased chance of subsequent pregnancy following *in vitro* fertilization [51-54]. The presence of HLA-G transcript in pre-implantation embryos is correlated with increased blastocystic cleavage rate and greater number of blastomeres per embryo following *in vitro* fertilization [55]. In contrast, the abortion rate increases for the women who previously received embryos with low or undetectable HLA-G expression [53]. A null HLA-G allele in a conceptus is also associated with recurrent miscarriage in women [56,57].

Transplanted allogeneic organs are always rejected by the host immune surveillance system unless immunosuppressive reagents are administered. sHLA-G proteins are usually detected at high levels in the blood samples from transplantation patients whose grafted tissues survive [58]. In addition, HLA-G positive infiltrating monocytes are observed in the grafted tissues due to the binding of sHLA-G molecule to its receptor on the immune cells. The above clinical correlation studies are consistent with the view that HLA-G expression plays an important role in immunosuppressive functions in pregnancy and allograft rejection. It is very likely that the similar molecular mechanisms may also be operated by ovarian cancer cells to escape from host's immune surveillance.

In the past several years, major advances have been made in delineating the molecular mechanisms of HLA-G in immunosuppression. Tumor development and progression are always accompanied by expression of novel tumor-associated antigens which can elicit an immune reaction by activating immune defenders including cytotoxic T cells, natural killer (NK) cells and macrophages. HLA-G expression in ovarian cancer cells represents one of the possible mechanisms to overcome the vigilant immune surveillance and attack. A growing body of evidence demonstrates a direct interaction between HLA-G and the leukocyte immunoglobulin (Ig)-like receptors, LIR1 (also known as LILRB1/ILT2/CD85J) and LIR2 (LILRB2/ILT4/CD85D), and between HLA-G and the killer cell-Ig-like receptor KIR2DL4 (CD158D) (Fig. 3) [59,60]. Both LIR1 and LIR2 belong to a family of immunoreceptors expressed on T cells, B cells, monocytes (macrophages), dendritic cells and NK cells that upon ligand binding can inactivate those immune effectors [61,62]. On the other hand, the killer cell-Ig-like receptor, KIR2DL4, belongs to the gene family of killer cell inhibitory receptors which upon binding to HLA-G inhibit NK cell-mediated cytolytic activity [63]. Although these Ig-

like receptors also interact with other HLA class I ligands, they show the highest binding affinity to HLA-G [64].

HLA-G binding receptors share a common cytoplasmic ITIM sequence (immunoreceptor tyrosine-based inhibitory motif) that upon HLA-G binding recruit intracellular protein-tyrosine phosphatases and trigger an inhibitory signal cascade. The negative signaling, in turn, down regulates the activation of various immunoresponsive genes and affects the cytokine/chemokine profiles secreted by the effector cells. As a result, cell maturation and clonal expansion of immunocompetent cells are significantly reduced. Interestingly, both membrane-bound and secreted forms of HLA-G are capable of upregulating the expression levels of its binding ligands, the Ig-like inhibitory receptors, forming a positive regulatory loop [65]. Thereby, the immune effector cells become more sensitive to the HLA-G mediated inhibitory effects, resulting in immune tolerance of tumor cells.

Clinical significance of HLA-G expression in ovarian cancer

Cancer detection in body fluid

Many patients who suffer from ovarian cancer presented with ascites which is an accumulation of fluid and floating tumor cells in the abdominal cavity. However, in addition to neoplastic diseases, ascites formation is related to a variety of benign diseases including inflammatory disorders, infection, cardiac, liver and renal disorders. Cytology examination of representative cells obtained from ascites is routinely performed for diagnostic purposes, but the sensitivity of cytology has been estimated to be 60% at best [15]. Thus, detection of tumor-associated biomarkers such as sHLA-G would facilitate the diagnosis in this clinical setting. Singer *et al.* applied a sensitive ELISA to determine sHLA-G levels in the supernatant of ascites from ovarian and breast carcinoma and benign controls [6]. They found that malignant ascites contained higher levels of sHLA-G than benign ascites. The investigators based on receiver operating characteristic (ROC) curves and demonstrated an excellent performance of HLA-G ELISA as the diagnostic tool to detect ovarian and breast cancer with an area under the ROC curve of 0.95. The presence of sHLA in ovarian carcinoma tissues strongly suggests that sHLA-G can be used as a tumor marker in blood for cancer detection. In fact, the serum levels of sHLA-G as detected by enzyme-linked immunoassay (ELISA) were elevated in patients with a variety of neoplastic diseases including ovarian cancer [25]. Although this study was preliminary, it provides another promising biomarker for future clinical trials to determine its potential clinical applications, especially for its role in detecting early stage and curable diseases.

Although sHLA-G may prove to be useful in assisting the diagnosis of malignant versus benign clinical conditions, there are several challenges ahead. For example, a large number of cases should be assessed to determine the optimal cutoff of sHLA-G for potential clinical applications. In addition, it is important to demonstrate a higher performance of sHLA-G than previously published or pre-existing ovarian cancer markers. Further studies are required to assess whether a combination of selected secreted biomarkers including sHLA-G may provide a better strategy in increasing both sensitivity and specificity than single marker alone for cancer detection in body fluids. Finally, it will also be interesting to address how age, menopausal status, histologic grade, and other clinical parameters affect HLA-G levels in body fluid including blood and ascites samples.

HLA-G as a marker associated with clinical outcome

Because of the biological implication of HLA-G in tumor immunology and biology, several studies have focused on clinical correlation between HLA-G expression and patient's clinical outcome including overall survival and the risk to develop metastatic diseases. Previous reports

have shown that such correlation depends on the cancer types and specific clinical settings as HLA-G expression can be associated with either a favorable or an unfavorable clinical outcome. In ovarian carcinomas, HLA-G expression in effusions has been shown as a possible marker of tumor susceptibility to chemotherapy [66]. In that study, the level of HLA-G expression in ovarian cancer cells was significantly lower in effusions obtained during or after chemotherapy than that without chemotherapy. Besides, the presence of HLA-G positive tumor cells in effusions obtained prior to the institution of chemotherapy significantly correlated with better overall survival. Similarly, Ishigami *et al.* found that the 5-year survival rate in gastric cancer patients whose tumors were positive for HLA-G was 78%, which was significantly higher than 51% in the HLA-G negative group [20]. In that study, HLA-G positive group had a more differentiated histology, less nodal invasion and earlier clinical stage than HLA-G negative patients. The above findings appear paradoxical to the expected role of HLA-G in propelling tumor progression. One of the possible explanations is that the HLA-G expressing cells may become more susceptible to elimination by the chemotherapeutic agents or other treatment modalities, which outweighs the immune protection effects conferred by HLA-G expression. It is uncertain if there is a causal role of HLA-G molecules in contributing to chemosensitivity in ovarian cancer. Obviously, a large series of multi-institutional study is required to further confirm the preliminary reports as discussed above.

Toward the HLA-G based therapy

The studies as discussed above suggest that HLA-G mediated immune suppression may represent the “Achilles' tendon” of cancer cells and appear as an attractive molecular target for developing new anti-tumor interventions. This feature is particularly important in ovarian cancer as most ovarian cancer patients are at advanced stages whose tumors almost always recur after chemotherapy. Novel therapeutics is urgently needed to improve the clinical management of ovarian cancer patients. Given the fact that the inhibitory effects mediated by HLA-G are mainly through the interactions with lymphocyte Ig-like receptors (Fig. 3), it is likely that blocking the interactions would have clinical benefit in treating HLA-G expressing cancers. Along this line, the antibodies against LIR-1 and LIR-2 are able to restore the proliferating activity in T cells based on *in vitro* allostimulation assays, suggesting the potential of such therapeutic approach [67]. Similarly, anti-FasL antibody which blocks the Fas/FasL pathway can reverse HLA-G mediated inhibition in T cell proliferation [67]. These data support a potential application of using HLA-G blockers, such as HLA-G neutralizing antibodies, soluble recombinant LIR1, LIR-2 and Fas, to abrogate HLA-G mediated inhibitory effects in immune cells (Fig. 3). It could be envisioned that the anti-HLA-G antibodies could also be applied in developing a cancer imaging system to locate tumors in anatomic sites and to monitor tumor load in cancer patients. As HLA-G molecules are cell surface proteins, HLA-G antibody alone or in combination with other antibodies can be used to decorate the surface of liposomes or nanoparticles that are packed with anti-cancer drugs. The reagents are expected to specifically target cancer cells to maximize the therapeutic effects and to minimize the untoward side effects that are frequently associated with anti-cancer reagents.

Summary

Escape from host immune surveillance represents an early and essential step in the development of human neoplastic diseases including ovarian cancer. The fact that HLA-G is frequently upregulated in ovarian serous (Type II) carcinomas supports the view that HLA-G participates in tumor development by executing its immunosuppressant role in tumor microenvironment. The complex immune regulatory networks that are involved in tumor-associated immune suppression indicate that HLA-G does not act alone. In fact, the active cross-talks between tumor cells and immune competent cells through various immunosuppressive cytokines and an array of ligand-receptor interactions create an immunosuppressive microenvironment to

protect ovarian cancer cells from destruction by the immune system [45,46]. Besides the biological significance in immunosuppression, HLA-G expression in ovarian carcinoma has potential clinical applications. sHLA-G may be potentially used as a surrogate tumor-associated marker in blood for early detection of ovarian cancer in high-risk group of patients. Similarly, sHLA-G test can also help in the differential diagnosis of malignant versus benign ovarian neoplasm by analyzing ascites supernatant. Future clinical studies will likely demonstrate the usefulness of HLA-G expression in predicting the clinical outcome in ovarian cancer patients. Finally, the development of HLA-G based target therapy may prove critical in the treatment of ovarian cancer patients. Anti-HLA-G therapy may be used in combination with conventional chemotherapy in ovarian cancer patients who receive cytoreduction surgery to prolong the disease free survival.

Although recent clinical and *in vitro* molecular studies provide new insights into the biological and clinical significance of HLA-G in ovarian cancer development, there are several key questions remained to be answered. For example, there seems a lack of pre-clinical animal model that allows for molecular dissection of immune networks to address if HLA-G plays a causal role in the pathogenesis of ovarian carcinomas. The animal model may take advantages of immune compromised mice that are reconstituted with genetically engineered immune cells and human ovarian tumor xenografts. The *in vivo* animal model would provide a valuable tool for the future study of roles of HLA-G in ovarian cancer and the development of a novel HLA-G targeted therapy in ovarian cancer.

Acknowledgements

This study is supported by a research grant from the National Institute of Health (CA-103937), Bethesda, MD, USA.

References

1. Greaves M. Darwinian medicine: a case for cancer. *Nat Rev Cancer* 2007;7:213–21. [PubMed: 17301845]
2. Hayakawa S. No cancer in cancers: evolutionary trade-off between successful viviparity and tumor escape from the adaptive immune system. *Med Hypotheses* 2006;66:888–97. [PubMed: 16455204]
3. Kovats S, Main EK, Librach C, Stubblebine M, Fisher SJ, DeMars R. A class I antigen, HLA-G, expressed in human trophoblasts. *Science* 1990;248:220–3. [PubMed: 2326636]
4. Shih Ie M. Application of human leukocyte antigen-g expression in the diagnosis of human cancer. *Hum Immunol* 2007;68:272–6. [PubMed: 17400063]
5. Tavassoli, F.; Devilee, P. World Health Organization Classification of Tumors- Pathology and genetics of tumors of the breast and female genital organs. Lyon, France: IARC press; 2003. Tumors of the ovary and peritoneum; p. 114-6.
6. Berek JS, Martinez-Maza O. Molecular and biologic factors in the pathogenesis of ovarian cancer. *J Reprod Med* 1994;39:241–8. [PubMed: 8040839]
7. DuBeau CE. Urinary incontinence management: new questions from old assumptions. *J Am Geriatr Soc* 2001;49:829–30. [PubMed: 11454125]
8. Shih Ie M, Kurman RJ. Ovarian tumorigenesis: a proposed model based on morphological and molecular genetic analysis. *Am J Pathol* 2004;164:1511–8. [PubMed: 15111296]
9. Shih Ie M, Kurman RJ. Molecular pathogenesis of ovarian borderline tumors: new insights and old challenges. *Clin Cancer Res* 2005;11:7273–9. [PubMed: 16243797]
10. Salani R, Kurman RJ, Giuntoli IR, Gardner GJ, Bristow RE, Wang TL, et al. Assessment of TP53 mutation using purified tissue samples of ovarian serous carcinomas reveals a much higher mutation rate than previously reported and does not correlate with drug resistance. *Int J Gynecol Cancer*. 2007In press
11. Crisa L, McMaster MT, Ishii JK, Fisher SJ, Salomon DR. Identification of a thymic epithelial cell subset sharing expression of the class Ib HLA-G molecule with fetal trophoblasts. *J Exp Med* 1997;186:289–98. [PubMed: 9221758]

12. Rouas-Freiss N, Moreau P, Ferrone S, Carosella ED. HLA-G proteins in cancer: do they provide tumor cells with an escape mechanism? *Cancer Res* 2005;65:10139–44. [PubMed: 16287995]
13. Kleinberg L, Florenes VA, Skrede M, Dong HP, Nielsen S, McMaster MT, et al. Expression of HLA-G in malignant mesothelioma and clinically aggressive breast carcinoma. *Virchows Arch* 2006;449:31–9. [PubMed: 16541284]
14. Hansel DE, Rahman A, Wilentz RE, Shih Ie M, McMaster MT, Yeo CJ, et al. HLA-G upregulation in pre-malignant and malignant lesions of the gastrointestinal tract. *Int J Gastrointest Cancer* 2005;35:15–23. [PubMed: 15722570]
15. Singer G, Kurman RJ, McMaster M, Shih IM. HLA-G immunoreactivity is specific for intermediate trophoblast in gestational trophoblastic disease and can serve as a useful marker in differential diagnosis. *Am J Surg Pathol* 2002;26:914–20. [PubMed: 12131159]
16. Singer G, Rebmann V, YC C, Liu HT, Ali SZ, Reinsberg J, et al. HLA-G is a potential tumor marker in malignant ascites. *Clin Cancer Res* 2003;9:4460–4. [PubMed: 14555519]
17. Bukur J, Rebmann V, Grosse-Wilde H, Luboldt H, Ruebben H, Drexler I, et al. Functional role of human leukocyte antigen-G up-regulation in renal cell carcinoma. *Cancer Res* 2003;63:4107–11. [PubMed: 12874014]
18. Urosevic M, Kurrer MO, Kamarashev J, Mueller B, Weder W, Burg G, et al. Human leukocyte antigen G up-regulation in lung cancer associates with high-grade histology, human leukocyte antigen class I loss and interleukin-10 production. *Am J Pathol* 2001;159:817–24. [PubMed: 11549573]
19. Wiendl H, Mitsdoerffer M, Weller M. Hide-and-seek in the brain: a role for HLA-G mediating immune privilege for glioma cells. *Semin Cancer Biol* 2003;13:343–51. [PubMed: 14708714]
20. El-Chennawi FA, Auf FA, El-Diasty AM, El-Daim MA, El-Sherbiny SM, Ali A, et al. Expression of HLA-G in cancer bladder. *Egypt J Immunol* 2005;12:57–64. [PubMed: 16734140]
21. Ye SR, Yang H, Li K, Dong DD, Lin XM, Yie SM. Human leukocyte antigen G expression: as a significant prognostic indicator for patients with colorectal cancer. *Mod Pathol* 2007;20:375–83. [PubMed: 17277760]
22. Amiot L, Le Fric G, Sebti Y, Drenou B, Pangault C, Guilloux V, et al. HLA-G and lymphoproliferative disorders. *Semin Cancer Biol* 2003;13:379–85. [PubMed: 14708718]
23. Rebmann V, Pfeiffer K, Passler M, Ferrone S, Maier S, Weiss E, et al. Detection of soluble HLA-G molecules in plasma and amniotic fluid. *Tissue Antigens* 1999;53:14–22. [PubMed: 10082427]
24. Rebmann V, van der Ven K, Passler M, Pfeiffer K, Krebs D, Grosse-Wilde H. Association of soluble HLA-G plasma levels with HLA-G alleles. *Tissue Antigens* 2001;57:15–21. [PubMed: 11169254]
25. Rebmann V, Regel J, Stolke D, Grosse-Wilde H. Secretion of sHLA-G molecules in malignancies. *Semin Cancer Biol* 2003;13:371–7. [PubMed: 14708717]
26. Gros F, Sebti Y, de Guibert S, Branger B, Bernard M, Fauchet R, et al. Soluble HLA-G molecules increase during acute leukemia, especially in subtypes affecting monocytic and lymphoid lineages. *Neoplasia* 2006;8:223–30. [PubMed: 16611416]
27. Nakayama K, Nakayama N, Jinawath N, Salani R, Kurman RJ, Shih Ie M, et al. Amplicon profiles in ovarian serous carcinomas. *Int J Cancer* 2007;120:2613–7. [PubMed: 17351921]
28. Mouillot G, Marcou C, Zidi I, Guillard C, Sangrouber D, Carosella ED, et al. Hypoxia modulates HLA-G gene expression in tumor cells. *Hum Immunol* 2007;68:277–85. [PubMed: 17400064]
29. Chang CC, Ferrone S. HLA-G in melanoma: can the current controversies be solved? *Semin Cancer Biol* 2003;13:361–9. [PubMed: 14708716]
30. Ibrahim EC, Morange M, Dausset J, Carosella ED, Paul P. Heat shock and arsenite induce expression of the nonclassical class I histocompatibility HLA-G gene in tumor cell lines. *Cell Stress Chaperones* 2000;5:207–18. [PubMed: 11005379]
31. Moreau P, Faure O, Lefebvre S, Ibrahim EC, O'Brien M, Gourand L, et al. Glucocorticoid hormones upregulate levels of HLA-G transcripts in trophoblasts. *Transplant Proc* 2001;33:2277–80. [PubMed: 11377528]
32. Yie SM, Li LH, Li GM, Xiao R, Librach CL. Progesterone enhances HLA-G gene expression in JEG-3 choriocarcinoma cells and human cytotrophoblasts in vitro. *Hum Reprod* 2006;21:46–51. [PubMed: 16210391]

33. Moreau P, Adrian-Cabestre F, Menier C, Guiard V, Gourand L, Dausset J, et al. IL-10 selectively induces HLA-G expression in human trophoblasts and monocytes. *Int Immunol* 1999;11:803–11. [PubMed: 10330285]
34. Urošević M, Willers J, Mueller B, Kempf W, Burg G, Dummer R. HLA-G protein up-regulation in primary cutaneous lymphomas is associated with interleukin-10 expression in large cell T-cell lymphomas and indolent B-cell lymphomas. *Blood* 2002;99:609–17. [PubMed: 11781245]
35. Wagner SN, Rebmann V, Willers CP, Grosse-Wilde H, Goos M. Expression analysis of classic and non-classic HLA molecules before interferon alfa-2b treatment of melanoma. *Lancet* 2000;356:220–1. [PubMed: 10963203]
36. Lefebvre S, Berrih-Aknin S, Adrian F, Moreau P, Poëa S, Gourand L, et al. A specific interferon (IFN)-stimulated response element of the distal HLA-G promoter binds IFN-regulatory factor 1 and mediates enhancement of this nonclassical class I gene by IFN-beta. *J Biol Chem* 2001;276:6133–9. [PubMed: 11087747]
37. Ugurel S, Rebmann V, Ferrone S, Tilgen W, Grosse-Wilde H, Reinhold U. Soluble human leukocyte antigen-G serum level is elevated in melanoma patients and is further increased by interferon-alpha immunotherapy. *Cancer* 2001;92:369–6. [PubMed: 11466692]
38. Rouas-Freiss N, Moreau P, Menier C, Carosella ED. HLA-G in cancer: a way to turn off the immune system. *Semin Cancer Biol* 2003;13:325–6. [PubMed: 14708712]
39. Moreau P, Mouillot G, Rousseau P, Marcou C, Dausset J, Carosella ED. HLA-G gene repression is reversed by demethylation. *Proc Natl Acad Sci U S A* 2003;100:1191–6. [PubMed: 12552087]
40. Boucraut J, Guillaudeux T, Alizadeh M, Boretto J, Chimini G, Malecaze F, et al. HLA-E is the only class I gene that escapes CpG methylation and is transcriptionally active in the trophoblast-derived human cell line JAR. *Immunogenetics* 1993;38:117–30. [PubMed: 7683306]
41. Mouillot G, Marcou C, Rousseau P, Rouas-Freiss N, Carosella ED, Moreau P. HLA-G gene activation in tumor cells involves cis-acting epigenetic changes. *Int J Cancer* 2005;113:928–36. [PubMed: 15514928]
42. Chang CC, Murphy SP, Ferrone S. Differential in vivo and in vitro HLA-G expression in melanoma cells: potential mechanisms. *Hum Immunol* 2003;64:1057–63. [PubMed: 14602236]
43. Ibrahim EC, Guerra N, Lacombe MJ, Angevin E, Chouaib S, Carosella ED, et al. Tumor-specific up-regulation of the nonclassical class I HLA-G antigen expression in renal carcinoma. *Cancer Res* 2001;61:6838–45. [PubMed: 11559559]
44. Onno M, Le Friec G, Pangault C, Amiot L, Guilloux V, Drenou B, et al. Modulation of HLA-G antigens expression in myelomonocytic cells. *Hum Immunol* 2000;61:1086–94. [PubMed: 11137211]
45. Khong HT, Restifo NP. Natural selection of tumor variants in the generation of “tumor escape” phenotypes. *Nat Immunol* 2002;3:999–1005. [PubMed: 12407407]
46. Mocellin S, Wang E, Marincola FM. Cytokines and immune response in the tumor microenvironment. *J Immunother* 2001;24:392–407.
47. Urošević M, Dummer R. HLA-G and IL-10 expression in human cancer—different stories with the same message. *Semin Cancer Biol* 2003;13:337–42. [PubMed: 14708713]
48. Dummer R, Willers J, Kamarashev J, Urošević M, Dobbeling U, Burg G. Pathogenesis of cutaneous lymphomas. *Semin Cutan Med Surg* 2000;19:78–86. [PubMed: 10892708]
49. Asadullah K, Docke WD, Volk HD, Sterry W. Cytokines and cutaneous T-cell lymphomas. *Exp Dermatol* 1998;7:314–20. [PubMed: 9858133]
50. LeMaoult J, Le Discorde M, Rouas-Freiss N, Moreau P, Menier C, McCluskey J, et al. Biology and functions of human leukocyte antigen-G in health and sickness. *Tissue Antigens* 2003;62:273–84. [PubMed: 12974794]
51. Rizzo R, Melchiorri L, Stignani M, Baricordi OR. HLA-G expression is a fundamental prerequisite to pregnancy. *Hum Immunol* 2007;68:244–50. [PubMed: 17400059]
52. Sher G, Keskin-tepe L, Fisch JD, Acacio BA, Ahlering P, Batzofin J, et al. Soluble human leukocyte antigen G expression in phase I culture media at 46 hours after fertilization predicts pregnancy and implantation from day 3 embryo transfer. *Fertil Steril* 2005;83:1410–3. [PubMed: 15866577]

53. Noci I, Fuzzi B, Rizzo R, Melchiorri L, Criscuoli L, Dabizzi S, et al. Embryonic soluble HLA-G as a marker of developmental potential in embryos. *Hum Reprod* 2005;20:138–46. [PubMed: 15498780]
54. Rebmann V, Switala M, Eue I, Schwahn E, Merzenich M, Grosse-Wilde H. Rapid evaluation of soluble HLA-G levels in supernatants of in vitro fertilized embryos. *Hum Immunol* 2007;68:251–8. [PubMed: 17400060]
55. Jurisicova A, Casper RF, MacLusky NJ, Mills GB, Librach CL. HLA-G expression during preimplantation human embryo development. *Proc Natl Acad Sci U S A* 1996;93:161–5. [PubMed: 8552596]
56. Pfeiffer KA, Fimmers R, Engels G, van der Ven H, van der Ven K. The HLA-G genotype is potentially associated with idiopathic recurrent spontaneous abortion. *Mol Hum Reprod* 2001;7:373–8. [PubMed: 11279300]
57. Aldrich CL, Stephenson MD, Karrison T, Odem RR, Branch DW, Scott JR, et al. HLA-G genotypes and pregnancy outcome in couples with unexplained recurrent miscarriage. *Mol Hum Reprod* 2001;7:1167–72. [PubMed: 11719594]
58. Creput C, Durrbach A, Menier C, Guettier C, Samuel D, Dausset J, et al. Human leukocyte antigen-G (HLA-G) expression in biliary epithelial cells is associated with allograft acceptance in liver-kidney transplantation. *J Hepatol* 2003;39:587–94. [PubMed: 12971970]
59. Rajagopalan S, Long EO. A human histocompatibility leukocyte antigen (HLA)-G-specific receptor expressed on all natural killer cells. *J Exp Med* 1999;189:1093–100. [PubMed: 10190900]
60. Ponte M, Cantoni C, Biassoni R, Tradori-Cappai A, Bentivoglio G, Vitale C, et al. Inhibitory receptors sensing HLA-G1 molecules in pregnancy: decidua-associated natural killer cells express LIR-1 and CD94/NKG2A and acquire p49, an HLA-G1-specific receptor. *Proc Natl Acad Sci U S A* 1999;96:5674–9. [PubMed: 10318943]
61. Cosman D, Fanger N, Borges L, Kubin M, Chin W, Peterson L, et al. A novel immunoglobulin superfamily receptor for cellular and viral MHC class I molecules. *Immunity* 1997;7:273–82. [PubMed: 9285411]
62. Colonna M, Navarro F, Bellon T, Llano M, Garcia P, Samaridis J, et al. A common inhibitory receptor for major histocompatibility complex class I molecules on human lymphoid and myelomonocytic cells. *J Exp Med* 1997;186:1809–18. [PubMed: 9382880]
63. Rajagopalan S, Fu J, Long EO. Cutting edge: induction of IFN-gamma production but not cytotoxicity by the killer cell Ig-like receptor KIR2DL4 (CD158d) in resting NK cells. *J Immunol* 2001;167:1877–81. [PubMed: 11489965]
64. Clements CS, Kjer-Nielsen L, McCluskey J, Rossjohn J. Structural studies on HLA-G: implications for ligand and receptor binding. *Hum Immunol* 2007;68:220–6. [PubMed: 17400055]
65. LeMaout J, Zafaranloo K, Le Danff C, Carosella ED. HLA-G up-regulates ILT2, ILT3, ILT4, and KIR2DL4 in antigen presenting cells, NK cells, and T cells. *Faseb J* 2005;19:662–4. [PubMed: 15670976]
66. Davidson B, Elstrand MB, McMaster MT, Berner A, Kurman RJ, Risberg B, et al. HLA-G expression in effusions is a possible marker of tumor susceptibility to chemotherapy in ovarian carcinoma. *Gynecol Oncol* 2005;96:42–7. [PubMed: 15589578]
67. Naji A, Durrbach A, Carosella ED, Rouas-Freiss N. Soluble HLA-G and HLA-G1 expressing antigen-presenting cells inhibit T-cell alloproliferation through ILT-2/ILT-4/FasL-mediated pathways. *Hum Immunol* 2007;68:233–9. [PubMed: 17400057]

Fig. 1. HLA-G expression in trophoblastic cells from a placenta (A) and in tumor cells from a high-grade ovarian serous carcinoma (B). Cell surface and cytoplasm show an intense HLA-G immunoreactivity (brownish chromagen deposits), indicating a high level of HLA-G expression. The stromal cells in both placenta and ovarian cancer do not express any detectable level of HLA-G immunoreactivity.

A. mRNA levels by Q-PCR

B. Protein levels by IHC

Fig. 2.

Comparison of HLA-G expression in ovarian cancer and other cancer types. A: HLA-G expression is analyzed by quantitative real-time PCR. B: HLA-G expression is assessed by immunohistochemistry. The X-axis indicates that percentage of tumors with HLA-G expression within each tumor type analyzed. Ovarian cancer is one of the cancer types that most frequently express HLA-G.

Fig. 3.

The biological effects of HLA-G expression in ovarian cancer cells. HLA-G is expressed in an ovarian tumor cell (left) and both membrane-bound and secreted (shed) HLA-G can interact with their receptors on the surface of a variety of immune cells including B- and T-lymphocytes, natural killer (NK) cells, dendritic cells and macrophages. Numerous of such immune cells are present in ovarian carcinoma tissues as well as in tumor ascites. The HLA-G receptors include the leukocyte Ig-like receptors, LIR-1 and LIR-2, that are expressed by all the immune effector cells while the killer-Ig-like receptor, KIR2DL4, is preferentially expressed by the NK cells. Upon HLA-G binding, the receptors on the immune cell surface initiates a cascade of signals that reduce cellular proliferation and compromise the immunological functions on the immune cells (summarized in the right column). (+): positive effects; (-): negative effects. Dimerization of HLA-G molecules can enhance the binding affinity between HLA-G and LIR1/LIR2 by stabilizing the protein complex. The factors that enhance HLA-G expression in ovarian cancer and the potential targets that can be blocked to restore local immunity in tumors are also listed in the box.