As to the general state of the patient during each of these conditions of treatment, restriction to an antidiabetic dietary produced improvement in thirst and mental activity. So long as the quantity of codeine was limited to six grains daily, this improvement was maintained; but when nine grains, and even more, when fifteen grains were being taken daily, the appetite failed, she became listless and apathetic, vertigo was occasionally experienced, and the patient remained for a considerable part of each day asleep in bed. The addition of atropine to the codeine did not produce any improvement, but rather added to the discomfort by impairing vision; and even when only one-sixtieth of a grain was being taken thrice daily, the pupils became slightly dilated. When codeine and atropine were stopped, and a restricted dietary alone adopted, the health was not improved. The symptoms referred to became, indeed, worse, and prevented a prolongation of the period of restricted diet without medicine to the extent that seemed desirable before a new plan of treatment was adopted. When opium was now given a marked improvement occurred, but the larger doses caused some drowsiness during the day. With one grain daily of hydrochlorate of morphine the condition of the patient became more satisfactory. The drowsiness soon disappeared, the appetite improved, and she became sufficiently active to engage in ward work. Constipation of the bowels was not produced by any of the medicinal agents employed.

A consideration of these averages seems to show that, under a daily administration of one grain of hydrochlorate of morphine, the quantity of fluids drunk, and of urine, urea, and sugar voided, was rather less than when three grains of opium, and decidedly less than when fifteen grains of codeine were being taken. In three other cases in which I have instituted a comparison between these substances in diabetes mellitus, morphine also showed a marked, though not so great, superiority over codeine. After this note had been prepared, I have seen a recent paper by Dr. Bruce, of London, in which similar results were obtained in two very carefully observed cases. So far as I know, also, the favour with which codeine is regarded in this disease has not been supported by any observations calculated to show its value relatively to opium or morphine so clearly as in the cases to which I have referred.

The evidence, therefore, seems to indicate that codeine is a less powerful remedy in diabetes than either opium or morphine, and to confirm the view that in its therapeutic value it ranks as a weak or diluted morphine.

The conclusion receives an importance (no doubt a subsidiary one) from the circumstance that codeine is about three times as expensive a substance as morphine. The great demand for it has led to its being manufactured from morphine so largely, that probably one-fourth of the codeine in the market is an artificial substance. When we consider the large doses that are required in diabetes mellitus, and the generally protracted duration of this disease, we are, I think, justified in asking for more clear evidence of its superiority over morphine than has as yet been produced.

NOTES ON THE VALUE OF SOME NEW DRUGS.¹

By JAMES F. GOODHART, M.D., F.R.C.P.,

Physician to, and Lecturer on Pathology at, Guy's Hospital.

THE drugs that I purpose to say a word or two about to-night are paraldehyde, urethan, antipyrin, and pilocarpin; and perhaps, in dealing with antipyrin, I may say a word or two upon my experience with other antipyretic drugs. In these go-ahead days I may be taken to task for calling these new drugs, and I had some doubt how to name my paper; but if I cannot call them new I most certainly cannot call them old, and no equivalent to the "subacute" of medicine came ready to my hand. Let me also say that I do not propose to treat of these drugs from any other than my own standpoint; it is simply my object to say in what I have found them useful, and ask you for such experience as you may have of the same agents in similar or other fields. Thus I shall hardly, if at all, discuss the actions of the remedies, nor is it my object to give even a summary of the diseases in which they have been used or recommended.

Now, first of all, *paraldehyde*. It has been *out*, so to speak, for several years, but I do not think it is at all in common use. I am rather fond of it, and have given it frequently now certainly for three or four years, and that means in a large number of cases. Originally recommended, I believe, as a hypnotic, one that combined the advantages, to the exclusion of the disadvantages, of chloral and bromide of potassium, I tried it first of all in cases of insomnia. I do not consider it at all a reliable drug for that state of things; but it is a drug of unquestionable value in the restlessness and cardiac asthma so often seen in aortic disease, and in the labouring dilated heart of chronic Bright's disease. I well remember the first case in which I gave it some years ago, a man who had long been subject to gout, and who had a granular kidney and a large dilated heart. His distress was extreme, and I had exhausted all the old-fashioned remedies (than which few are better), and many new, including caffeine, and with a marked condition of Cheyne-Stokes's breathing, he had all the aspect of speedy dissolution. As a last resource I gave him half-drachm doses of paraldehyde, with marked relief to the end of the case, which occurred about a fortnight later. I have also found it occasionally useful in the headache of renal disease. In two cases, where this was especially troublesome in the early morning, it was taken on and off for many months, with great relief. As a general rule, nitro-glycerine is the more reliable remedy for the headache of Bright's disease, but the symptom is often a troublesome one, and paraldehyde may well be kept in memory for occasional cases. But there is another class of cases in which I think it of even more use. It was recommended some few years ago, I think by Dr. Strachan, for certain cases of mania, and some three years or so ago I had to deal with a very troublesome case of alcoholic delirium. The patient was a very free drinker, had had delirium tremens twice before, and was now in a low, muttering, sleepless condition, which nothing seemed to quiet or control. He had been in a semi-comatose state already for five days. He was quite unheeded of anyone about him, talked in an unintelligible mutter, and his muscles were tremulous, flabby, and wasted. As his wife was his equal, if not his better half, in drinking capacity, he was sent into Guy's Hospital, and for eighteen days he remained in much the same state. During that time his treatment consisted of free aperients, iodide and bromide of potassium, opium, and latterly of subcutaneous injections, first of $\frac{1}{10}$ grain, and then of $\frac{1}{20}$ grain of hyoscyamine; none of them seemed to do him any good. I then tried half a drachm of paraldehyde three times a day, and within a day or two he was much clearer in his mental condition, and before a week was out he was practically well. He remained somewhat feeble-minded for some little time, but he quite recovered eventually. Since then I have given it in similar cases several times. Whether it acts as a stimulant or a sedative I do not know, but it certainly sometimes quiets the delirium and clears the mental faculties. My friend Mr. Wornum lately ordered it in another such case, but it was only given in ten-minim doses. The general condition improved decidedly, but the mental condition still remains cloudy. However, the patient has come out of a semi-comatose state, and is now sitting up, although she still suffers from delusions.

Another case of what I think may be called a similar ailment was treated in a similar manner. A young fellow, aged 27, had typhoid fever badly; he became excessively emaciated, and not long after the fever left him he went off his head; he was very delirious and troublesome. Free stimulation failed to restore him, and he was, therefore, ordered half-drachm doses of paraldehyde. This was on the 8th of the month. The stuff was so nasty, he said, that he would take no more, and when I again saw him, on the 11th, it had not been persevered with. He then began it regularly, and within four or five days he was much clearer and less delirious, and at the end of a week he was practically well.

I suppose that the state of malnutrition after a prolonged attack of typhoid fever is not very unlike that produced by alcoholic poisoning, and, if so, might be expected to be benefited by the same remedy. The main objection to paraldehyde is its strong taste. I have usually given it with almond mixture, but it is, after all, not so very nauseous combined with some tincture of orange and syrup and water. The patient taking paraldehyde emits a very peculiar and characteristic odour, which is very tenacious, and I have known it to linger about the person two or three days after the drug has been discontinued.

Of *urethan* I have not much to say, but that little is in its favour. I have used it in a large number of cases as a hypnotic, and although it often fails, I think it is one of the best remedies

¹ Read before the North London District meeting of the Metropolitan Counties Branch at Tottenham.

of the kind since the introduction of chloral. I give it in fifteen-grain doses in water, and have often found it useful; and once I remember it was very so in relieving the headache of early typhoid. A newer hypnotic still is sulphonal, but my experience of this as yet only extends to one case.

Of *pilocarpin* I have only two points that I wish to make. I have, of course, used it quite commonly in Bright's disease, and still do as a usual thing, and it seldom fails to induce a copious diaphoresis; but one day it occurred to me to order a third of a grain as a subcutaneous injection in a case of chronic jaundice with intense itching. It occurred to me that a drug which so uniformly was productive of speedy diaphoresis must profoundly modify the functions of the skin for the time being, and might in doing so relieve the itching which has hitherto, in my experience, defied treatment. It did so quite beyond my expectation, and kept the patient comfortable until she died. In the next case it was equally successful. The patient had one-third of a grain injected subcutaneously many times, and always with this result, that the first twenty-four hours he was quite free, the second he was fairly free, and the third day he was getting bad again, and the dose had to be repeated. I next tried it on an old lady who suffers from very frequent attacks of gall-stones. She is equally decided as to the relief she obtains, but her pleasure is to some extent marred by her skin being for the time converted, as she describes it, into the similitude of that of a laundrywoman. I have had six cases in all, and in none has it failed; and when we consider that really there is nothing that can be relied upon to relieve this most distressing feature of jaundice, I hope the suggestion may prove of service. The other class of cases in which it seems to me to be occasionally of value is where the lightning pains of locomotor ataxy are severe. I have two or three times found it useful in these cases when many other means had been tried and failed.

Of *antipyrin* I am almost afraid to speak, for just now it is such a fashionable drug that it would well-nigh be easier to say for what diseases it had *not* been recommended than to enumerate those for which it had. But it does seem that it is likely to achieve a permanent name for abnormal nerve discharges of various kinds, and chiefly those that have been, I think happily, designated as the "paroxysmal neuroses." For instance, there can, I think, be no doubt that it is a very valuable means of combating some migrainous headaches. From what others have told me, and from what I have occasionally seen myself, it would appear to be a remedy of value in some cases of dysmenorrhœa—cases which might, I think, not unfairly be designated as pelvic catamenial neuralgia, if the gynaecologists will forgive me for venturing to have an idea outside my own lines. And fever, too, though you may not, perhaps, have looked at it in that light, is surely much of it a paroxysmal neurosis; and as that light is, I think, a leading one, let me follow it for a moment. Pyrexia is the medical Aunt Sally just now. Up goes a temperature of 103°, and down goes a dose of antipyrin or one of its congeners; and no doubt in most cases the shot tells and down goes the temperature, but it speedily goes up again, and, like the old lady I have mentioned, is soon ready for the next shot. And what I have learnt from the study of pyrexia is this, that it is much more special to the individual than it is to the disease. I am shown a chart of a doubtful case, and asked if it is not like typhoid. But what is a typical typhoid chart? There is little else typical than its duration; and the proper duration of typhoid, as everyone knows, is as often honoured by the breach as by the observance; and taken, as temperature charts are usually nowadays, every three or four hours, or oftener, there are few charts typical of anything. And I maintain that, given a certain uniformity in length of certain fevers, according to the life-history of particular organisms, the rest of the pattern is due to the individual resistance or want of it on the part of the tissues and centres provoked. Well, then, who does not know that fever (the tendency to pyrexia) runs in families? Only yesterday I heard a mother say so. All her children are noted by their doctor as becoming sharply pyrexial in all the minor ailments and disturbances of daily life. We can all endorse the truth of that observation. Fever runs in families, then, which is only another way of saying that it is the individual you must take stock of in treating the malady; and I take it that that is in some sort evidence of the predominant influence of the nervous system in the regulation of the production and the discharge of heat, and I would certainly hold that pyrexia is in many cases a neurosis. It is from this point of view that there seems, I think, the strong probability that drugs will come to be of value in checking fevers

by acting upon and restoring the tone of the diseased nervous centres.

I throw this out for discussion. According to my own experience, I doubt whether we have got far yet in this direction. We require to know much more than we as yet know about the particular fault in each case that needs a remedy. By-and-bye, no doubt, we shall be able to tell these and apply now one antipyretic, now another, according to the special need; but it is hardly so as yet. My own experience would lead me to say that there are two kinds of fever: one that does harm to the patient, the other associated with no ill result. In the latter, little good results from the administration of antipyretics; and in the other class my own impression is that little good results either; in both the temperature is depressed for a time by the remedy, but it soon rises again, and the duration of the disease is not appreciably influenced thereby.

Antipyretics have been advocated by some in enthusiastic terms for the sudden and sharp febrile states that are met with in childhood, and I am far from saying that in such they may not be of use. I believe they may be of great use in that special condition known as hyperpyrexia—or rather in the states which threaten it—because in it the rise is very sudden, the attack is of the nature of a sudden stormblast, and the tissues may be irreparably withered by it. In the same way in children sudden fever may do irreparable damage to their succulent tissues, and a dose of antifebrin or antipyrin may avert this. Nevertheless, it will occur to all of us that the attacks in which they seem to be so beneficial are just those whose nature is to subside as suddenly as they come, and I do not think that the evidence they afford is as yet conclusive in favour of these remedies.

But to go back to antipyrin in its power to depress abnormally high temperatures and to dissipate a migraine or pelvic neuralgia, we have evidence of some powerful influence on the nervous centres, and it is not far thence to the question, Might it be useful in other neurotic states? And it has been tried of late in two very troublesome ailments, namely, chorea and whooping-cough. I have no personal experience to relate as regards the latter, but it has been spoken of very favourably by Sonnenberger,² Wendt,³ and Genser.⁴ Sonnenberger has tried it in about seventy cases, the dose varying from one-seventh of a grain in very young children to fifteen grains in adults three times a day. Wendt followed the former writer's suggestions, and states that the drug served him better than any other; he claims no cures, but it favours an easy course to final recovery, a mitigation of the paroxysms and a reduction in their number, and certainly a freedom from complications. Genser also asserts that antipyrin always diminishes the number of the coughs in the twenty-four hours, and controls the intensity; the duration of the disease seldom extended over twenty-four days—a very good record, as all will allow, if it be borne out by other observers. The mean dose was a grain and a half a day for each year of age; but I should say myself that the drug might be pushed to the extent of 1½ grain per year three or four times in the twenty-four hours if necessary.

As regards chorea, I can speak—to some extent and tentatively—favourably. I have given it in the last five cases that have come under my observation. In four the result may, I think, be considered to have been good. Thus a girl, aged 8, was admitted for her fourth attack of chorea. It was general and rather severe, and she had mitral disease (probably contraction) also. She was put upon two-grain doses of antipyrin three times a day, and after five days it was noticed that there was a marked improvement in the choreic condition, the face muscles especially being quieter. The dose of the drug was then increased to four grains three times a day. She continued to improve for ten days, and by that time the movements had lost all their violence, when the drug was discontinued for arsenic. The minor movements remained for some time, as is usually the case with any treatment. She left the hospital practically well after seven weeks.

In another case a boy of 12, who had had rheumatic fever twice and chorea on and off for a long time, was admitted to Guy's Hospital with marked but not violent chorea of face and extremities and aortic regurgitation. He was put upon five-grain doses of the drug three times a day on the 24th of the month; on the 29th the movements were noticed to be less marked, and on

² *Deutsch. Med. Wochenschrift*, 1887, 14.

³ *Philadelphia Med. News*, June 2nd, 1888.

⁴ *Allgem. Wiener Med. Zeit.*, 1888, 15.

the 31st they were not noticeable, and he remained well afterwards. He took the drug for twelve days altogether, and it was then discontinued because of an erythematous eruption that appeared. He was in the hospital three weeks.

A third case, a girl of 8, under the care of Dr. Cooper, of Rotherhithe—a very bad case indeed, associated with much wasting, imbecility, paralysis, and pericarditis—was given also five grains three times a day. She took it for eight days, and was by that time so much improved that it was discontinued for iron and digitalis. The improvement began within forty-eight hours after the commencement of the antipyrin.

A fourth case, now in the Evelina Hospital, a girl of 9½, came in on November 26th with general rather violent chorea. I gave her five-grain doses every six hours, and on the 29th increased it to ten grains every six hours. On December 3rd she was lying in bed hardly choreic. In this case, however, although the chorea was manifestly better, she suffered from antipyrin poisoning. After three or four days a copious measly rash appeared, with high fever and albuminous urine, and then it was necessary to stop the drug.

The other case, under the care of Dr. Steele Perkins, of Streat-ham, was made worse by the treatment, and it was given up after ten days. I shall not indulge in any exuberant anticipations, but, remembering how very troublesome chorea is, I think there is enough evidence at hand to make us believe that, at any rate in some cases, antipyrin may prove of service. Legroux and Dupré⁵ have given it in twenty cases in doses of about seven grains five or six times a day, and they conclude that it is one of the most certain and harmless remedies for chorea, the improvement usually commencing in four to six days.

Before closing, perhaps I may be allowed to add a word not upon a new remedy, but upon an uncommon method of administration of an old one. I refer to the subcutaneous injection of *strychnine*. There is, of course, nothing novel in this. Nevertheless, I am under the impression that strychnine is not often administered in this fashion. Dr. Habershon has advocated its use thus in cases of cardiac failure, and during the past year I have had three troublesome cases of more or less general paralysis that have, for such cases, rapidly improved by its means. One was a case of peripheral neuritis due to alcohol and lead-poisoning combined. He was in a very tremulous condition when admitted. Many of his muscles were much wasted. He had a good deal of pain in the course of various nerves, and he was both sleepless and demented. Various drugs were administered, and he was assiduously galvanised, but he became decidedly more powerless. By-the-bye he had a subcutaneous injection of hydriodate of hyoscine, first $\frac{2}{10}$ of a grain and then $\frac{1}{10}$, and this certainly seemed to quiet his nocturnal delirium, and procure him sleep. On November 4th I ordered him $\frac{1}{10}$ grain of strychnine to be injected hypodermically twice daily. On the 14th it was increased to $\frac{2}{10}$, and by the 31st he was taking $\frac{3}{10}$ twice daily. By this time it was noted that the movements of the patient's hands and arms were much more lively, and there was a gain in the grasping power of both hands. From this time he made continuous improvement. The injection was continued for another month, and was then omitted, chiefly because it caused, or he thought it did, severe pain in his ulnar nerves, which, he insisted, came on soon after the injection, and caused him great misery for several hours afterwards. He had had this pain before the commencement of the injections, and my clerk, Mr. McIlwaine, had doubts about the reality of the cause, but the man was very positive, and I think he was very probably correct. His ulnar nerves were clearly those most affected by his disease, and it is fair, therefore, to suppose that the remedy acted upon them in excess.

A second case was one of general paralysis, dysphagia, diaphragmatic palsy, and reaction of degeneration in the muscles in a young man aged 27. All the history that he could give was that, being a carman, and of late carrying sacks of coal, he had, a month before his admission, felt generally out of sorts and easily fatigued. A week later his finger-tips became numb, and then his toes, and gradually he lost all power in his legs, and then in his arms. He was at first given iodide and perchloride of mercury, and he steadily got worse. Then he had three-minim doses of liq. strychnine given internally; but, as he was steadily going down hill, and was swallowing and breathing so badly that I quite feared for his life, I changed the internal administration of strychnine to the hypodermic method after a week. He had

$\frac{2}{10}$ grain injected night and morning. This was commenced on the fifth, and on the ninth the report stated that there was a marked improvement in the strength of the right arm, and there was considerable improvement in sensation. By the thirteenth he was beginning to have slight control over his legs. His pains, before rather troublesome, were not so severe, and the patient seemed altogether better; and he rapidly improved from that time, so that at the end of a month he was able to get up and about feebly. He is now, many months later, quite well. It might be added that the real nature of the case still remains obscure to me. The completeness of the paralysis, coupled with his occupation, and a doubtful thickening and prominence over the second cervical vertebra, made me think at first of some cervical pachymeningitis, but I subsequently inclined more to the diagnosis of peripheral neuritis, albeit the cause of it could not even be suggested.

I must only inflict one other case upon you. It is that of a man aged 41, who was admitted for loss of power in his right leg, and incontinence of urine. He came in with these symptoms, and there was, on careful testing, some loss of sensation in the opposite leg. His bladder was running over. This and the paralysis made one suppose that there was disease in the cord, rather than outside it, and I feared he had some transverse myelitis, worse on the right side. He had not been in the hospital long when he lost the use of the other leg. He had at first some iodide of potassium and perchloride of mercury. But not finding any marked improvement after a fortnight, I changed it to two-minim doses of the liquor strychniae of the *British Pharmacopæia*, twice a day, and at the end of five weeks he was up and walking about the ward. He is getting about fairly well now, with some clonus and exaggerated reflexes on both sides; but his chief trouble is frequent micturition, possibly due to some slight vesical irritation still remaining.

AN ADDRESS

ON A

USEFUL TECHNIQUE IN REMOVING TUBERCULAR DISEASE FROM THE HIP-JOINT.

*Delivered before the Brighton Medical and Chirurgical Society,
December 6th, 1888.*

By ARTHUR E. BARKER, F.R.C.S.,

Surgeon to University College Hospital; Hunterian Professor of Surgery and Pathology, Royal College of Surgeons.

MR. PRESIDENT AND GENTLEMEN,—In the short communication which I have now the honour to lay before you it will be necessary to start by assuming that I need not plead here for the now almost universally received doctrine of the identity of scrofula with tubercular disease; nor need anything special be said of the details of that irritative hyperplasia induced by the presence of the bacillus which gives rise to the production of those bodies generally known as tubercles. If I may be allowed to say so, I have alluded to these points in sufficient detail to define my own position in my Hunterian Lectures at the Royal College of Surgeons last June. But as the procedure now to be described is the outcome of some deductions put forward in those lectures, perhaps I may be pardoned if I briefly refer to them.

In the first place, the identity of the tubercular processes met with in joints and those observed in other organs was made out. In the hip the tubercular hyperplasia was shown to begin primarily either in the young tissue at the junction of the neck and head of the femur, or in the same tissue at the lines of union between the three portions of the acetabulum, or finally in the synovial membrane of the joint. Starting in any one of these situations, it may spread to any or all of the other parts of the articulation.

Wherever it makes its appearance, tubercle in joints is subject to the same changes as in other parts of the body, such as the brain, the lungs, the kidney, or the lymphatic glands. In the first place, if the vitality of the individual or affected tissue be

⁵ *Revue Mens. des Mal. de l'Enf.*, March, 1888.