1885 has been at school, not only in a particularly healthy part of the country, but also under a gentleman who loyally carries out the great principle of equal culture for the body as well as for the mind. The lad is not only good at most games, but also an excellent shot at a rabbit. While C. C. M.'s eyes have been thus improving, it is interesting to note that his head has increased an inch and a half in circumference, and doubtless the depth of his orbital fossæ and his eyeballs have grown in an equal proportion. He has never used his spectacles except in school hours, and now frequently reads ordinary print without them.

This treatment was based on the opinion that hypermetropia is the normal, or at any rate the common, condition of children's eyes; but that, if healthy growth and development of all parts of the body progress in regular order, hypermetropia gives way to emmetropia or normal vision by the time a boy has reached his sixteenth or eighteenth year. On the other hand, children reared in our large cities are too often defective in their tissue-forming power; not a few are rachitic to a greater or less extent. Many of these children are almost from infancy set to work in gloomy rooms. Conditions such as these, with perhaps faulty food, that is defective in tissue-forming elements, especially in phosphate of lime, tend to prevent proper development, and therefore to perpetuate the hypermetropia. The use of glasses in such cases is, however, only palliative, and, I think, tends to obviate the necessity for growth of the sclerotic and cornea, and therefore may rather fix the formation of the child's eye on into manhood, especially if glasses are constantly worn. As I have before remarked, we should, if possible, relieve such children from work and attend to their surroundings. But, supposing a child has reached his eighth year and has difficulty in his studies on account of hypermetropia, we may order him glasses so as to enable him to work, but he should only wear them when in school; and it is well, if practicable, to reduce the power of his lenses about every two years until the lad has ceased growing, when we can expect no further improvement. I am convinced that treatment of this kind may in some cases lessen the degree of hypermetropia, if it cannot completely overcome it. Hypermetropia is obviously a growing evil, due, it may be, in some measure, to the causes to which I have referred.

ON RABIES:

ITS

TREATMENT BY M. PASTEUR,

AND ON

THE MEANS OF DETECTING IT

IN SUSPECTED CASES.

Read before the Epidemiological Society, February 13th, 1889.

By VICTOR HORSLEY, B.S., F.R.S.,

Surgeon to the National Hospital for the Paralysed and Epileptic: Assistant-Surgeon to University College Hospital: Professor of Pathology in University College, London: Professor Superintendent of the Brown Institution.

THE lecture this evening being on the subject of M. Pasteur's work, it is fitting that what I have to say should be preceded by a brief sketch of the pathology of rabies, since it is owing to his extraordinarily brilliant investigations that we are now rejoicing in full possession of such particulars as make completely clear what till three years ago was somewhat obscure and apparently contradictory. Moreover, I could not treat (as I wish to do for the sake of time) some points by simply scheduling the results of investigations unless I supplied you with the connecting facts.

Rabies is an acute specific disease, the virus of which behaves so exactly like a micro-organism that there can be no doubt that it is indeed one. Micrococci have been described by all observers as occurring in such numbers and position as to suggest a causal nexus, but with the exception of a single doubtful observation by Fol, these have never been isolated in cultivation and proved to be pathogenic. While, therefore, henceforth speaking of the

cause of the disease as a virus, I wish to be understood as considering it established that that virus is a microbe.

We will now proceed to consider the conditions under which the virus at present exists and from time to time produces its destructive effects. Like most pathogenic organisms, it is transmitted from one living animal to another living animal; but if exposed to drying, extremes of temperature, absence of pabulum (probably in rabies necessarily living tissue), it rapidly becomes inert, that is, dies. The practical application of this fundamentally important fact is obvious, namely, that if we could at one blow destroy all animals infected by the disease the malady would immediately become extinct. (I need hardly say that no case of asserted "spontaneous" generation of the disease bears investigation.) It is a fortunate thing that this rapid destruction of the virus should evidently occur, since as it frequently happens that the saliva of a rabid dog is dropped upon the streets, etc., the prevalence of the infection would be much greater than it now is. The virus attacks all animals, is more deadly in its action upon young individuals, and seems to acquire a specially virulent quality during its development in the cat and wolf. Insistence upon the first of these points is important, as in a recent trial the counsel for the plaintiff asserted that horses could not take the disease, and the judge appeared to acquiesce, although every textbook on the subject describes such cases. In 1887, 4 horses, 11 cattle, 8 sheep and swine were returned as affected with the disease in England, and in the same year 99 cattle, 13 horses, 1 mule, 21 sheep and swine were affected in France. No doubt these figures are much smaller than the actual facts, inasmuch as failure to recognise the disease has lately been shown to occur more frequently than is desirable. In all cases the natural mode of transmission of the virus from one animal to another is by subcutaneous inoculation; for example, by being bitten or by the reception of the saliva on to a recent scratch or wound.

I now come to that most important point, the incubation period. Upon this naturally hangs a great deal, and it must therefore be considered at some length. First, let us see what accurately (that is, experimentally) measured, is the universal incubation period, that is the period elapsing between the inoculation and the appearance of (a) excitability (if the disease assume the maniacal form); (b) dulness (if the paralytic form).

The shortest time is 6 to 8 days if the inoculation be subdural; 8 to 9 days is also not infrequent; but while 7 days is an accurate mean, that, though of much practical interest, need not concern us specially for the present. We will take the minimum of 6 days.

I have said that this has been found with subdural inoculation. Now, cases of such extraordinarily short incubation period have been stated to occur in young children from bites upon the face. On account of the age and part attacked this is obviously quite possible, and not only is credence to be given to such a statement, but this interval also serves as a critical point in testing the value of M. Pasteur's treatment. Any case submitted to treatment after 6 days had elapsed should clearly fall under the category of "possibly too late." And, strictly speaking, such cases should be placed in a class by themselves, the death-rate of which would be specially high. That this most particularly concerns patients in this country attacked by a dog certified by a veterinarian to be rabid is seen in the fact that of the few English patients who have died after the Pasteurian treatment, most belong to the "too late" class. Such a case, for instance, was that of Lord Doneraile, who arrived at Paris 11 days after the bite, and who refused to go through the treatment ordered.

The next incubation period of interest is the mean for subcutaneous inoculations, as, for instance, through bites received in the street. The evidence is nearly unanimous in placing this at 6 weeks.

Finally, the maximum period is one which of course we are all extremely desirous to see determined, since its uncertainty constitutes the most painful aspect of the torturing anxiety from which these patients suffer. Unhappily, we can only say that it appears to be actually as much in some cases as two years.

Symptomatology.—The symptoms of rabies vary in degree in different animals, but their sequence is identical as follows: 1. Mental disturbance (excitement, delusions); 2, hyperæsthesia, spasms; 3, hebetude, with mania and melancholia; 4, paralysis. Death, customarily followed by syncope or coma.¹ Clinically speaking, the disease assumes commonly (1) a mainly explosive

¹ The sudden arrest of the heart, which is so very common in this disease, seems in some cases to be due to extreme reflex excitation of the bulb.

form, (2) a mainly paralytic form. Until the last few years, the latter form was not recognised as a clinical entity in man, although numerous cases have been published; and it was no doubt owing to the general ignorance of the frequent existence of this kind of the disease that M. Peter and others made an absurd series of attacks on M. Pasteur, asserting that paralytic rabies was introduced by his method of treatment. I now throw photographs of different rabid animals on the screen, and you see the various stages of the malady.

I will now proceed to the discussion of M. Pasteur's mode of attempting to cure, or rather, to prevent the development of this malady. The cases he receives at Paris are divided into classes, according as to whether the evidence of the rabidity of the dog rests upon that of experimental inoculation, on veterinary certificate, or only a general report. Although the death-rate in the two latter classes amounts to nearly that in the first class, I shall only allude to the first class, and deal in the following remarks with those persons unquestionably the prospective victims of rabies. Hence, we are concerned with but one death-rate—namely, that of groups of persons bitten by rabid dogs, the rabidity being established by the fact of one or more individuals of the group dying of rabies or hydrophobia. Now the true death-rate among persons thus bitten amounts to 15 per cent. This important figure is the result of analyses by three different and independent observers.

LEBLANC, in 1873, carefully analysed the results of cases of rabies known to him personally, and found that the death-rate in unquestionable cases was almost 14 per cent.

DUJARDIN-BEAUMETZ, "Rapport Spécial au Conseil d'Hygiène et de Salubrité du Département de la Seine, 1887," found that 7 deaths occurred among 44 persons attacked by rabid animals, and not treated by M. Pasteur, the mortality thus being 15.90 per cent.

MYSELF.—Through the kindness of Colonel Moorsom, chief constable of Lancashire, I obtained the following facts. Six deaths occurred among 36 persons attacked; the mortality therefore was 16.6 per cent.

There can be no doubt, therefore, from examination of these figures, independently ascertained, that the true death-rate among persons bitten by indubitably rabid dogs, and not treated by M. Pasteur, is on the average about 15 per cent. With this definite fact before us, we learn that in exactly the same class (Class A) of patients M. Pasteur has obtained a death-rate of only 1.36 per cent. Though the success of his treatment is thus assured, it will not be uninteresting, I think, to dwell upon its scientific bearing.

Treatment.—Since the details of the pathology of rabies were dubious until M. Pasteur cleared up the whole subject, no treatment worth the name was ever proposed. In fact, the only attempt made was the lowering of the hyperexcitability of the congested reflex centres by empirical dosing with narcotics. While, of course, in no sense curative, I nevertheless believe that, after the appearance of the first symptom, this, for the sake of humanity, is the only justifiable course to be pursued.

M. Pasteur from the first recognised the true nature of the malady, and realised that the only possible mode of coping with the virus would be by attempting to arrest its development within the body. Arguing that it, the virus, was a microbe, he considered that we should be able to obtain the waste products of its metabolic processes, and that then such waste products (known to inhibit the growth of organisms) might be injected in sufficient quantity to check their progress. Every fresh investigation shows the clearness of M. Pasteur's insight into this question, since the validity of his theory receives daily confirmation from the record of the behaviour of the virus of other infective diseases.

The merit of proving the correctness of this theory by perfectly independent research is due to Dr. Wooldridge, who isolated a chemical substance which protected rabbits against anthrax. M. Pasteur believed it possible to obtain these waste products (unaltered by any mode of chemical extraction) by destroying the virulence of the microbe and not the protective substances. He therefore subjected portions of the central nervous system—that is, spinal cord by preference—to the combined action of heat and drying. By inoculation he discovered the rate of loss of virulence, according to the heat employed, etc. By injection into dogs of successive quantities of the material, he found that they became proof against the bites of rabid dogs, and at the same time, by their remaining in health, that they were in no wise affected injuriously by the treatment. Having in view that the average length of the incubation period of subcutaneous inoculation, as in the streets, was no less than six weeks, it seemed to M. Pasteur that, if the nature of

the case were recognised at once, the development of the virus in the body of a human patient might be prevented, if it were only possible to inject sufficiently rapidly a large quantity of the chemical protective substance. This he proceeded to do in the manner I have already indicated, and I shall only dwell upon the two great divisions into which this subject may be divided. These divisions are what are commonly known as the "simple treatment" and the "intensive treatment" respectively.

The "Simple Treatment."—By this, as you know well, M. Pasteur meant the injection of small quantities of the protective substance contained in cords, dried not less than from five to seven days, and this treatment he employed only in apparently mild cases.

The "Intensive Treatment."—This consists in injections, frequently repeated, of a considerable quantity of the substance from cords dried for shorter periods than in those of the simple treatment. This intensive treatment was, perhaps, the part of the method most attacked. It was, indeed, asserted by M. Pasteur's irresponsible opponents to be fatal in operation. The actual facts, however, are as follows: The only case completely investigated in which death was not averted by thorough employment of the method was that of the patient Goffi. This case I have fully published with Dr. Bristowe in the Clinical Society, November, 1888, and we there showed that the death was to be attributed to the cat-bite and not to the treatment. The intensive treatment is employed for severe cases, and, consequently, especially used in Russia, where the excessive prevalence of the disease and the severity of the wolf-bites make the statistics particularly valuable.

In the following table I contrast the results of treatment by the old or simple method and by the later or intensive plan.

Table showing the contrast between the Old or Simple Treatment and the Intensive Treatment.

	Simple Treatment, 1886.	Intensive Treatment, 1888.
Odessa.....	3.39 per cent.	0.64 per cent.
Warsaw	4.1 ..	0 per cent. (although figures include 16 months' work, and 30 individuals bitten in the face, and 4 by wolves).
Moscow	8.42 ..	1.60 per cent.

It is evident that the intensive treatment is very successful in coping with the worst cases, and that, instead of being itself a source of death, as asserted by those who gain notoriety and subsistence by vilifying and misrepresenting scientific progress, it is a powerful agent in saving life.

Summary.—To sum up, it is clear that M. Pasteur has succeeded in diminishing the percentage of risk in being bitten by a rabid dog from 15 to 1.36, and that this result is obtained by injecting a chemical protective substance.

Means of recognising the Disease in Suspicious Cases.—This is a point of considerable importance to medical officers of health and veterinary inspectors, since they are frequently called upon to authoritatively decide as to whether an animal is or has been suffering from rabies. For the correct detection of the disease in the living animal I need only refer you to textbooks and to the few remarks which I have already made on that point. But supposing one is shown a dead animal supposed to have been rabid, upon what is reliance to be placed for a correct diagnosis? Clearly (1) the *post-mortem* appearances; (2) the result of the method of experimental inoculation discovered by M. Pasteur. (1) *Post-mortem* appearances: It is very curious to see the amount of confusion and doubt concerning this branch of the subject which still seems to exist in some minds. The facts really are extremely simple and straightforward. In the first place, an animal may be killed soon after the disease has shown itself by the initial symptoms—that is, on the first or second day of the illness. At this stage the virus will not have had sufficient time to produce the characteristic changes in the tissues, and therefore nothing but a few patches of congestion may be found here and there in the central nervous system. The same negative result will also attend similar investigations in the cases of human beings who commonly die in the early stages from syncope. It is neglect of these very obvious facts that has led to the oft-repeated assertion that rabies has no characteristic morbid anatomy. In such cases, correct diagnosis rests alone upon experimental inoculation. But in a fully-developed case of rabies running its complete course the following changes, gross and minute, are to be observed. I shall simply sum them up categorically, and in illustration show

² This unusually high rate was found to be also due to imperfections in the mode of preparing the cords for the inoculation material.

you a few photographs upon the screen. For those which are microscopic I am indebted to my friend Mr. Boyce, they being made by him from my specimens.

Central Nervous System.—Early stage, congestion, followed later by exudation of leucocytes into the perivascular lymphatics, and interstitial neuroglia. Hæmorrhages also are frequent. (All as previously described by Allbutt, Gowers, and others.)

Peripheral Nervous System.—Congestion.

The changes in the central nervous system are frequently concentrated in localities, so that the whole must be carefully examined.

Respiratory System.—Congestion of laryngeal mucous membrane; congestion of tracheal mucous membrane; frequently patches of broncho-pneumonia.

Alimentary System.—Tongue furred and dry (especially in paralytic cases); pharynx congested, markedly at root of tongue. Stomach congested; hæmorrhages in the mucous membrane, specially in the cardiac end. Contents, coffee-ground material and foreign bodies.

Renal System.—Bladder distended in paralytic stage.

2. *The second method*, or that of inoculation, performed according to M. Pasteur's description, is the most valuable, since it forms an incontestable evidence of the presence or absence of the disease.

The extraordinary statements made by some writers, such as Spitzka, etc., to the effect that the symptoms of rabies in a rabbit are not peculiar or in any degree characteristic, and can be evoked by non-rabic material, etc., I only allude to because they are frequently paraded in our journals by writers who have obviously never consulted the original description of these so-called experiments, nor are possessed of any personal experience of the disease of which they treat. I here show you a photograph of a rabbit with rabies following a bite from a rabid dog of the streets. The paralysis is well marked. Suffice it to say that the experience of every careful observer on this subject entirely annihilates such loose and incorrect assertions.

Method.—The following is the method now generally adopted. About two inches of the upper cervical region of the spinal cord is removed as soon as possible after death, and placed in 10 per cent. solution of glycerine and water. This, while preventing decomposition if the fluid is frequently changed, does not destroy the specific rabic virus. When it is convenient to inoculate, a piece of the cord is washed in recently-boiled distilled water, then crushed in a glass mortar in *bouillon*, and injected subdurally through a very small trephine opening. The whole operation can be performed in a few minutes under local anæsthesia, produced by the application of 5 per cent. carbolic acid, which at the same time ensures the immediate union of the little wound. In some hundreds of cases I have never yet seen suppuration occur.

The animals most suitable for the inoculation test are rabbits, since in them the disease assumes a paralytic form, as a rule, from the commencement; and, as we know from cases in man, this form, besides being painless, terminates early by coma. As a rule, the first symptom appears on the fourteenth day, after which the animal commonly dies in four days. I think, however, that in those cases where the animals remain healthy they should be kept for inspection for three months after inoculation. I have carried out this method in thirty-six specimens sent me for diagnosis. Of the thirty-six, twenty-five, or nearly 70 per cent., proved to be genuine rabies.

In the present outrageous state of the law it is, however, impossible for any medical officer of health or veterinary inspector to perform this simple test, unless he is possessed of (1) a licence, (2) a certificate B, (3) and performs the experiment in a registered place, under 39 and 40 Vict. How long the profession and the public are willing to keep hygiene and science thus hampered remains to be seen.

Although M. Pasteur's mode of treating rabies has met with rather more virulent opposition than his mode of treating anthrax, it is fortunate that he has seen the adoption of the latter and the gradual perfection of the former.

At the present time this country is content to accept the charity of M. Pasteur, the very large majority of patients sent from England having made no return (in most cases they have been in poor circumstances) for the treatment received. Not only is our present position in regard to this disease humiliating for this reason, but the actual need for treatment ought not to exist in Great Britain, and it is a scandal that it does so exist. Clearly if by (1) proper measures we prevent its importation into

the country from infected lands abroad; (2) by muzzling we prevent dogs from communicating the disease by biting each other and other animals; (3) by destroying stray animals, notably dogs and cats, we prevent the frequent accumulation of sources of ready infection, the disease must die out, since, as I have already shown, the virus is apparently only transmitted from living animal to living animal. But the actual proof of this is now present in overwhelming degree; I need only give you two or three examples.

This photograph shows you the result of applying such measures as I have indicated in Berlin. You see the death-rate among men and cases among dogs were rapidly rising when the police regulations were enforced, and the disease was temporarily stamped out; a number of years later it reappears by introduction from without, and by confessed neglect of the regulations, but only to be again eradicated by the former means, and this time apparently permanently, for the same laws having been made practically universal throughout the German Empire, this disease is now becoming extinct, and would be so at once were it not constantly being imported over the Russian frontier.

In Scandinavia, formerly decimated, rabies is now unknown, by reason of their strict adherence to restrictive legislation. Paris furnishes us with the most recent evidence, a curve published by Dr. Grancher showing the immediate effect of the laws in cutting down the disease by merciful extermination of the already infected.

The records of the provinces, especially Nottingham (as shown by Dr. Whitelegge's figures) and Lancashire (report by Colonel Moorsom), exhibit very strikingly the fact that rabies prevails or disappears as the regulations are respectively neglected or properly enforced. But by far the most striking results are those obtained here in London, although they illustrate in a lamentably striking manner the folly of applying laws to isolated districts, which only purge that district, and do not provide against the reintroduction of the evil from the surrounding country.

Rabies in London destroyed 27 lives in 1885. The authorities then put on the muzzle, and ordered the slaughter of stray and unclaimed dogs; the result being, that by the end of 1886 it was practically extinct. The regulations were then allowed to lapse in accordance with the present imperfect legislation. Since that moment the disease has been steadily creeping into London again, especially from the south and south-west suburbs; in 1886-87 it raged among a herd of deer in Richmond Park, destroying some hundreds of them; and since that it has also been epidemic in Kingston, Surrey, on the borders of the park. Deaths among human beings are now gradually becoming more numerous. I have myself seen two cases, and had to do with a third, during the last three months. These cases came from different parts of London, so that the disease is as usual steadily becoming universal.

The Select Committee of the House of Lords, which reported on this subject in August, 1887, by the unfortunate influence of two of its members, Lord Onslow and the late Lord Mount Temple, stated that they considered it better to leave to the local authorities the initiation of necessary preventive measures, but they say, para. 26, that "the time may come when the public may call upon the Privy Council to act on its own authority throughout the kingdom." That that time has come when the public ought so to call, indeed, has always been present, is obvious from such a history as that of London, which I have just detailed to you, and we can only hope that the public interest may be sufficiently awakened to destroy for ever, as it easily might, so horrible a scourge.

The legislative machinery for this end has been drafted in the form of a workable Act by the Society for the Prevention of Hydrophobia and the Reform of the Dog Laws. Whether any Government will think it worth while to take the matter up and stamp out once and for all this, the most easily eradicated and yet the worst of diseases, remains to be seen. That it should exist among us is a standing disgrace to our civilisation.

DR. PHILLIPS, medical officer of the Leighlinbridge Dispensary, has resigned, and the Dispensary Committee have unanimously recommended that he should receive two-thirds of his salary and emoluments. Dr. Phillips has served in the district during the past thirty-eight years.

THE Duke of Cambridge has consented to preside at the festival dinner in aid of the funds of the Royal Westminster Ophthalmic Hospital at the Whitehall Rooms, Hôtel Métropole, on April 22nd.