material. Microscopically it contained hyaline cartilage and acinous adenoid tissue lying in young connective tissue. The glandular epithelium was cubical.

Fig. 8.—A persistent thyroid duct.

Shattock sums up his clear account of this tumour to this effect: "The specimen is doubtlessly related to the congenital sacral tumours, and has characters indicative of an early intra-uterine formation in some aberrant embryonic structure." Shattock also informed me that he could not find a thyroid body distinct from this tumour. This is a point of some importance, for I am of opinion that the tumour originated in the thyroid diverticulum, and must be regarded as an aberration analogous to those occurring in the post-anal gut. In support of this opinion I will adduce the following evidence.

The museum of St. Thomas's Hospital contains the upper part of the body of a mature foetus; the neck is occupied by a very large tumour which anatomically corresponds to the thyroid body, and resembles it in general configuration. Through the kind co-operation of Mr. Shattock I have been enabled to investigate this interesting specimen, and on making a sagittal section through it, the naked eye characters of the specimen recall in a striking manner those presented by congenital sacro-coccygeal tumours. The analogy however does not rest here, for on examining the tumour microscopically it is seen to be composed of loculi, varying in size from a millet seed to a small nut; the cavities contain mucoid material, but it is only here and there that evidence can be obtained of a parietal epithelium; this may be accounted for by the length of time the specimen has been preserved. The matrix of the tumour consists of richly cellular connective tissue in which tracts of well-developed hyaline cartilage occur; in many places this cartilage is undergoing direct transformation into bone. The cartilage in this specimen is typical of that found at the end of a foetal long bone, and differs from that so common in chondrifying sarcomata. In the latter case the cells are stellate, and do not present the sharp contour so characteristic of well-formed hyaline cartilage.

Thus in their naked eye character, in their minute structure and embryological associations, this particular form of congenital thyroid tumour corresponds so closely with the sacro-coccygeal tumours, that they ought to be grouped together.

It is an interesting fact that the six obsolete canals existing in the embryo of a mammal, namely, the infundibulum, neurenteric passage, post-anal gut, cranio-pharyngeal canal, thyreo-lingual duct, and the duct of the yolk-sac, should all have direct relation with the alimentary canal, and each be directly associated with dermoids often of considerable complexity, and with a peculiar form of tumour identical in structure with the thyroid body.

THE Marchioness of Dufferin and Ava has signified her willingness to become a patroness of the New Hospital for Women, in recognition of the value of the training it affords for medical women in India.

PRESENTATION.—On January 15th the male members of the Honiton class formed in connection with the St. John Ambulance Association presented their lecturer, T. W. Shortridge, M.D.Brux., with a handsome case of silver-mounted carvers as a token of appreciation and esteem.

INTUBATION OF THE LARYNX AND AIR-PASSAGES, WITH A DESCRIPTION OF A NEW INSTRUMENT AS AN AID TO CERTAIN OPERATIONS.¹

By THOMAS ANNANDALE, F.R.C.S.,

Regius Professor of Clinical Surgery, University of Edinburgh.

I DESIRE to consider the subject of intubation of the air-passages as concisely and as practically as possible, and, in order to do so, I propose to study the procedure under four heads. I cannot say that I have much that is new or original to tell, but as the subject is now attracting considerable attention, I hope that this short note of the practical and clinical facts connected with it may not prove uninteresting.

1. *Intubation as an aid in certain operations.*—In operations, involving the mouth or naso-pharynx, in which bleeding may take place into the air-passages, and interfere with the respiration, or in operations in which the respiration is not satisfactory during the administration of an anæsthetic, preliminary tracheotomy or laryngotomy, with or without plugging of the throat, is a useful and well-recognised procedure. Experience has, however, taught me that intubation of the trachea through the mouth can, in the majority of instances, be substituted for tracheotomy, and is a simpler and safer means of aid. I give a note of the following case in illustration. About nine months ago, a man was brought to the infirmary from the country to me suffering from great difficulty of respiration, caused by the pressure of a very large and malignant tumour of the thyroid gland. The larynx and trachea were completely buried in the tumour, and after some search I was only able to feel the top of the upper part of the thyroid cartilage, and it was upon a level with the lobe of the left ear. It was necessary at once to do something to relieve his breathing; and I, accordingly, administered chloroform, with the intention of cutting down upon the thyroid cartilage, opening into the larynx, and introducing a tube. Before operating, I had provided myself with a gum-elastic catheter (No. 10), having an opening at the extremity and an efficient mouth gag. It was well I did so, for during the operation his breathing ceased, and it was only by introducing the catheter through the glottis that his respiration was restored. Through this catheter the anæsthetic was further given, and the operation completed with success. I have since employed a similar plan in operations attended with a risk of bleeding into the air-passages. The tube used in these cases was, as has been stated, a gum-elastic catheter, and it is as well to have in it a stilette of strong wire—the point of which should not project through the opening at the end—as a certain amount of rigidity makes the introduction of the tube easier.

Dr. Macewen, of Glasgow, appears to have been the first surgeon to use intubation on the principle I have just detailed, and I would refer you to an important and practical paper by him upon this subject in the BRITISH MEDICAL JOURNAL for July 24th and July 31st, 1880. Dr. Macewen's first operation was performed on July 5th and 6th, 1878.

2. *Intubation as a means of restoring or carrying on respiration in cases of sudden obstruction in the larynx or trachea.*—There will, I think, always be cases of this kind, in which rapid laryngotomy or tracheotomy will most quickly, and consequently most successfully, restore the respiration; but many of these sudden emergencies can also be treated rapidly and successfully by the introduction of a tube through the mouth into the trachea, and this tube may, if necessary, be retained there for some time. A few weeks ago a young man was admitted into my wards, on account of a large swelling of the neck, apparently of glandular origin. The larynx and trachea were slightly displaced to the left side, but the respiration was not much interfered with. Two days after admission, a rapid increase, owing to an attack of acute inflammation, took place in the swelling, and the breathing became most difficult. Instead of performing laryngotomy or tracheotomy, operations which would have been difficult in this case, my house-surgeon, Dr. Simpson, introduced a tube into the trachea, through the mouth, and this at once relieved the patient. The tube was retained for twenty-four hours until I saw him, when chloroform was administered through the tube, and a deep incision made, which gave exit to a collection of pus. This having

¹ The greater part of this paper was read before a meeting of the Northumberland and Durham Medical Society.

taken off the pressure from the air-passages, the tube was removed and the man progressed satisfactorily. Still more recently, in a case of scald of the mouth and throat from inhaling hot steam, a similar introduction was most successfully practised. The patient was a child, aged 5, who, four hours before admission, had attempted to drink out of the spout of a kettle filled with boiling water. When admitted, the difficulty of respiration was great, and there was cough and great restlessness. The inhalation of warm vapour charged with benzoic acid was at once employed, but the symptoms became gradually worse, and at 2 A.M. next morning they were most urgent. Dr. Simpson accordingly inserted a No. 9 elastic catheter by the mouth through the glottis. The introduction of the catheter caused some irritation and coughing, and a considerable amount of muco-serous fluid was discharged through it. After this the child fell into a calm sleep, being only occasionally disturbed by the discharge of more of a similar fluid. About two hours after the introduction of the tube, the patient had an attack of vomiting, and the tube was withdrawn, as the respiration was now much relieved. The symptoms continued favourable and breathing satisfactory for the next few days, and two weeks after the scald all anxiety in connection with the case was at an end, and a perfect recovery prognosed. In the paper already referred to, Dr. Macewen records several cases of oedema glottidis successfully treated in this way; but Dr. Huch, of Freiburg, had previously treated one case of acute stenosis of the larynx by introducing one of the tubes suggested by Schrötter.

The intubation under these two first heads is, in my opinion, best carried out by the long catheter, or similar tube, one end of which protrudes through the mouth. It has been suggested that such tubes should be passed through the nose; but it is much more difficult to pass a tube in this way through the glottis into the larynx and trachea, and, therefore, I prefer to pass it through the mouth.

One disadvantage of the gum-elastic catheter or soft tube is its liability to be compressed, and its canal consequently obstructed by a twist, or by the pressure of the teeth; and it therefore occurred to me that a special tube might be used, in order to avoid this risk of interference with the respiration, more particularly when it is being used in operations, or temporarily, as in cases of sudden laryngeal or other obstruction of the air-passages. Accordingly, when recently in New York, I asked Messrs. Tiemann and Co., the well-known instrument-makers, to make for me an instrument which would carry out this principle. Messrs. Tiemann, with their usual ingenuity, have now constructed for me an elastic tube—not too pliable—shaped after Schrötter's one, and having an arrangement to prevent the compression of its canal from contact with the teeth, or otherwise. To the end of the tube protruding from the mouth there is also attached a piece of india-rubber tubing, through which an anæsthetic may be administered, at a convenient distance, in the case of an operation.

3. *Intubation in cases of acute inflammatory affections of the Larynx, more particularly in croup and diphtheria.*—It is in connection with these affections that intubation has attracted and is attracting much attention; and, already, considerable experience of it, particularly in America, has been recorded.

In 1858, Bouchut read, before the Academy of Medicine, in Paris, a paper entitled "On a New Method of Treating Croup by Tubage of the Larynx." Bouchut's few attempts were unsuccessful, and the operation was condemned by a committee of his colleagues.

In 1885, Dr. O'Dwyer, of New York, published a paper in the *New York Medical Journal* for August 8th, 1885, and it is to him that we owe, as a result of observations and experiments carried on for five years prior to this date, the successful establishment of the operation.

The principle of Dr. O'Dwyer's operation is to introduce into the larynx, through the glottis, a metal tube of a shape to fit comfortably in the canal, and of a size according to the age of the patient and capacity of the cavity, and to retain the tube as long as may be required, provided it allows the respiration to be properly carried on. In fact, it is proposed by this proceeding to relieve, in many cases, the laryngeal obstruction to respiration, instead of performing tracheotomy.

When attending the recent Congress of Physicians and Surgeons in the United States, I had an opportunity of listening to several papers on the subject, and also of discussing the question with gentlemen who had had large experience of the operation; and although no one considered that intubation would take the place of tracheotomy in every case, there was a very general opinion

that in patients under 4 years of age intubation gave better results than tracheotomy. Dr. Ingals's (*New York Medical Journal* for July 2nd and 9th, 1887) reports that he had collected 514 cases, with 134 recoveries, or 26.7 per cent., whereas the recovery after tracheotomy in patients under 4 is only from 15 to 20 per cent.

For a well-illustrated and excellent paper upon the performance of this operation, I would refer to one by Dr. Waxham, in the *BRITISH MEDICAL JOURNAL* for September 20th, 1888. Dr. Waxham has collected notes of 1,072 operations performed in America, and out of these 287 recovered, about 27 per cent. He also gives an interesting table proving the advantages of intubation over tracheotomy in young children. Taking only one example from this table, it is found that, in children between 2 and 2½ years, the recoveries after tracheotomy were 12 per cent.; after intubation 19.46 per cent. Dr. G. U. Gay, surgeon to the City Hospital, Boston, U.S., has published, in the *Boston Medical and Surgical Journal* for October 11th, 1888, an interesting paper, in which he discusses fairly and independently the question of the "Comparative Merits of Tracheotomy and Intubation in the Treatment of Croup;" and I think that the following quotation from the paper is important in connection with the consideration of the subject:—"I have thus in a very imperfect way endeavoured to show that, while intubation is a most valuable operation, it is not as free from objections and complications, and is not so far superior to the old and time-honoured operation of tracheotomy as some of its advocates would lead us to suppose. The method has great advantages. In favourable cases it is easily and quickly performed; there is no cutting, and hence no hæmorrhage; no anæsthetic is required; as it is not looked upon by the laity in the same light as an ordinary surgical operation, consent for its performance is more readily granted; it can thus be done earlier in the disease; many practitioners will undertake the new method who would shrink from the old one; the tube takes care of itself; it is often coughed out when no longer required; it is especially adapted to young children. But, on the other hand, the operation may be difficult and even dangerous to perform; it may not relieve the dyspnoea; the tube may be repeatedly coughed out or it may be swallowed; it may 'gum,' thus requiring frequent removals; it may become suddenly occluded, and, unless quickly ejected or removed, death will ensue; and there may be great difficulty in feeding the patient. Both operations are often attended with difficulties, and occasionally with danger; both are liable to complications; the mortality attending each is about the same; the fatal results in a majority of instances are due to the extension of the disease to the lungs, and this occurs as frequently after one operation as after the other; in neither case is the treatment the cause of the complication, nor does it prevent, or cure it. In conclusion, I would say that, in a majority of cases of membranous laryngitis, intubation may be done with a fair prospect; that it will effectually relieve the dyspnoea for the time being. That it is to be preferred in young children and in all cases living at a distance from skilled aid, where the tube must be allowed to take care of itself. That it may be resorted to preliminary to tracheotomy. That it may be done for euthanasia, providing the operator is reasonably expert and can do it quickly without producing collapse. Tracheotomy is indicated in those cases in which intubation cannot be done, or in which it fails to give relief to the dyspnoea. In severe cases, situated at such distances, or under circumstances in which only ordinary and not skilled assistance can be obtained in such an emergency, tracheotomy is the safer method. It is also to be preferred in those cases of intubation which cannot be fairly nourished either in the natural way or by enemata, etc. It may be resorted to when the O'Dwyer tube is frequently ejected, or when it requires frequent removal to prevent obstruction. I would repeat that as regards patients residing at a distance, if competent aid is to be at hand during the convalescence, intubation is advisable as it is if no care whatever is to be given to the tube. But if ordinary care and no other in case of accident can be commanded, and I mean by that, a fairly good nurse, or any clever person, then tracheotomy is the better procedure."

It may be well now to consider the advantages claimed for intubation in this class of case, and then to consider any of its disadvantages as compared with tracheotomy.

ADVANTAGES.

- (1) No anæsthetic is required, and it can be done at once without shock or the risks attending tracheotomy, especially in young patients.
- (2) No external wound is made, and, as the breathed air passes

through the mouth and is so warmed, there is less risk of pneumonia or bronchitis.

- (3) No after-treatment is required when the tube is removed.
- (4) Consent of friends is more readily obtained than in case of tracheotomy.
- (5) Tracheotomy can at any time be performed if intubation fails.
- (6) In many cases the tube can be removed early—third, fourth, or fifth day.

DISADVANTAGES.

- (1) When introducing the tube, the membrane may be pushed down and obstruct respiration.
- (2) The tube may be obstructed by membrane passing into it.
- (3) Food may pass down the tube and cause pneumonia or other complications.
- (4) The tube may slip down into the trachea.
- (5) The tube may be coughed up.
- (6) If the tube becomes displaced or obstructed, it is not easily reintroduced by a nurse.

The first of these accidents would in all probability give rise to immediate symptoms, so that the membrane might be removed, either by means of laryngeal forceps or by the performance of tracheotomy. The second is a rare occurrence, according to the experience of operators, and can only be remedied by the removal of the tube and replacement after its canal had been cleared, or by the performance of tracheotomy, if the symptoms were urgent. To prevent the third, it is advised that food, especially fluids, be at first given in small quantities, and the nourishment assisted by nutrient rectal injections, or by means of a tube passed into the oesophagus. There is a considerable difference in different patients in connection with the power of swallowing after this operation, and many swallow well after the first day or two. I believe much depends on the exact position of the tube, and the amount of interference with the action of the epiglottis. Dr. T. E. Waxham has suggested a modification of O'Dwyer's tubes, with the addition of a soft rubber false epiglottis; and, still more recently, Dr. J. Mount Bleyer has employed a tube, more cup-shaped at its upper end, and having also a false rubber epiglottis. Dr. Bleyer is of opinion that these latter tubes have afforded him better results than he obtained by O'Dwyer's original ones. The fourth and fifth very rarely occur. The sixth point should not, I think, be a bar to the operation, for in such cases it is usual to have some skilled assistance at hand which should be able to replace the tube if required. If such skilled aid cannot be available, tracheotomy would be preferable.

I conclude this section of my paper with a brief note of a case (my only case as yet), recently treated by intubation. A little girl, aged 3, suffering from great difficulty in respiration, the result of an attack of croup, was brought to me at the Royal Infirmary by Dr. Sloan, about 12 o'clock on the forenoon of November 9th. She was attacked with the croup upon the 6th, three days before, and although carefully treated, her symptoms grew worse, and the laryngeal obstruction became so marked that immediate relief was demanded. Instead of performing tracheotomy, I at once introduced one of O'Dwyer's tubes. When the tube entered the glottis, there was a violent spasmodic cough, and a piece of membrane was ejected through it. Immediately after this, when the tube had been placed in position, the breathing was relieved, and ten minutes after, when I visited the child in the ward, she was respiring naturally, and was asking, in a whispering voice, for a "piece." For the first few hours there was a difficulty in swallowing, much of what was taken being ejected; but after this she was able to swallow with ease, and took milk in abundance. On the following day, the 10th, the respiration was more rapid, and the face flushed, but the tube was causing no irritation, and there was no laryngeal obstruction. The symptoms, however, pointed to commencing pneumonia. On the 11th, the lung condition was worse, but the tube was retained, and there were no signs of laryngeal obstruction. She had no difficulty in swallowing. On the 12th she became much exhausted, and died the same forenoon, the tube having been retained until the last. The *post-mortem* examination, which was confined to the throat, showed that the laryngeal condition had been quite relieved, and that the tube had caused no ulceration of the portion of the canal upon which it rested. Death was the result of pneumonia, which had probably been caused by the exposure of the patient when brought to the hospital. The points which convinced me of the value of intubation in connection with the experience of this case were:

- (1) The ease with which the tube was introduced.

(2) The complete relief to the respiration when it had been properly adjusted.

(3) The fact that the tube caused no irritation, although retained for some days, and carried on efficiently the laryngeal respiration.

(4) The fact that after a few hours the patient was able to swallow fluids well.

(5) The satisfactory condition of the larynx as proved by the *post-mortem* examination.

4. *Intubation for Stenosis of the Larynx, the result of Chronic Inflammatory conditions, or of accidental or surgical wounds.*—The introduction of O'Dwyer's tubes in this class of case has been followed by better results than have been obtained by any other plan of treatment, and these tubes have been worn in the adult and also in the child for a week, without requiring to be changed, and in a few cases for several months. In one case, related by O'Dwyer, the tube was worn for ten months. Intubation in these cases not only relieves any temporary obstruction to respiration, but the permanent retention of the instrument, when it can be borne, is likely to have a beneficial effect in restoring the proper calibre of the canal.

A. A.—Tracheal end and opening.
B.—Free end to which india-rubber tube is connected.
C.—Gag of hard vulcanite, which is movable along the tube.

FURTHER RESEARCHES ON THE TREATMENT OF TINEA TONSURANS, ILLUSTRATED BY MICRO-PHOTOGRAPHS.¹

By A. J. HARRISON, M.B.Lond.,
Physician, Bristol General Hospital.

At our Association meeting held at Cardiff in 1885, I had the honour of reading a paper before the Section of Medicine on a "New Method of Treating Tinea Tonsurans," and I then alluded to the unsatisfactory modes adopted in the treatment of this very troublesome complaint, and I referred to the multiplicity of remedies so-called; and I formulated a statement that the efficacy of the treatment of any disease may be considered to stand in an inverse ratio to the number of remedies propounded. No one, except very rarely, ever thinks of treating itch by any other remedy than sulphur, or some compounds of sulphur, and why? Because sulphur almost invariably cures the complaint by destroying the acarus, or mite, on which the ailment depends.

I also went on to state that the great difficulty of getting at ringworm of the scalp was a physical one, the parasitic fungus revelling and running rampant in the secure nidus of the hair and hair-follicle, a soil so congenial that it seems to be in some cases ever fertile and ever fertilising. To overcome this difficulty, and so get at this secure fungus retreat, I indicated that no remedies could be considered highly efficacious which did not act chemically upon the hair-matter itself, or perhaps, to define more accurately, which could act so, and at the same time carry with them a parasiticidic agent; and I showed that I had found the caustic alkalies the most useful hair-solvents, but as they are not conidia destroyers, it was necessary to combine other agents with them, and I came to the conclusion that the application of two solutions, the first, composed of liquor potassæ, spirits of wine, and iodide of potassium; and the second, mercuric chloride dissolved in spirits of wine, and water, constituted a very efficacious mode of treatment. The hair-matter is softened by the

¹ Read in the Section of Medicine at the Annual Meeting of the British Medical Association, held in Glasgow, August, 1888.