through the dental arches. Pressure over both condyles produced pain and some slight crepitus, more marked on the right side. The appearances were in favour of dislocation, but there was no

Fig. 4.—Special Bandage.

hollow behind the condyles. I obtained the aid of Mr. Davies-Colley, and under chloroform a thorough examination was made, revealing fracture through the neck of both condyles. The deformity was reduced, and a chin bandage was applied in the hope of holding the jaw in place till a splint could be made; but directly the patient began to come to the deformity began to recur, and very soon it was necessary to cut the bandage to relieve the pain. In consultation with Mr. Davies-Colley I made a full-size Gunning splint, with deep vulcanite rims, taking its fixed point from the upper jaw, and thus holding the inferior maxilla forward. The models were taken under chloroform, and the splint inserted under ether. A perfect position of the fragments could be obtained under the anæsthetic, but partial relapse took place when the patient came to, in spite of the deep rims of the splint and firm pressure applied by a special chin bandage. The latter was made of strips of webbing united by buckles (Fig. 4). The obliquity of fracture and the muscular action were such as to defy our efforts to bring the fragments of the jaw into their correct position in the splint by any pressure that the patient could readily bear. Increased pressure at once produced pain, swelling of the lower lip, and ulceration of the mucous membrane in contact with the splint. By degrees the lower jaw settled down into an improved position, and the patient was prepared for some slight separation of the front teeth at first. In five weeks the splint was removed under ether, and it was found that good union had taken place on both sides. The cusps of the molar teeth impinged on one another, and when these were smoothed off the separation of the upper and lower front teeth was an eighth of an inch, and this has gradually disappeared. The other injuries from which the patient was suffering were injury to ribs (fracture) and fracture of neck of scapula.

CASE OF ACTINOMYCOSIS OF THE LIVER.

By FREDERIC S. EVE, F.R.C.S.,

Assistant-Surgeon to the London Hospital, etc.; Curator of the Pathological Museum, Royal College of Surgeons.

THE following case was alluded to in an article on actinomycosis I contributed to the *Practitioner* of May, 1888, page 327. At that time the diagnosis was not confirmed, actinomycetes not having been found in the discharge from sinuses; but at the request of the editor a reprint of my paper appeared in the *Veterinarian* of September, 1888, in which I inserted a note stating that the diagnosis had been confirmed by naked-eye and microscopic examination of the liver.

W.S., aged 60, a sailmaker, came as an out-patient to the London Hospital, complaining of pain and swelling over the hepatic region. He was spare almost to emaciation. I found distinct enlargement of the liver, which extended some distance below the costal margins. In the right hypochondrium a rounded nodule could be felt, which moved with the diaphragm, and appeared at that time to be unattached to the abdominal wall. The swelling was thought to be either a gumma or an abscess, projecting from the surface

of the liver. Seven to fourteen days later, on May 11th, 1887, he was admitted under the care of Mr. McCarthy. The patient stated that three months before he began to suffer severe pain in the right side, just under the ribs. Fourteen days subsequently he noticed a swelling, at first painless, but which became painful as it grew larger. He admitted having a sore on the penis some years ago, while abroad as a sailor. There was a tender, softish swelling, three inches in diameter in the right hypochondrium. The skin over it was discoloured and œdematous.

On May 12th an incision was made into the swelling, but only soft granulation tissue came away. A drainage-tube was inserted, and the wound dressed with iodoform.

May 17th. Very slight discharge. Skin around opening discoloured, and dense. There was slight pyrexia, the temperature rising 1° to $1\frac{1}{2}^{\circ}$ at night.

Iodide of potassium was prescribed, and he was discharged from hospital on May 27th. I readmitted him a week later, on finding that the skin around the incision was undermined, and that sinuses were forming. I laid open two sinuses, one on each side of the opening, and scraped the infiltrated tissues.

June 27th. The infiltration of the integuments had increased. Small subcutaneous abscesses continued to form, which intercommunicated by sinuses. Pyrexia very slight, the temperature occasionally rising to 100° .

After leaving the hospital he continued to attend as an out-patient, and I ordered liq. hyd., perchlor. and potas., iodid., but without marked benefit. Fresh sinuses formed at intervals, until the right hypochondrium and epigastrium were marked by some six or more openings, from which a thin watery fluid, but no pus, exuded. He suffered no pain. The enlargement of the liver appeared to decrease rather than the reverse. Seeing that anti-syphilitic remedies had no distinct effect, and taking into consideration the general aspects of the case, I began to suspect that the disease was actinomycosis, and on several occasions examined the discharge and scrapings from the sinuses for the ray-fungus, but without success. I continued liq. hyd. perchlor. in $\mathfrak{3j}$ doses, in the hope that it might retard the progress of the disease. He gradually got weaker and more emaciated, and finally in March, 1888, he became confined to his bed. On visiting him I found ascites, but no other change in the general condition. He died from exhaustion at the end of April.

Fig. 1.—Actinomycosis from human liver. (It is drawn rather too large).

The friends would not permit a necropsy, but I succeeded in persuading them to allow me to make an incision into the abdomen, through which I removed a piece of liver. This was studded with the characteristic nodules of actinomycosis. They had not undergone softening, and did not present the usual honeycombed

structure. Scrapings of the nodules showed, under the microscope, the well-known clubs, two or three clubs being often attached to a single filament. Sections made after embedding in celloidin were stained with Plaut's method, and also with fuchsin and methyl-blue. They were studded with numerous actinomycetes, the circles being large, and surrounded by the character-

Fig. 2.—Actinomycosis from tongue of ox for comparison.

istic rays or club-shaped bodies (see Fig. 1) I could not observe prolongation of filaments into the centre of the mass, as I have figured them in Fig. 2. I did not detect any essential difference between these specimens and those I had examined from oxen (see Fig. 2). The circles, however, appeared wider, the granular centres more extensive, and the separate centres of growth remained distinct, instead of forming irregular and mulberry-shaped masses.

REMARKS.—This case appears quite typical of the usual course of actinomycosis, namely, in its slow progress—the duration of the disease being a year and three months—in the absence of acute symptoms, severe pain or fever—in the appearance of a fluctuating point from which only granulation-tissue was evacuated by incision, and the subsequent formation of numerous sinuses, with undermining of the skin. This combination of symptoms, coupled with the exclusion of syphilis, abscess, and hydatid, convinced me that I had to deal with a case of actinomycosis, although I could not find the fungus during the life of the patient. The pain complained of at the early period of the disease was probably due to local peritonitis, and ceased when adhesions had formed between the liver and abdominal parietes. The organism probably gained access to the liver through the intestine.¹

A CASE OF OSSIFICATION OF THE CHOROID.²

By E. F. DRAKE BROCKMAN, F.R.C.S.,
Surgeon-Major, Madras Army.

THE case which I wish to submit for the consideration of the members of this Branch will, I think, prove of interest in two chief ways, first, in reference to the rarity of the affection; and, secondly, in reference to the extraordinary changes to which, under some circumstances, the vascular structures of the eye are liable. I have brought here this afternoon a section of an eyeball in which an ossific change has occurred; also some dry sections of the bony formation mounted for microscopic investigation, and two sketches in sepia and chalk of the bony structure.

1. As regards the rarity of the affection. Among fifteen authors who have treated on diseases of the eye, I find eight have referred to this subject in a casual manner, four have given a fuller account of the pathological condition, and three have omitted mention of it altogether. The most elaborate description appears

in De Wecker's classical work, entitled, *Traité Complet d'Ophthalmologie*. I have also searched in the records of the Ophthalmological Society, in the *Transactions of the Pathological Society*, in the reports of the Ophthalmic Hospital at Moorfields, and in the *Archives d'Ophthalmologie*, and I find that the reported cases of this nature are, comparatively speaking, few in number. Among the large number of patients who have passed under my observation at the Ophthalmic Hospital, Madras, and among the numerous eyeballs I have dissected after enucleation, this is the first specimen of ossific change in the eyeball with which I have met. At one of the meetings of the Ophthalmological Society in London in 1884, a similar case was exhibited by Mr. Adams Frost, which I was fortunate enough to see. To us, therefore, who are here present, and who, perhaps, have not many opportunities of studying eye-diseases, this specimen will, I trust, be of interest on account of its rarity at least.

2. The second point of interest has reference to the peculiar metamorphosis which the soft vascular coat of the eyeball is capable of undergoing under diseased conditions. Such decided ossific changes are, as far as I am aware, unknown in other parts of the vascular system of the body, and the primary cause for its appearance in the choroid has been usually regarded as inflammatory in its nature. Fibrinous lymph appears to have been thrown out at first as a result of the inflammatory action in the vascular tissue, which afterwards undergoes developmental changes into fibrous tissue, and subsequently into bone structure. This process must not be confused with petrification or calcification of a pre-existing structure. The latter, being an infiltration of a tissue with calcic phosphates and calcic carbonates in a solid form, is distinctly degenerate in its character, and must be regarded as a retrogressive change in a tissue of a more elaborate nature to one of less histological importance. Although these processes are of so decidedly opposite a nature, either can prove equally productive of mischief when present in the tissues of the eyeball.

Ossification of choroid, dry section $\times 50$.

It is a remarkable fact, which has received verification by the observations of many oculist-pathologists, that neither in the iris nor in the ciliary body has true ossification been known to occur, although these structures are regarded as the direct anatomical continuation of the choroid anteriorly. These tissues are known to be liable to calcification and pseudo-osseous formations, but not to the development of true bony structure. The layer of the choroid in which the initial changes are said to take place is that which is known as the *lamina vitrea*, which lies in apposition with the chorio-capillaris layer and separates this layer from the retina.

"The chorio-capillaris, so rich in vessels, sets free the essential materials for the formation of the osseous masses, which terminate in their development at the point where the chorio-capillaris terminates." De Wecker, from whose work I have translated the preceding passage, goes on to say: "The diseases, which chiefly predispose to an ossification of the choroid are those forms of chorioiditis which follow on injuries." Others are of opinion that an irido-choroiditis of idiopathic origin is equally powerful in bringing about osseous formation in the choroid; and in the case which I am about to relate a preceding syphilitic irido-choroiditis

¹ The above account was written six months ago.

² Read at the meeting of the South Indian and Madras Branch of the British Medical Association on October 12th, 1888.