FOUR CASES OF FRACTURED INFERIOR MAXILLA: TREATMENT.

By F. NEWLAND-PEDLEY, F.R.C.S., L.D.S.,
Dental Surgeon, Guy's Hospital.

A CONSIDERABLE number of patients suffering from this injury seek treatment annually at Guy's Hospital, and three of the cases I purpose recording were incomplete a month ago. Each of the four illustrates a whole class of fractures, and all of them required mechanical contrivances applied within the mouth.

CASE I.—A middle-aged man received a kick from a horse, and presented himself with a lower jaw that was literally collapsed by three fractures, one vertically on each side of the mouth in the region of the canine tooth; and a third running transversely above the chin separating the lower front teeth with their sockets from the base of the bone. The sides of the jaw fell inwards, and it required moderate digital pressure to bring the lateral segments into position, though the central piece was quite loose. This was a typical case for the Hammond wire splint, for the parts could be brought into correct position by means of moderate

Fig. 1.—Model showing displacement of fragments before treatment.

Fig. 2.—The same, with deformity corrected by sawing through the lines of fracture, and rearticulating, with model of upper jaw. Hammond wire splint applied.

pressure, and there were firmly implanted teeth in each fragment. Plaster-of-paris models were taken of both jaws, and a true model of the lower jaw with deformity corrected was obtained by sawing through the plaster along the lines of fracture, and rearticulating the fragments with the cast of the upper teeth (Figs. 1 and 2). A loop of steel wire was made to pass round all the teeth of the lower jaw along the gum margin, and was affixed to the fragments in corrected position by fine binding wires in the usual way. The splint was removed in seven weeks, the parts being restored to normal position. Lately I have used piano-wire and fine knitting needles for making the frame; the steel possesses greater strength, and

diminished bulk as compared with iron. Gold solder should be used.

Fig. 3.—Gunning splint.

CASE II was a boy aged 8, with a fracture by the side of the canine tooth on the right side. He had been run over, and his face and head were very much swollen and injured. The posterior fragment was much raised, and required considerable pressure to reduce the displacement. The only teeth erupted were the first permanent molars and the central and lateral incisors, and these were very short. The temporary molars were carious and the condition of the mouth generally was not favourable to Hammond's splint. Models were taken, and the deformity was corrected on the plaster cast. A Gunning's splint (Fig. 3) was applied, consisting of a vulcanite plate covering the palate, teeth, and gums of the upper jaw and united by vulcanite pillars to a similar plate covering the teeth and gums of the lower jaw. The separation of the jaws allows spaces to be left between the columns of vulcanite for speech and the passage of food and saliva. For more than a week this splint did little good beyond limiting the movement of the fragments, for the displacement recurred as often as it was corrected, and the boy's face was so swathed in surgical dressings it was out of the question to apply pressure by a chin bandage. Gradually the fragments settled down into their position in the splint and, with the rapid union common in childhood, the case was complete in five weeks.

CASE III was a young married woman admitted into the wards of Guy's Hospital with necrosis of the lower jaw. Nearly the whole depth of the horizontal ramus from the median line to the angle was removed as a sequestrum. A few days afterwards the arch of the jaw yielded and collapsed, the sound side deviating towards the affected side and constituting a remarkable deformity due to "spontaneous" fracture. At the request of a surgical colleague, I constructed a form of Gunning's splint. A vulcanite appliance was made to cover the whole available surface of the upper jaw and gums, and to the lower surface of this there was attached a vulcanite cap that fitted over the teeth and gums of the lower jaw on the sound side only. The splint was inserted under chloroform, and the space left free on the affected side gave ample room for the introduction of food. This appliance was worn continuously for five months, at the end of which time the power to depress the lower jaw was obviously restored, with absence of pain. On the removal of the splint the teeth of the upper and lower jaw were in their correct relation to one another, and the maxillary arch was again firm.

CASE IV was a patient in private practice. On her wedding trip to London she was knocked down and run over by a trap, in which two men were seated. The wheel passed over her, and she was so injured it was thought she was dead. She consulted lady doctors, especially about deformity of her face, but was assured there was no maxillary fracture. A fortnight elapsed and she returned home to the West of England, where she saw her family doctor, who advised her to consult a local dental surgeon. This she did, and at his instance she returned to London, three weeks after the accident, to be placed under my care. The face showed separation of the upper and lower front teeth to the extent of about three-fourths of an inch, and the lower jaw receded greatly. The mouth could be opened, and there was no deviation of the median line. Examination of the oral cavity revealed no fracture

through the dental arches. Pressure over both condyles produced pain and some slight crepitus, more marked on the right side. The appearances were in favour of dislocation, but there was no

Fig. 4.—Special Bandage.

hollow behind the condyles. I obtained the aid of Mr. Davies-Colley, and under chloroform a thorough examination was made, revealing fracture through the neck of both condyles. The deformity was reduced, and a chin bandage was applied in the hope of holding the jaw in place till a splint could be made; but directly the patient began to come to the deformity began to recur, and very soon it was necessary to cut the bandage to relieve the pain. In consultation with Mr. Davies-Colley I made a full-size Gunning splint, with deep vulcanite rims, taking its fixed point from the upper jaw, and thus holding the inferior maxilla forward. The models were taken under chloroform, and the splint inserted under ether. A perfect position of the fragments could be obtained under the anæsthetic, but partial relapse took place when the patient came to, in spite of the deep rims of the splint and firm pressure applied by a special chin bandage. The latter was made of strips of webbing united by buckles (Fig. 4). The obliquity of fracture and the muscular action were such as to defy our efforts to bring the fragments of the jaw into their correct position in the splint by any pressure that the patient could readily bear. Increased pressure at once produced pain, swelling of the lower lip, and ulceration of the mucous membrane in contact with the splint. By degrees the lower jaw settled down into an improved position, and the patient was prepared for some slight separation of the front teeth at first. In five weeks the splint was removed under ether, and it was found that good union had taken place on both sides. The cusps of the molar teeth impinged on one another, and when these were smoothed off the separation of the upper and lower front teeth was an eighth of an inch, and this has gradually disappeared. The other injuries from which the patient was suffering were injury to ribs (fracture) and fracture of neck of scapula.

CASE OF ACTINOMYCOSIS OF THE LIVER.

By FREDERIC S. EVE, F.R.C.S.,

Assistant-Surgeon to the London Hospital, etc.; Curator of the Pathological Museum, Royal College of Surgeons.

THE following case was alluded to in an article on actinomycosis I contributed to the *Practitioner* of May, 1888, page 327. At that time the diagnosis was not confirmed, actinomycetes not having been found in the discharge from sinuses; but at the request of the editor a reprint of my paper appeared in the *Veterinarian* of September, 1888, in which I inserted a note stating that the diagnosis had been confirmed by naked-eye and microscopic examination of the liver.

W.S., aged 60, a sailmaker, came as an out-patient to the London Hospital, complaining of pain and swelling over the hepatic region. He was spare almost to emaciation. I found distinct enlargement of the liver, which extended some distance below the costal margins. In the right hypochondrium a rounded nodule could be felt, which moved with the diaphragm, and appeared at that time to be unattached to the abdominal wall. The swelling was thought to be either a gumma or an abscess, projecting from the surface

of the liver. Seven to fourteen days later, on May 11th, 1887, he was admitted under the care of Mr. McCarthy. The patient stated that three months before he began to suffer severe pain in the right side, just under the ribs. Fourteen days subsequently he noticed a swelling, at first painless, but which became painful as it grew larger. He admitted having a sore on the penis some years ago, while abroad as a sailor. There was a tender, softish swelling, three inches in diameter in the right hypochondrium. The skin over it was discoloured and œdematous.

On May 12th an incision was made into the swelling, but only soft granulation tissue came away. A drainage-tube was inserted, and the wound dressed with iodoform.

May 17th. Very slight discharge. Skin around opening discoloured, and dense. There was slight pyrexia, the temperature rising 1° to $1\frac{1}{2}^{\circ}$ at night.

Iodide of potassium was prescribed, and he was discharged from hospital on May 27th. I readmitted him a week later, on finding that the skin around the incision was undermined, and that sinuses were forming. I laid open two sinuses, one on each side of the opening, and scraped the infiltrated tissues.

June 27th. The infiltration of the integuments had increased. Small subcutaneous abscesses continued to form, which intercommunicated by sinuses. Pyrexia very slight, the temperature occasionally rising to 100° .

After leaving the hospital he continued to attend as an out-patient, and I ordered liq. hyd., perchlor. and potas., iodid., but without marked benefit. Fresh sinuses formed at intervals, until the right hypochondrium and epigastrium were marked by some six or more openings, from which a thin watery fluid, but no pus, exuded. He suffered no pain. The enlargement of the liver appeared to decrease rather than the reverse. Seeing that anti-syphilitic remedies had no distinct effect, and taking into consideration the general aspects of the case, I began to suspect that the disease was actinomycosis, and on several occasions examined the discharge and scrapings from the sinuses for the ray-fungus, but without success. I continued liq. hyd. perchlor. in $\mathfrak{zj}$ doses, in the hope that it might retard the progress of the disease. He gradually got weaker and more emaciated, and finally in March, 1888, he became confined to his bed. On visiting him I found ascites, but no other change in the general condition. He died from exhaustion at the end of April.

Fig. 1.—Actinomycosis from human liver. (It is drawn rather too large).

The friends would not permit a necropsy, but I succeeded in persuading them to allow me to make an incision into the abdomen, through which I removed a piece of liver. This was studded with the characteristic nodules of actinomycosis. They had not undergone softening, and did not present the usual honeycombed