now arises, Has the animal, the recipient of a sufficient dose of these products, become refractory in consequence of their being present in the tissues, and thus preventing the growth of the microbe? Upon this point, while we do know that in cultivation the growth of certain microbes is arrested by the accumulation of the products which they form there, we must carefully avoid forming a definite opinion as to what happens in the living body upon the basis of phenomena which have been observed to take place in culture tubes. For example, if we take a little blood from a sheep which has been rendered refractory to anthrax, and place in it anthrax bacilli, they will grow there rapidly and abundantly, thus showing that there is in the blood of this protected animal no substance capable of destroying the life of the bacteria. This experiment is, of course, an extremely crude one, since from a chemical point of view there is an enormous difference between blood while still retained within the living vessels, and the same blood drawn from the body and placed in a culture flask. If any positive result were obtained, it could only be from an absolutely enormous chemical change in the composition of the tissues.

A more delicate experiment may be performed by injecting into the anterior chamber of the eye of a protected sheep a few virulent bacilli, which, while growing well in the aqueous humour, will confine themselves to that special spot. It is, therefore, evident that in this aqueous humour, notwithstanding its being a part of the animal, and consequently, of necessity, participating in its chemical modification, there is no substance present capable of resisting the local development and vital activity of the anthrax bacillus.

Besides the chemical, we have also the physiological, question to deal with, as may be seen from the following experiment. If the virus of "quarter evil," that is, *charbon symptomatique*, be injected into the thigh of a rabbit, an animal which is by nature refractory to the disease, no characteristic tumour will develop, and its immunity appears to be absolute. But if now any lesion be first produced in the tissues, either by a blow or by the injection of some caustic substance, and the inoculation be made at that point, then an anthrax tumour will soon appear, and the rabbit, though habitually exempt from the disease, may chance in this manner ultimately to succumb to it, the reason being, of course, that the injured tissues have formed an inert soil for the microbe to start its growth without hindrance. The immunity, therefore, of a rabbit from "quarter evil" does not arise from its possessing in its body any simple chemical substance inimical to the cultivation of the virus, as we see that it is only by artificially causing necrosis of the tissues that the virus is able to gain a footing in it.

What then, does happen on the injection of active virus into an animal which is refractory? What becomes of the microbes? M. Metschnikoff has taught us that they are soon destroyed, and that the foremost agents of the destruction are the white corpuscles or phagocytes, which swallow up the microbes and digest them, while, on the other hand, the white corpuscles in the bodies of non-refractory animals do not swallow the microbes, or that, if they attempt to do so, the latter develop notwithstanding.

Any satisfactory explanation of the problem of immunity should comprise all these facts, and should be based upon the relative importance of the action of the chemical products and the resistance of the tissues. I believe that the best explanation, in fact, consists in considering immunity as the result of the habituation of the cells to the poisons secreted by the microbes.

When the virus begins to develop in the body of an animal subject to the malady it forms its poisons, and when the white corpuscles begin to struggle against them their activity is arrested by this toxic production. The microbe continues to develop and the malady progresses. But in the case of an animal rendered refractory by previous injection of the soluble substances or by inoculation of attenuated virus, the corpuscles have already got accustomed to the microbic poison, and since the small doses given at the beginning of the cultivation of the virus do not impede their activity, they commence the struggle and devour the parasites. But if, as in the experiment with "quarter evil," some cause should prevent this corpuscular intervention, the cultivation of the microbe would succeed, and in this local centre there would soon be a sufficient quantity of poison to render the corpuscles powerless, notwithstanding that these have been previously accustomed to it, or possess natural powers of resistance against it.

There is no means of habituating the system to large doses. It is therefore in the time immediately following inoculation that

the decisive battle is waged. Hence the great importance of clearly understanding the importance of the condition of the seat of inoculation, and of the quantity of virulent matter introduced. When once we are thoroughly acquainted with these poisonous substances formed by pathogenic microbes, we shall perhaps be in a position to find antidotes for them capable of paralysing their action even within the tissues themselves; but in offering this suggestion I perceive that I have deserted the domain of fact for that of hypothesis, and that it is therefore time I drew my remarks to a close.

It seems to me that the foregoing view of this wide question reconciles the various works upon it, which have been multiplied in the last few years, and though it is possible that it may be modified by time, one thing will remain unchanged, and that is the gratitude of all to him who by his studies of attenuated viruses and preventive vaccinations has enabled us to deal successfully with that problem of immunity which has hitherto remained impenetrable.

THE CROONIAN LECTURES

ON THE RELATIONSHIP BETWEEN

CHEMICAL STRUCTURE

AND

PHYSIOLOGICAL ACTION.

Delivered before the Royal College of Physicians of London.

By T. LAUDER BRUNTON, M.D., D.Sc. EDIN., LL.D. ABER., F.R.S.

Fellow of the Royal College of Physicians; Assistant-Physician and Lecturer on Materia Medica at St. Bartholomew's Hospital.

INTRODUCTION.

MR. PRESIDENT AND GENTLEMEN,—Allow me before commencing these lectures to return you my most sincere thanks for the honour you have done me in asking me to deliver them, for the stimulus you have given me to collate the facts relating to a subject of the greatest practical importance, and for the substantial aid which you have afforded me in doing it. The motto of this College—"Art is long and Time is fleeting," as it has been translated by Longfellow, is one which, while it tends to repress too sanguine expectations of individual success in furthering medicine, is yet a strong incentive to individual effort, and its teaching has been well put by the same poet in the words:

Let us then be up and doing
With a heart for any fate,
Still achieving, still pursuing,
Learn to labour and to wait.

The period of waiting on the advance of medicine has indeed been a long one, and in the Goulstonian Lectures which I had the honour to give in this place twelve years ago, I tried to show to some extent why the progress of medicine has been so slow.

The Causes of Retardation.—Its advance has been mainly hindered, as I believe, because men have been guided in their practice more by their own fancies than by objective facts. I expressed a hope that the advance of medicine would become more and more rapid as the knowledge gained by close study of objective phenomena became more and more the basis of our practice. I also ventured to echo the prediction made by Dr. B. W. Richardson in 1868,¹ that, ere long, we should be able to manufacture in our laboratories drugs which would produce in the animal body any effect that we might desire, without our having to obtain them, as one might say hap-hazard, from various plants, or other sources.

ADVANCEMENT OF MEDICINE AND SURGERY.

Anæsthetics.—Already we owe a great deal to chemistry for the potent medicines which it has prepared and supplied to us. Probably the two most striking advances which have been made in the healing art during the present century are the introduction of anæsthetics and of antiseptics. Anæsthetics have not only relieved the pains of labour and lessened the agony which occurs in various diseases coming within the province of the physician, but they have taken away the terror formerly associated with surgical operations, and have allowed the surgeon, in place of try-

¹ British Association Reports, 1868, p. 186.

ing to finish his work in a few seconds, to spend minutes, and even hours, in doing operations carefully and thoroughly. They have thus not only made patients more ready to submit to surgical interference, but they have enabled the surgeon to attempt and perform with success operations which would formerly have been impossible.

Antiseptics.—The introduction of antiseptics has greatly lessened the spread of infectious diseases, and it has diminished the risk from surgical operations to an extent that is almost incredible.

INTRODUCTION OF NEW REMEDIES.

Another great advance which has been made still more recently is the introduction of various new antipyretics, some of which have the power not only of lessening fever but of relieving pain. At the present moment new drugs are being constantly introduced by the chemist to the notice of the physician—so rapidly, indeed, that it becomes a little difficult to keep *au courant* with all the recent introductions. Most of the new drugs belong to one or more of three classes, namely, antipyretics, analgesics, or hypnotics. Several of them may be said to belong to all three classes at once, and will lessen temperature, relieve pain, and produce sleep; but generally one or other action predominates and determines the class to which they are assigned, the other effects being more or less subsidiary or not existing at all. These remedies are not taken haphazard from the innumerable substances known to the chemist, but are selected from certain groups, the members of which have been ascertained to possess the desired action in varying degrees.

OBJECT OF THESE LECTURES.

In these lectures I propose (1) to try and show what principles guide us in selecting new remedies from the enormous number of substances known to chemistry, and (2) to indicate the lines upon which further work is desirable in order that we may be able to manufacture substances which will not only act upon any part of the body we wish to influence, but will produce upon it any particular effect we may desire. In order to fulfil this twofold object I shall have to mention a number of facts already ascertained; but I shall be obliged also, in indicating the directions for future research, to introduce hypotheses which require to be tested by experiment, and either proved or disproved.

The Prospects of Therapeutics appear to me very bright. At present most of the drugs which we are obtaining from the chemist belong to the classes of antipyretics, analgesics, or hypnotics; but I think it is highly probable that before long we shall have a series of drugs which will stimulate the biliary secretion of the liver or modify its glycogenic function, arranged in order of their comparative strength, in much the same way as we have now the class of antipyretics. We may also look for a series of remedies which will modify the circulation by dilating the blood-vessels, not only temporarily but more or less permanently, and will thus afford relief, not merely in transient conditions like angina pectoris, but in the prolonged high tension of chronic Bright's disease. We may also, I think, fairly expect to obtain a series of remedies having an action upon the heart and vessels like digitalis, strophanthus, or erythrophloeum, and which will possess an advantage over these drugs in one most important point. For the chemical structure of the artificial cardiac tonics will be known to us, and we shall consequently be able to modify them in the laboratory in one way or another until we secure the one which will have the particular effect desired in any given case.

SUBJECT OF THE LECTURES.

The regulations of the Croonian lectures have decided that they should be upon some subject in anatomy, physiology, or pathology in relation to the prevention, control, and cure of disease. My subject is a very wide one, for although it may be said to deal chiefly with physiology, it relates also to anatomy and pathology.

Development of Anatomy, Physiology, and Pathology.—All three subjects—anatomy, physiology, and pathology, as well as a new one, pharmacology, which may be looked upon as a subdivision of physiology—either have passed or are passing through similar stages in the course of their development.

Anatomy.—The old anatomists dealt with the organs and described the relationships of the bones and muscles, of the lungs and liver without troubling themselves about their composition. From Bichat the study of anatomy received a new impulse and the tissues of which the organs are composed became the subject of investigation. Next the anatomy of the tissues was taken up, and the importance of their constituent elements, the cells, was

pointed out with regard to vegetable tissues by Schleiden, to animal tissues by Schwann, who has only recently passed away, and in pathological formations by Virchow, who we rejoice to know is still alive and active. The anatomy of the cell itself is now beginning to receive attention. The cell is coming to be regarded as "an elementary organism with the properties of a complicated organism," and the relationship of its different parts to one another are now examined with as much care as the old anatomists devoted to the relationship of the viscera.

Physiology.—Physiology has followed a similar course to anatomy, dealing first with various functions, as related in a general way to the viscera, and afterwards with the functions of the tissues. But physiology, like anatomy, is now beginning to deal not merely with the functions of organs or tissues but with the functions of the cell. It examines now the phenomena of respiration, circulation, tissue change, motion, growth, reproduction, decay, and death, in relation to single cells. Instead of the cell being regarded as a structureless mass of proteid matter it is now considered to have a reticular structure, perhaps as complicated as the skeleton of one of the higher animals, with soft parts lying between which might be regarded as corresponding to the muscles and viscera. Instead of reproduction being a simple process of division we know that it is in many instances preceded by a most complicated series of movements within the cell, to which the name of "karyokinesis" is given, and which remind us of the movements which occur in a gas at the critical moment when it is condensing into a liquid, and are seen by passing a beam of light through it.

Pharmacology.—Pharmacology, like physiology, dealt first with the action of drugs upon general functions, and if we look into such a book as Orfila's *Toxicology*, we find it noticed that after the administration of certain drugs the respiration was accelerated, enfeebled, or stopped entirely, the circulation was quickened or slowed, or that vomiting or purging occurred, without any reference to the mechanism through which these symptoms were produced. Magendie and Bernard did for pharmacology what Bichat did for anatomy, and began to localise the action of drugs to the tissues on which they act. The number of drugs to be investigated is so great, and the work required to do it is so laborious, that even now our knowledge of the exact mode of action of most of our drugs, and even of the most important of them, is far from complete, although a vast mass of information has been accumulated.

Cell-Pharmacology.—Though much remains to be done in ascertaining the action of drugs upon organs and tissues, a new department of pharmacology is growing up—the pharmacology of the cell—corresponding to its physiology. Indeed, without the pharmacology of the cell, the study of its physiology would have been very limited, for even the highest powers of the microscope would not have enabled us to discover many of the most important points both in structure and function, were it not that various aniline compounds have been found to affect differently the various parts of cells, and we have thus been enabled not only to observe the movements which occur within the cell, but to form some ideas regarding the activity of vital processes in various parts.

The Relative Activity of Nucleus and Cell-Protoplasm.—Dr. Sheridan Delépine has made numerous observations on this subject, and has kindly furnished me with the following condensed account of the most important facts regarding it:

"The most active part of a cell, as far as essentially vital phenomena are concerned, is undoubtedly the nucleus, and in the nucleus the part of the network which Fleming has called chromatin. The protoplasm of the cell, although practically undifferentiated in young or embryonic cells and differing little from the nucleus, becomes with age distinctly differentiated by transformation into certain globulines as in the case of muscle and of the crystalline lens, or by the accumulation within its meshes of various products such as albuminoids, pigments, zymogenous substances, fats, calcareous salts, etc. Finally, some of these substances may either entirely replace the protoplasm, form a cell-wall, or accumulate between the cells, and these masses of products of cell activity, having lost all the properties of living protoplasm, can be considered as dead, or differentiated to the utmost. Thereactions of these various parts of tissues correspond to the physiological peculiarities we have just mentioned. Of the aniline dyes two great classes can be recognised: (1) acid fuchsin may be taken as a type of the first; (2) basic fuchsin as a type of the second. The main difference between these two classes is that the first

class stains deeply the most differentiated parts of the cells; the second class stains deeply the least differentiated parts of the cells. Thus if a drop of blood after proper treatment be stained by acid fuchsin, the red corpuscles will be intensely stained; the substance of the leucocytes very faintly stained; the nuclei not stained at all. If, on the contrary, basic fuchsin be used, the nuclei will be stained most deeply, the body of the leucocytes stained less, the red corpuscles least of all." In the case of the aniline colours we can see that different cells or different parts of the same cell exercise a selective power, taking one and refusing another. It is probable that the same thing occurs in the case of other substances which we cannot see under the microscope, and that strychnine, for example, unites with the cells of the spinal cord rather than with those of bone, cartilage, or muscle. What I have just called selective power is no doubt in many instances merely a chemical attraction between the components of the cells and the chemical substances brought in contact with it, for it occurs after the cell is dead. It resembles the apparent selective power of the grains of gold-dust in the process of amalgamation, for when mercury is mixed with powdered quartz, sand, and gold, the other substances remain unaffected, and it only combines with the gold. In other instances it may possibly be connected with the life of the cell, and, at all events, the alteration in colour which some of these aniline colours undergo when oxidised or reduced also render visible to the eye the chemical changes which occur in the cell, and allow us to observe its respiratory functions in the same way as the redness or blueness of the comb in a fowl, or of the nails or mucous membranes in a man, indicates the arterial or venous character of the blood and the completeness or imperfection of the respiratory functions in the fowl or in the man.

Pathology.—Turning to pathology we find a similar development occurring as in the kindred sciences of anatomy and physiology. Morgagni described the seats of disease in the organs. Virchow investigated pathology as depending upon the cells. Cohnheim threw a new light upon the process of suppuration by observations on the movement of wandering cells; and the researches of

FIG. 1.—A piece of the anterior part of the body of a daphne with a number of spores, some of which are still in the intestinal canal; others are penetrating the intestinal wall, and others are free in the abdominal cavity, where they are attacked by leucocytes. The cells which eat up others are termed phagocytes.

FIG. 2.—a. A spore which has penetrated the intestinal wall and entered the abdominal cavity, where four leucocytes have surrounded its end. b. and c. Spores surrounded by leucocytes, which have become confluent in c. d. A spore, one end of which is being digested by a leucocyte.

FIG. 3.—Different stages of the digestion of spores by phagocytes.

Metschnikoff on digestion within cells both wandering and fixed form a new point of departure for investigations on the relation between living organisms and infective diseases.

FIG. 4.—A germinating spore with leucocyte adhering to it.

FIG. 5.—a. Leucocyte of a frog eating up an anthrax bacillus. b. The same leucocyte after it has completely enveloped the bacillus. c. A connective tissue phagocyte containing three fungi cells.

We have not succeeded in tracing all infective diseases to micro-organisms, and it is possible that further researches may even modify the views now commonly held with regard to the relation of microbes to disease.

Phagocytes and Microbes.—The phenomena which Metschnikoff has observed may not be of such universal occurrence as he has supposed, but nevertheless they probably play an important part in the struggle against the disease. From them we see that a likeness may be observed between the body corporeal and the body corporate, and that the death of a mouse from anthrax may be compared with the destruction of the Roman empire by the savage hordes who invaded it. The anthrax bacilli after invading the organism enter into conflict with the cells belonging to it, and if the invaders be few and weak they are destroyed and the organism preserved, but if the invading bacilli are numerous and powerful they destroy the cells which they encounter and death of the organism is the result. Sometimes instead of completely over-

FIG. 6.—A spore germinating and forming conidia, which drop off and become free in the abdominal cavity.

FIG. 7. a, b, etc., stages in the process of a leucocyte eating up conidia.

FIG. 8.—A group of conidia which have caused the leucocytes surrounding a spore to dissolve, leaving only an empty vesicle and fine detritus.

mastering and destroying the organism into which they enter, the microbes appear to make only a local settlement where they interfere with the growth of the cells of the part attacked, and may either lead to their destruction or may possibly induce them to

take on a new manner of growth and thus lead to the formation of local tumours, or local mischief of other kinds.

Nature of Cancer.—In many instances we are able to trace disease directly to the introduction of microbes from without, but occasionally, as in cancer, we find the cells of the organism itself take on an independent action, and grow in a manner not only independent of the interests of the organism as a whole, but so as to actually destroy it. The question is a natural one—What causes the cells of the morbid growth to take on this peculiar action? And some have sought to answer it by looking in the cancer for some microbes which had induced the disease, as the ovum of an insect leads to the production of galls in the oak. The success of this search is as yet uncertain, and the statement that the bacillus of cancer has been discovered² must be confirmed by further observation before it can be definitely accepted, but it is not improbable that ere long the nature of cancer may be finally determined, and thus by and by lead up to the discovery of a cure for that dreadful disease.

Influence of Heredity.—We sometimes speak of disease as hereditary, but this is not strictly correct. The tendency of the disease is no doubt hereditary, but the disease itself is not, at least in those cases where it is due to a microbe. Such diseases require two factors for their occurrence, just as a crop of grain does for its growth. The first factor is the microbe, which may be regarded as the seed, and the second is a condition of the organism which we may compare to the soil suitable for its reception. The soil is hereditary, but the seed may or may not be sown in it; and just as a field without the seed will produce no crop, however good the soil may be, so, if the microbes be absent, disease will not occur, however great the susceptibility of the individual may be. For example, we not infrequently speak of the hereditary nature of consumption, and yet we are constantly meeting with persons belonging to very consumptive families who escape the disease by living under conditions where the bacillus tuberculosis is likely to be absent; and, on the other hand, persons, such as nurses, are in all probability frequently inhaling the microbe, and yet are not attacked by the disease. In the first case, immunity is probably due to the absence of the seed, notwithstanding the favourable nature of the soil; in the second it is probably due to the barrenness of the soil, notwithstanding the presence of the seed. But just as the seed falling on a soil adapted to its growth will germinate and yield a crop, so microbes which find within an organism conditions suitable to their growth will multiply and cause disease. As two factors are required for an attack of infective disease, namely, a disease germ and a suitable soil, it is evident that we may try to prevent it, either by destroying the germ or by rendering the soil unsuitable. Both plans are now largely employed, but the consideration of them belongs to a later division of these lectures.

Life-processes in Disease Germs and in the Human Body.—In these lectures I have, in accordance with the regulations of the College, to consider my subject in relation to the (1) prevention, (2) control, and (3) cure of disease. I may be able, to a certain extent, to treat these three separately, but the last two—the control and cure—are so intimately connected, that it is almost impossible to separate them completely. I have already mentioned that many diseases depend upon the invasion of the organism by microbes; and, in discussing the action of chemical agents in their treatment, we must consider their effect upon both factors, namely, the organism and the microbes. We were formerly accustomed to look upon mankind as entirely different and quite apart from the rest of the animal kingdom; but ideas have entirely changed; and we now see that in the lowest organisms, even in those microbes which are such potent causes of disease, the processes of life are not unlike those which go on in the human body. In both we have nutrition, respiration, growth, reproduction, decay, and death. These functions are dependent, both in man and in microbes, upon substances which may not always have exactly the same composition, but which are essentially of a proteid nature.

Concerning Protoplasm.—The name protoplasm has been a household word ever since Huxley's famous lecture in 1862, but since the time when he then sketched out with a master's hand the great features of protoplasmic life, many details have been filled up by laborious investigations. This protoplasm is able to absorb and disintegrate various substances which it uses for its

own nutrition. Within its substance respiratory functions are performed, oxygen being absorbed and carbonic acid being eliminated, and by the energy thus liberated the protoplasm may move about or may carry on processes of chemical synthesis, and build up complex organic substances from simple materials. The waste products of its tissue change are excreted and, just as the poor captives in the Black Hole of Calcutta were poisoned by the respiratory products they had themselves formed, so the lowest organisms will be destroyed if the products of their own tissue waste accumulate sufficiently around them.

"Extractive Matters."—The best known products of tissue waste in man are carbonic acid, urea, and uric acid, but in old books on the chemistry of the urine one constantly met with the expression "extractive matters." This term was used to indicate the smeary mass, resembling a vegetable extract, which remained on evaporation of the urine after the urea and uric acid had been removed. These extractive matters have now been more fully examined, and in them recent investigators have discovered powerful poisons. Indeed the amount of poison excreted by a man in four and twenty hours is frequently so great that it would have been sufficient to kill him if injected into his veins at once (Bouchard). Amongst the products of the tissue change in low organisms have been found a number of substances which destroy these organisms, or which, in other words, are antiseptic. Amongst these are phenol, phenyl-acetic acid, cresol, indol, phenyl-propionic acid, and skatol. These are all equally powerful as antiseptics, for phenol, better known as carbolic acid, is weakest, the others becoming progressively stronger in the order just mentioned.³

APOLOGY FOR THE ELEMENTARY CHARACTER OF SOME PARTS OF THESE LECTURES.

In the list of products of the tissue change of microbes, there are three names—phenol, phenyl-acetic acid, and phenyl-propionic acid—whose similarity of sound suggests a relationship to one another, but many men will not understand the nature of this relationship, either because they have forgotten the chemistry of these substances, or perhaps have never known it. For chemistry is a very young science, very little over a hundred years old, and a knowledge of the phenyl compounds is one of its newest acquisitions. The question what antiseptics have the greatest power to destroy the microbes and are at the same time least injurious to the human organism is one of vital importance. And yet if I tell you that bodies containing a diphenyl nucleus are amongst the most powerfully antiseptic and, at the same time, least poisonous substances, many of you might reply by asking me: "What is diphenyl?" On the other hand it is quite possible that many, especially of the younger Fellows of this College, who have studied chemistry recently, may laugh me to scorn for attempting to explain what is as simple to them as A B C. But it was my misfortune to study chemistry in the old days, when one spoke of equivalents, and atomic weight was not heard of. In trying to keep up, even to a very limited extent, with the progress of the science, I have had not only a great deal to learn, but a good deal to unlearn. The difficulties I have had in understanding the subject have made me try as far as possible in the present lectures to give such brief information regarding modern chemical ideas and facts as may be helpful to those who have had the same difficulties as myself, and I trust that those whose more recent and fuller knowledge renders such an attempt useless to them will nevertheless forgive me for making it.

Course followed in the Lectures.—I know it is almost impossible to apprehend many statements regarding the relation of chemical substances by simply hearing them once. I also know that, although it may be little more than a waste of time and a tax upon your patience to make such statements in a lecture, yet without a knowledge of chemical relationships it is impossible for you to understand the subject of which I am treating. I have, therefore, resolved to follow a middle course, and to put down a number of relationships in the form of tables, to which either you or I can refer at will, without being under the necessity of going over them from end to end. But while they will save me from going into details, there are some general considerations into which I must enter in order to render my subject at all intelligible to those who, like me, had the misfortune to learn chemistry long ago, and whom other engagements have prevented from keeping up with its progress.

Kiubassoff, Proc. of 3rd Gen. Meeting of Russian Med. men at St. Petersburg, 1889, 2141. Abstracted in BRIT. MED. JOURN., March 23, 1889, p. 665.

³ Wernike and Salkowski, *Virchow's Archiv*, 1879, vol. 78, p. 51.

DIVISION OF CHEMICAL SUBSTANCES INTO ORGANIC AND INORGANIC.

The composition of the universe was formerly regarded as very simple, its component elements being only four, namely, fire, air, earth, and water. Now, we regard as elements some sixty odd substances, and we give them this title because we cannot split them up by the means at our disposal. But it is possible that new methods may effect their decomposition, and then they will lose their title, just as potash did when Davy succeeded in showing it to be composed of oxygen and a metal, and it was accordingly removed from the class of elements to that of compounds. The substances which we at present regard as elements, numbering between sixty and seventy, form by their union with one another an innumerable host of compounds. These used to be divided into two great groups, namely, organic substances which were supposed to owe their origin to life, and inorganic which had no such origin. Many organic substances can now be manufactured in the laboratory, and so the old reason for their name no longer exists; but all living things, however high or low they may be in the scale of creation, contain carbon, and the name organic is now retained for compounds of carbon, and inorganic for compounds of other elements.

Organic Matter the Basis of Life.—Darwin described the evolution of the higher forms of life from the lower, and Lockyer and Crookes have described the evolution of the elements from primitive matter. But there is still a great gap between the elements and organic matter, and another between organic matter and living matter. The evolution of organic matter has still to be described, for we require its presence on the earth before we can have life at all. In another place⁴ I have discussed the various beliefs regarding the origin of living things; but first of all there must be organic matter. For, if we take the Mosaic account of the formation of animals from the dust of the ground, we still want the organic dust. If we assume that living things were gradually developed out of highly complex organic elements upon the earth itself, or suppose with Sir William Thomson that the germ of life was brought hither by a meteor from some distant planet, we still require the presence of complex organic matter in order to form the protoplasm which was to serve as a physical basis for the life which was gradually developed on the one supposition, and which was implanted, took root, and spread in it on the other.

PROTOPLASM.

The highly organised protoplasm which forms the physical basis of life has, in all probability, a most complicated chemical structure, and Plüger supposes that the relation which the size of a protoplasmic molecule bears to that of an ordinary chemical molecule is like that of the sun to the smallest meteor. We do not know how this gigantic molecule is built up, but we are learning something about its component parts by a study of the substances which it yields when decomposed. Amongst these products of decomposition we find various acids containing carbon, and compounds in which nitrogen plays an important part. If we subject any living matter to great heat we find that it becomes charred, showing us the presence of carbon as one of its most important constituents, while the unpleasant smell indicates the presence of nitrogen. The more powerful the decomposing agencies to which protoplasm is subjected, the more simple are the compounds into which it falls, until it is reduced to the elements themselves or to some of their simplest combinations.

EVOLUTION OF THE ELEMENTS.

It will probably facilitate our study of chemical structure if we begin at the very beginning and take the genesis of the elements themselves before we proceed to the compounds which they form. But it must always be borne clearly in mind that our hypotheses on this subject refer only to the how and not to the why, to the mode of making and not to the Maker.

Darwin and Lyell have taught us to look for the explanation of a natural phenomenon to regular sequences of a similar kind, and we shall probably be right if we assume that the combinations by which new substances are formed in the present day are not altogether unlike the processes by which the elements themselves were evolved, although the conditions of temperature are, of course, utterly different. At any rate it will, I think, help us if we take a look at the mode in which the elements are supposed to have been evolved, and I will try to repeat to you as nearly as

possible the simple words in which Mr. Lockyer described his hypothesis to me. This hypothesis is still regarded by many as a mere hypothesis, yet, like Dalton's atomic theory, it may be most useful in enabling us to string together known facts, as well as indicating the way to search for others yet unknown.

Let us suppose the universe to be filled with atoms of matter, each one like the other. By and by some of them coalesce, forming groups of two amongst the majority of single atoms. While some of the twos remain as they are, others, meeting single atoms form groups of three, and others, meeting double atoms, form groups of four. Aggregation goes on unceasingly until we have single atoms, groups of two, three, four, and five atoms, and so on. Moreover, differences will occur in these groups, according to the way in which the atoms are aggregated together, for some may be joined lineally to each other, while others form cubes, triangles, or irregular clumps, as shown in the accompanying diagram. Thus we may have groups of the same atomic weight, with quite different properties. The elements next combine to form meteoric dust, the dust to form meteors and the meteors to form stars and solar systems.

EVOLUTION OF ORGANIC MATTER.

Carbon Compounds.—Amongst the compounds we may include those formed by the combination of carbon with hydrogen and nitrogen, as well as compounds of these latter elements amongst themselves. That carbon may be very different in its physical characters we can see at a glance by comparing the three forms in which we are acquainted with it—diamond, graphite, and charcoal. There seem to be differences quite as great in its atomic characters. Each atom appears to have four affinities, and these may unite, either with four atoms of another element having only one affinity each, or all four affinities may unite with those of another atom of carbon, forming a molecule of carbon such as we have in charcoal, with no free affinities at all. Between those conditions we have various intermediate stages in which the carbon atoms are united to one another by three affinities, two affinities, or one affinity respectively, while the other affinities remain free for combination with other elements. The simple experiment of throwing a piece of egg into the fire or of lighting a jet of gas shows us that complicated compounds of carbon cannot exist at a high temperature. Consequently, organic matter could not exist on this earth until a comparatively late period in its history, when it had cooled down greatly from its original condition of intense heat. But let us suppose that the temperature has fallen sufficiently to allow compounds of hydrogen and carbon to exist. In the diagram to which I have referred Mr. Crookes looks upon electricity, or something akin to it, as one of the great factors in the evolution of the elements.

Acetylene.—Now if sparks be passed from carbon point to carbon point while they are surrounded by an atom of hydrogen, a new compound will be formed to which the name of acetylene

FIG. 9.—Diagram of a charcoal molecule consisting of two atoms united by four affinities of each.

has been given. In it the carbon atoms, instead of being united to one another by all their four affinities, as in the charcoal of which I have just spoken, are united to one another only by three, so that each atom has one free affinity by which it can combine with an atom of hydrogen, thus:

Acetylene

Ethylene; Methane.—Suppose that acetylene thus formed were to meet with nascent hydrogen, the affinities by which the carbon atoms were linked would be diminished first to two, and then to one. The affinities thus set free would combine with hydrogen to form ethylene,

⁴ Lauder Brunton, *The Bible and Science*. London, Macmillan and Co.

Nucleoplasm; nucleus Protoplasm; body of Periblast; red blood cor-
of white blood corpus- white blood corpus- puscles.
cles.

FIG. 10.—Diagram to show the relative intensity of staining of more or less differentiated proteid matter. The background represents the blood plasma.

1. Aniline blue-black (sloelin) and acetic acid.
2. Acid Magenta.
3. Eosin (tetrabrom-fluorescein), $C_{20} H_3 Br_4 O_5$.
4. Sloelin (aniline blue-black), $C_{30} H_{25} N_5$?
5. Gentian violet.
6. Rosein.

Benzene.—You will notice that in the three substances just mentioned—acetylene, ethylene, and methane—the carbon atoms unite like the links of an open chain, and these bodies all belong to one of the great classes into which carbon compounds can be divided, the so-called fatty or alcoholic series, because oils, fats, and alcohol belong to it. But when acetylene is strongly heated

the carbon atoms unite in another way, so as to form a closed chain in which the carbon atoms of acetylene are supposed to

FIG. 11.—Diagram of the union of three molecules of acetylene to form one of benzene.

have as it were opened out their connecting affinities, so that no two are connected by three affinities, or bonds, as they are often termed, but are alternately connected by two, as in the accompanying figure.

Three molecules of acetylene unite to form one of benzene, and this substance, in which the six atoms of carbon form a closed ring or nucleus, as it is termed, is a type of the second great class of carbon compounds—the aromatic series.

The Intermediate Class.—Between the two great classes of carbon compounds which I have just mentioned, the fatty series in which the compounds are linked so as to form an open chain, readily broken, and the aromatic series in which the carbon atoms form a closed chain or ring which is difficult to break, a third or intermediate series may be placed. In this the carbon atoms form a close chain, but instead of this consisting of six carbon atoms closely linked together, as in the aromatic series, it contains only three or four carbons, either linked to one another or having the chain completed by an atom of oxygen, sulphur, or nitrogen. As examples of this series we have

Three Classes of Carbon Compounds.—We have thus three great classes of carbon compounds: 1. The fatty or alcoholic class, with an open chain of carbon atoms; 2, the aromatic class, with a firm closed chain or ring containing six carbon atoms; 3, an intermediate class.

(To be continued.)

POOLE (Population, 13,781).—**Measles Epidemic: Question as to best Method of Sewage Disposal.**—The returns of this borough for the year 1888 are not quite so favourable as those of the previous year, there being a decrease of 58 births and an increase of 25 deaths compared with 1887. Mr. H. A. Lawton reports that there was an epidemic of measles during the spring and summer, and that the wet summer followed by the warm autumn gave rise to an outbreak of typhoid fever. Not a single death was caused by infantile diarrhoea, the 3 fatal cases of diarrhoea recorded being among aged people. One death was attributed to dysentery. Phthisis caused 24 deaths, bronchitis and other lung diseases 47, and heart disease 14. The death-rate from all causes was 17.47 per 1,000 of the population. There are several questions requiring the attention of the sanitary authority, chief among them being the method of disposing of the sewage. The by-laws also seem to need entire revision.