MEDICAL PARIS OF TO-DAY.

NOTES MADE IN DECEMBER, 1888.

By ERNEST HART.

XV.

DR. DÉJÉRINE AT THE HOSPICE DE BICÊTRE.

Cases illustrative of Nerve Pathology: 400 Beds: Fibrous Rheumatism of Nervous Origin: Cases of Paget's Disease: A File of Ninety Ataxic Patients: Symptomatology of Progressive Atrophic Myopathy; the Myopathic Facies: Thomsen's Disease: Male Hysterics and Hypnotics: the Diathesis of Contracture: Psychical Paralysis (ab irâ).

THE Hospice de Bicêtre or Hospice des Vieillards et Incurables (men) comprises various categories of wards; some are devoted to cases of mental alienation, epilepsy, or idiocy; others to the surgical and medical affections of old men. Nowhere except in Paris, and there only at the Salpêtrière, have I found any service comparable with this for its richness in illustration of affections of the nervous system. I shall speak for the moment only of the medical wards under the charge of Dr. Déjerine. They afford a mass of material which furnish the elements in the hands of this accomplished physician for a highly

various nature, and specially from affections of the nervous system. The old men are there by reason of their age, and come to end their lives. The chronic patients come there as incurable, and remain there also frequently till death. These are the two categories—old men and cases of chronic disease in the male—which furnish the patients in Dr. Déjerine's service. The Bicêtre receives only men.

His medical wards comprise about 200 beds, a considerable proportion of which I found occupied by patients suffering with interesting and often rare affections. Let me add that many of the nervous patients do not occupy the infirmary itself, but remain in other wards connected with this service in which regular visits are not required. These patients, who may be estimated at about 200, are nevertheless at the disposal of the physician for the needs of his *clinique*, etc. The whole number of patients affected with maladies of the nervous system and belonging to these medical wards reaches, therefore, about 400. It will not be without interest to students of neuropathology who may propose to visit Paris if I begin by briefly enumerating some of the patients affected with different chronic affections of a rare and interesting kind whom I specially noticed in my visit.

Fig. 1.—General hypertrophy of the bones (Paget's disease).

interesting series of studies. To explain the organisation of this service, the method of recruiting its wards must be mentioned. Besides the insane, imbecile, and epileptic who constitute, as I have mentioned, the special services, there are at Bicêtre aged men and adults of all ages suffering from chronic affections of

Fig. 2.—Atrophic myopathy (facio-scapulo-humeral type).

Among them was, first, a case of chronic fibrous generalised rheumatism attacking all the articulations of the body, including those of the vertebral column and of the jaw in a man about 50 years of age. This was accompanied by muscular atrophy, with

glossy skin, by vitiligo and interstitial nephritis. This, then, was probably a case, as M. Déjerine pointed out, not of ordinary chronic rheumatism, but much more of an affection of nervous origin, and of a kind yet undetermined.

Two other cases (1 and 2) exhibited general hypertrophy of the bones (Paget's disease), a very rare affection, the pathological anatomy of which is as yet little known, and which in these cases presents the following characters: considerable hypertrophy of the bones, especially of the long bones, going so far as to more than double their diameter; determining in some of them (the humerus, femur, tibia) anomalous curvature in consequence of unequal deposit of bony substance of new formation in the whole circumference of the bone. In one of these patients the cranium, as well as the lower maxillary bone, had acquired considerable dimensions. The evolution of this singular affection is very slow, but continuous. It began to appear in these two patients at about the age of 40. It is especially the considerable number and great variety of diseases of the nervous system which constitute the most interesting study of the service of M. Déjerine.

I passed in review about ninety ataxic patients, presenting all possible forms of that affection. Some of them (about fifteen) are blind in consequence of complete papillary atrophy, remaining, however, in the first period of tabes, and thus confirming the opinion of Benedikt of Vienna that blindness from papillary atrophy arrests the development of the sclerosis of the posterior columns. Others present visceral symptoms, gastralgic, vesical, ureteral, nephritic crises, etc. Others, again, are attacked with strongly-marked muscular atrophy—atrophy due not to polio-myelitis, but to peripheral motor neuritis, as M. Déjerine has recently shown.¹ We examined several of these atrophic ataxic patients; the atrophy in these different cases affects sensibly the same topography, predominating throughout, as M. Déjerine pointed out, in the muscles of the extremities, in particular in the small muscles of the hands and feet, whence the production of deformities and very pronounced vicious attitudes of the extremities of the limbs.

I may mention, too, a rather rare case of unilateral ataxy. The proportion of syphilitic persons among the ataxics of the Bicêtre is about half. To finish with locomotor ataxy, let me further cite a very well-marked case of Friedreich's disease (hereditary ataxy), presenting the deformity of the feet which has so often been observed in this disease. I may mention, without stopping to describe them, a very great number of patients, hemiplegic from various causes, of whom some, however, are interesting from the existence of muscular atrophy on the hemiplegic side. About twenty among these hemiplegic patients had been attacked with hemiplegia in the earliest years of their existence (infantile hemiplegia). These patients are all atrophied on the corresponding side of the body. In some the hemiatrophy of the face is particularly pronounced; others present athetosis in the extremities of the paralysed limbs. Let me mention, too, several cases of motor aphasia, and a case of sensorial aphasia, verbal blindness with homonymous hemianopsia.

The various forms of paraplegia are largely represented; we saw many cases of spasmodic tabes of syphilitic origin, of uni- or bilateral paraplegia, due to injuries of the spinal column. I may specially mention a case of paraplegia, with muscular atrophy and disturbance of sensibility due to compression of the cauda equina, consecutive to traumatic dislocation of the fourth lumbar vertebra. If the vertebral column of this patient were not examined it might have seemed, from seeing him walk, that it was a case of alcoholic paralysis. Further may be mentioned a very remarkable and very complete case of hypertrophic cervical pachymeningitis, an affection first studied by Gull. The report of the case of the patient was published some years ago by Vulpian, in his *clinique of La Charité*.

In the wards of this service may be seen the most varied forms of cerebro-spinal sclerosis *en plaques* and of Parkinson's disease.

Progressive muscular atrophy is represented by upwards of a dozen patients. It is known that since the published work of MM. Landouzy and Déjerine the classification of the different species of progressive muscular atrophy has been greatly modified, and that these authors have established that muscular atrophy of myopathic origin (the progressive atrophic myopathy of Landouzy and Déjerine) is much more common than muscular atrophy of myelopathic origin. They have shown, moreover,

how difficult it is to establish a differential diagnosis between these two species when the muscles of the face do not participate in the atrophy. The cases of progressive atrophic myopathy were very interesting. Three of them presented the type which has been designated by the name of facio-scapulo humeral type (Illustration 2). The atrophy of the muscles of the limbs is extreme in these three cases. It dates back to the earliest years of infancy. It is more pronounced in the muscle roots of the limbs than in those of the extremities. It is accompanied in one of the patients with muscular retraction, immobilising the limbs in a vicious position, but no traces exist of muscular hypertrophy. It is important to study the *facies* of these patients, for if one were not forewarned there would be a great chance of not recognising it, and of making thus so far an imperfect diagnosis. The "myopathic facies" (Illustration 3), as Landouzy and

Fig. 3.—The myopathic facies.

Déjerine have designated it, is characterised by a special flaccidity of the physiognomy, which is wanting in expression. They often give to the patient an unintelligent look. The patient can neither completely close his eyes nor wrinkle his forehead or his eyebrows; he cannot whistle, and when he is made to laugh he laughs transversely; often the upper lip bulges forward ("lip of the tapir"). The affection is essentially hereditary, but may, however, skip a generation. Its evolution is very slow, its progress fatal. Often, however, in these patients the progress of the atrophy may be so slow that the affection remains, so to speak, stationary for a long time.

Thomsen's disease is represented in this service by a very remarkable case. This name is used, as is known, to designate an affection of very peculiar symptomatology, characterised by a state of special rigidity of the muscles when the will puts them in motion. If one of these patients is made to execute any movement, the muscles are observed to be seized with a sort of contraction, lasting for a certain time before the phases of decontraction occur. In this patient the muscles of the lower limbs are affected with considerable hypertrophy, which is frequent in this affection. Here, as is always the case in Thomsen's disease, the affection is congenital; but, contrary to rule, one

¹ J. Déjerine: Etude Clinique et Anatomico-pathologique sur l'Atrophie Musculaire des Ataxiques (Névrite Motrice Périphérique des Ataxiques) *Revue de Médecine*, February, 1889.

does not find in these cases similar heredity in the antecedents. I may add finally that in this patient there are certain special electric reactions, among others the myotonic reaction indicated by Erb. Thomsen's disease, now well known from the clinical point of view, is as yet only very incompletely studied from the point of view of pathological anatomy. No necropsy has, I believe, yet been published of this affection. The only information which we possess comes from Erb, who, by excising small fragments of the muscle in a patient, has ascertained the existence of considerable hypertrophy of the primitive fasciculi, with slight thickening of the connective tissue. The state of the peripheral nerve and of the central nervous system is still unknown.

There are in these wards a great number of neuroses of various kinds. First may be mentioned several cases of chorea of the aged, and several cases of functional spasm of various symptomatology. Hysteria of the male is represented by some very remarkable types, among others, by a young man, aged 21, hysterio-epileptic, with sensitivo-sensorial hemianæsthesia of the left side. This patient is very remarkable from another point of view. He is, indeed, hypnotisable and suggestible to a very high degree; but one does not observe in him the three phases described by the school of the Salpêtrière; and the symptoms which he presents in the hypnotic state appear conformable to the ideas put forward in this respect by the school of Nancy. In any case it is one of the most perfect examples that can be met with for the study of suggestion and cerebral automatism. I saw produced in him, besides psychical and traumatic paralyses, a whole series of phenomena of the highest interest from the psychological point of view; duplication of personality; reversion in consciousness to an anterior period of his life; suggestions of various kinds of the most striking effect are produced in him with surprising rapidity and precision. Such cases, which have been much more carefully studied in France than elsewhere, present not only considerable psychological interest (as such a case as this shows), but an interest of marked character from the medico-legal point of view. It is certain indeed that a subject such as this, going through the extraordinary phases of mental impression which can be witnessed in him, is from the social point of view an eminently dangerous being, not from his own impulse, but because, from the very character of his neurosis, an evil-minded person might use him as an instrument, and make him do almost anything that he wished.

In speaking of hysteria, there should be mentioned, also, the case of a man, 59 years old, the subject of attacks of hysteria, and suffering from pronounced left hemiplegia, with sensitivo-sensorial hemianæsthesia on the same side—phenomena which have existed for the last fifteen years. Another patient, equally subject to attacks of hysteria, is not less interesting from other points of view. He is a man, 30 years of age, suffering with blindness from papillary atrophy, arising probably from a cerebral tumour. This man is affected with what the school of the Salpêtrière has designated under the name of the diathesis of contracture. When he takes hold of an object, he cannot let it go; nor can he open his hands when you close them by passive flexion; in order to open his hand in either case, it is enough to slightly rub the back of the hand or the muscles of the posterior region of the forearm. This patient presents, moreover, a symptom which is rarely observed out of the hypnotic state, namely, neuro-muscular hyper-excitability in the waking state. This is an interesting peculiarity, and tends to prove that the different symptoms which this man presents were probably connected with the mechanism of auto-suggestion.

M. Déjerine showed me another case which has a more complicated interpretation; it is that of a man, 46 years of age, having monoplegia of the inferior left limb, with left sensitivo-sensorial hemianæsthesia, and loss of muscular sense. The case is not one of organic lesion of the brain, but it is impossible to say that it is certainly due to hysteria, for sensitivo-sensorial hemianæsthesia does not necessarily call for the diagnosis of hysteria. The patient, indeed, does not present the stigmata of hysteria, and is not hypnotisable. To conclude, among the series of paralyses *sine materid* may be cited a case of paraplegia, which has lasted for ten years in a very vigorous man, about 40 years of age. This paraplegia occurred suddenly after a fit of rage. Here also hysteria cannot be affirmed, this man not presenting any other symptom of that affection. The case is all the same one of psychical paralysis, and the paraplegia, which improved at first, became as intense as ever again after a fresh fit of anger. I may add, finally, that there is a laboratory annexed to the service

of Dr. Déjerine, containing a great number of pathological preparations illustrative of the anatomy of the nervous system, and a photographic service which supplies the *clinique* with a rich series of illustrations of the phases of disease in various stages of progress; and to which I am indebted for the reproduction of some of the most striking cases which I saw, and from which I have selected the materials for the illustrations here furnished. The necropsies in this service number from 300 to 350 annually. I have thus enumerated rapidly some of the most interesting cases met with in this service. From the point of view of its richness of material, I do not think that outside the Salpêtrière there can be found in any of the great cities of Europe so rich a collection of all possible varieties of nervous disease; nor do I doubt that this opinion will be confirmed by those who may visit this hospice. I can but advise neurologists and physicians in general to make such a visit, for they will find there the most desirable facilities for completing their studies in neuropathology, and in Dr. Déjerine an accomplished and courteous clinician and professor.

ROYAL COLLEGE OF PHYSICIANS.

THE annual meeting of the College for the election of a President was held, as required by statute, on the 15th instant, being the Monday after Palm Sunday. There was a large attendance of the Fellows.

A report was received from the Building Committee with respect to the public statue of the Queen, in which it was stated that the statue was approaching completion, and that it was hoped that some member of the Royal Family might consent to unveil it when erected.

The PRESIDENT, in accordance with custom, read an address to the Fellows, in which he commented on the life and work of the eight Fellows deceased during the last twelve months (the best-known of these being Dr. Greenhow and Dr. C. J. B. Williams); drew attention (especially with reference to recent occurrences) to the methods of action adopted by the Censors' Board; and referred to the newly-presented portraits of the two foregoing Presidents, the late Sir George Burrows, Bart., and Sir William Jenner, Bart., adding an eloquent testimony to their character and devotion to the College. Reference was made to the progress of the new building behind the Examination Hall, and to the committee appointed to consider the arrangements for scientific medical work therein, also to the committee recently nominated for the purpose of rendering, if possible, the fever hospitals of London available for clinical study. The address concluded with a strong protest against the difficulties, which of late have been greater than ever, in the way of investigation by experiment on animals. Finally, the President announced, to the regret of every Fellow present, that Sir Henry Pitman, after thirty years' service as Registrar, desired permission to resign his office, but that it was hoped that some plan might be adopted whereby his wisdom and experience might still be available to the College.

The address was received with much applause; and the regulations for the election of a President having been read, the ballot was proceeded with. The result was that Sir Andrew Clark was almost unanimously re-elected, having obtained 112 votes out of 116. He was again formally installed, and promised to maintain the rights and privileges of the College. In a few appropriate words he thanked the Fellows for the honour again bestowed on him.

THE ROYAL COLLEGE OF SURGEONS AND ITS MEMBERS.

At an ordinary meeting of the Wigan Medical Society, on April 11th, the following resolution was submitted by the President, and carried *nem. con.* "That this Society cordially approves of the Bill about to be presented to Parliament, by which it is proposed to restore to the Members of the College of Surgeons a free and unrestricted exercise of their ancient rights and privileges."

At a meeting of the Cardiff Medical Society held on Thursday, April 4th, Dr. Alfred Sheen, President, in the chair, on the motion of the President, the following resolution was passed, and a copy ordered to be sent to the Association of Fellows and Members: "That this Society cordially approves of a Bill being presented to Parliament by which the free and unrestricted exercise of the ancient rights and privileges of the Members of the College of Surgeons of England shall be secured."