natural life on healthy, natural food; in marrow from the long bones of the ox; and in ox-blood, served up in the popular form of black puddings. Snail-oil, also, and linseed-tea were commended as salutary, if less palatable, remedies. With this line of treatment was conjoined the use of such herbs as might stimulate healthy appetite and promote healthy digestion; while in the later stages of the disease that great factor of folk skill called "rule of thumb" had somehow stumbled into the use of such simple remedies as modern science now guides us to prescribe in aid of the vicarious action of reins and liver.

ON INFLAMMATORY DISEASE OF THE SEMINAL VESICLES.

By JORDAN LLOYD, M.B., M.S., F.R.C.S.,

Surgeon to the Queen's and Children's Hospitals, Birmingham; Visiting Surgeon to the Birmingham Workhouse Infirmary.

A FEW years ago the term inflammatory diseases of the Fallopian tubes was scarcely to be found in the clinical vocabulary of practical surgeons. Such conditions were casually referred to by erudite pathologists rather as morbid possibilities than as everyday realities. During the past few years, however, this latter class of maladies has had "daylight let into it," and now stands revealed to us as a common incident in the morbid history of civilised woman. Of the liability of the Fallopian tubes to inflame there can be no doubt, and of the part played by gonorrhoea in initiating salpingitis most authorities are agreed. Whether time will prove that the analogous structures in males are similarly prone to affection I do not know, but I am already convinced that inflammatory diseases of the sexual apparatus lying at the base of the male bladder are far from uncommon, and that in seminal vesiculitis the key is to be found to much that is puzzling in many of the so-called prostatic and bladder disorders. A brief reference to the anatomy of the parts concerned will not, I trust, be considered altogether out of place.

The vesiculæ seminales are two membranous receptacles, situated one on each side, beneath the base of the bladder between it and the rectum. Their length is usually about two inches, and their greatest breadth from four to six lines; but they vary both in size and shape in different individuals. Their posterior extremities are separated widely from each other, but anteriorly they converge so as to approach the two vasa deferentia, which run forwards to the prostate between them. With the vasa deferentia thus interposed, they occupy the two diverging sides of the triangular portion of the base of the bladder, which lies upon the rectum, and is bounded behind by the line of reflection of the recto-vesical fold of the peritoneum. The seminal vesicles themselves rest upon the rectum, but are separated from it by a layer of the recto-vesical fascia which attaches them to the base of the bladder. Their posterior ends lie beneath the openings of the ureters.

The common seminal or ejaculatory ducts, two in number, are formed on each side by the junction of the narrowed extremities of the corresponding vas deferens and vesicula seminalis, close to the base of the prostate. From this point they pass side by side through the prostate between its middle and lateral lobes. After a course of nearly an inch they end in the floor of the prostatic portion of the urethra by two valve-like slits placed in the veru montanum, one on each prominent margin of the opening of the prostatic sinus.

Seminal vesiculitis is usually secondary to mischief in the urethra. It is a common accompaniment of gonorrhoeal epididymitis, and originates in a precisely similar manner. When the inflammatory process has crept from the urethra along the common ejaculatory duct to its termination, it is as likely to proceed along the short seminal tube to the vesicle, as along the twenty-four inch vas deferens to the epididymis. I believe it commonly extends along both these structures.

I have seen the vesicles inflame secondarily to urethritis, set up by the passage of a bougie, by the presence of a stricture, and after coitus with a leucorrhoeal woman. I have seen both vesicles suppurate in one case where their inflammation appeared to be primary; at least there was no antecedent cause discoverable in the urethra.

When acute inflammation attacks a vesicle it gives rise to a

swelling at the side of the base of the bladder, the greater part of which is due rather to effusion of inflammatory products into the perivesicular connective tissue than into the cavity of the vesicle itself, just as we see that in epididymitis the bulk of the enlargement depends upon inflammatory infiltration into the connective tissue between the tubules of the epididymis.

Seminal vesiculitis, like acute epididymitis, most frequently terminates in resolution. It sometimes ends, however, in suppuration, and, when this occurs, pus may make its way laterally into the ischio-rectal fossa, or may diffuse itself deeply around the rectum (constituting one of the varieties of peri-rectal suppuration), or may discharge itself by the ejaculatory duct, or may open either into the bladder or rectum, but never into both cavities together.

The following case is one of suppuration of both seminal vesicles without an antecedent urethritis.

On May 23rd, 1888, I was asked by Dr. Ravenhill, of Bordesley, to see a case of acute prostatic abscess. The patient was a healthy man, aged 30, married, three children, wife again eight months pregnant. He had never had any venereal disease. He felt pain in the rectum about twelve days before, which came on without any known cause. Within forty-eight hours he had difficulty with his water, with frequent desire to micturate and pain at neck of bladder. There was a cloud of mucus in healthy urine. He had no orchitis. With a finger in the rectum I found on the right side a deep-seated, smooth, round, fluctuating swelling, with its longest diameter in the direction of the bowel, and its lower end about one inch and a half from the anal margin. Pressure on this was painful, and gave rise to a desire to micturate. Under chloroform I made a puncture through the perineum, inserting a long-bladed knife about one inch to the right, and in front of the anus, and guided by the finger in the rectum, entered the blade to the depth of two inches, tapping a cavity containing about four drachms of pus. A finger passed through the wound entered a cavity which lay altogether to the right of the middle line, up towards the sides of the pelvis. The cavity was drained, and all symptoms at once disappeared. The wound healed up completely in eight or nine days, and for three or four days he appeared to be perfectly well. He then developed a train of symptoms precisely like those of his former illness, and on June 8th I was asked to see him again. I then found an elongated fluctuating swelling on the left side of the base of the bladder, the right side being perfectly normal. Under chloroform I made a deep puncture in the left side, and entered a cavity as large as a hen's egg, containing thick yellow pus. A finger exploring the cavity found it close to the base of the bladder, and passing upwards so that the finger tip could almost be felt from the left groin above. The cavity was drained, and in two or three weeks had soundly healed.

Some cases of vesiculitis pursue a chronic course, and distend the cavity of the vesicle, a condition implying necessarily permanent choking of the ejaculatory duct, and giving rise to a cystic swelling behind and beneath the bladder. We are familiar with a similar state of things in connection with the gall-bladder, and in hydro- and pyo-salpinx. A case of this kind is reported by Dr. Smith, of Baltimore, in the *Lancet*, vol. ii, 1872, p. 559, under the title of hydrocele of a seminal vesicle. Dr. Smith's patient had a cyst behind the bladder and in front of the rectum which completely filled the pelvis, and reached up to the umbilicus. An ounce of urine having been drawn off by a long catheter, the cyst was tapped *per rectum*, and ten pints of brown serous fluid withdrawn.

Dr. Ralfe describes¹ the case of a sailor, aged 25, whose symptoms were of five months' duration. He had a cystic tumour filling the pelvis, pressing on the rectum, and presenting in the middle line of the abdomen above the pubes. After death this was found to be due to a large cyst connected with the left seminal vesicle.

Another somewhat similar case is recorded by Mr. Mitchell Henry in the *Medico-Chirurgical Transactions*, vol. xxxiii, p. 307. A sailor, aged 20, was admitted into the Middlesex Hospital with severe pains in the loins and left hip-joint, attributed to exposure to a severe wetting six months before. He had never had syphilis or gonorrhoea, and had usually no difficulty in micturition. The urine, specific gravity 1029, was thick, scanty, and loaded with pus. Hæmaturia followed, and a little later symptoms of peritonitis, which proved fatal. *Post-mortem* examination showed general early peritonitis; bladder contained purulent fluid, and

¹ *Lancet*, vol. ii, 1876, p. 782.

was perforated by an ulcer as large as a sixpence. Pelvis and hip-joints were healthy. The remains of a large abscess were discovered in the position of the left seminal vesicle, which was nearly destroyed by suppuration. The urethra and other genital organs were healthy.

Another case of the kind was communicated by Mr. Cock to Mr. Henry, in which it is probable that an abscess formed in one of the vesicles, a fatal result having been prevented by a timely evacuation of the pus by an incision through the rectum.

I have had no opportunity of testing the accuracy of my own clinical observations by pathological examination, because no death from gonorrhœa or its early complications has occurred in my practice. A search, too, in the literature of the subject has not furnished me with much positive evidence on the point. In the *Medico-Chirurgical Review*, vol. i, 1857, p. 270, a reference is made to 48 cases of orchitis, under the care of Velpeau, of which 46 were cured, 1 was discharged uncured, and 1 died. In this case the orchitis was not very severe, and succeeded to a mild gonorrhœa contracted by a young man with hypospadias. He died of peritonitis, and on examination all the seminal passages were found to be the seat of blennorrhagia. The seminal vesicles were in a state of suppuration, and the peritonitis had originated at the recto-vesical *cul de sac*. This is a rare case, but M. Velpeau had long since observed that inflammation of the seminal vesicles is by no means a rare affection in gonorrhœa.

The symptoms of vesicular disease are essentially those of vesical irritability—increased frequency of micturition, attended by more or less pain. They are like those of prostatitis, according to Professor Humphry, and consist in uneasiness about the perineum, painful defæcation, frequent and rather painful micturition or retention, painful emissions at night, bloody semen, persistent gleet discharge, and irritability of the bladder and the sexual organs. Zeissl says that the subjective symptoms differ but little from those of prostatitis; but there is one symptom which belongs exclusively to this disorder, namely, erections are well-nigh constant, and so painful as to constitute priapism. According to Lallemand, Gosselin, and Pitha, involuntary seminal emissions occur.

Objective symptoms are, in my experience, much more reliable, and are easily elicited by careful methodical examination. A finger in the rectum recognises an elongated swelling beyond the prostate, running obliquely upwards and outwards at the side of the base of the bladder. This swelling is made quite evident to the examining finger when a metal bougie is passed into the bladder and moved from side to side across the tumour. I have demonstrated a distended vesicle in this way, and have verified the diagnosis by aspirating seminal fluid through the rectum. The following case illustrates several of the above-mentioned facts:

H. J. T., aged 27, was seen with Dr. Leech; he had been under treatment several months with symptoms of irritability of the bladder. He complained of frequent micturition, twelve or fourteen times daily and three or four times at night. He was in constant pain referred to the root of the penis; this pain became aggravated as the bladder distended, and subsided again after micturition. The patient was married, and the father of two children, his wife being again eight months pregnant. He had no rectal symptoms of any kind; had had nocturnal emissions at least once a week during the past six months; had been troubled also during this time by too frequent erections. Last intercourse a month ago; pains unaffected by it. Had had pains across loins, more particularly on right side. Had ridden a bicycle for five or six years, and had noticed that the point of the saddle pressed unduly on his perineum and produced a desire to micturate. Urine acid, 1020, free from albumen and deposit; no hæmaturia or urethral discharge at any time. The bladder was sounded negatively, and a finger *per rectum* discovered tenderness at vesical base. A few days later—three days after a nocturnal emission—I gave him chloroform, and carefully explored the bladder with Leiter's endoscope. The mucous membrane looked perfectly healthy in every part. I then passed a finger into the rectum, and could feel an elongated swelling, commencing just beyond the prostate and running upwards and outwards for about three inches. This swelling was fluctuant, smooth, and the rectal wall was movable over it. I then passed a bougie into the bladder, and by combined manipulation between it and the finger I could detect most easily an elongated fluctuating tumour between the two. Aided by a Hilton's speculum I aspirated 2 drachms of characteristic seminal fluid—leaving

much more behind—by puncture through the rectum. Microscopical examination of this fluid showed large, roundish, opaque bodies and numerous motionless spermatozoa. The patient was relieved temporarily of his pains by the operation.

So far as I am aware, this is the first case where the diagnosis of a distended seminal vesicle has been confirmed in this manner. The microscopic examination put the matter beyond dispute. In the absence then of all trace of disease in the bladder—and by no other way than by visual inspection with an endoscope can we positively determine this—and in the absence of disease in the rectum, I think we are justified in looking upon the distended vesicle as the cause of this patient's symptoms. Nor are these symptoms difficult to explain. The distended vesicle must produce a certain pressure on the neck of the viscus in the neighbourhood of the trigone, and this pressure excites the frequent desire to micturate. The pain increases as the bladder fills, because the pressure on the vesicle is increased with the distension of the viscus above it. The cause of the distended vesicle is to be found in the obstruction of its outlet; the occurrence of a copious nocturnal emission three days before my examination negatives mere physiological distension. The absence of primary urethral inflammation make it less clear to determine the wherefore of the duct's obstruction, but I think it probable that the pressure of the elongated front of a bicycle saddle may have had an influence upon the most anterior part of the ejaculatory tubes. The patient had noticed that bicycle riding had excited a frequent desire to micturate. Nocturnal emissions and undue erections were both present. The occasional lumbar aching, of which he complained, might be due to the pressure exerted by the distended vesicle on the overlying ureter, damming up as it were for a time the urine in the kidney pelvis.

Gonorrhœal vesiculitis occurs at least in two forms, one mild and the other severe. The mild cases are as common as gonorrhœal epididymitis, and are distinguished by symptoms usually attributed to the neck of the bladder or the prostate, such as deep-seated aching, and sensation of fullness in the perineum; irritability of the bladder with healthy urine, more or less relief to discomfort after micturition; absence of scalding and of urethral discharge, slight increase of troubles during defæcation. With symptoms such as these, a finger passed high up the rectum recognises heat, tenderness, and swelling at the base of the bladder, beyond the position of the prostate, and the diagnosis is in this way very easily made. These mild cases usually subside without any other treatment than that suggested by the gonorrhœa.

The more acute variety is well illustrated by the following case. At the beginning of this year I saw a middle-aged gentleman, who, three weeks before, had had intercourse with a prostitute, and eight days subsequently began to develop symptoms of an ordinary gonorrhœa. A week later the discharge rapidly lessened in quantity, and he felt deep-seated pain in the perineum. He then consulted his medical attendant, who regarded his condition as one of acute prostatitis, and owing to the large amount of swelling felt about the base of the bladder, my opinion was sought as to whether there was not some deep collection of pus to account for it. The symptoms at this time were micturition every hour; pain constant, increasing as the bladder distended, and being always relieved after micturition. Rectal and vesical tenesmus. Urine neutral and free from mucus, blood, and pus. Temperature slightly raised, defæcation painful. Finger in rectum found the sphincter slightly contracted. Parts about base of bladder swollen and bulging backwards to sacrum and flattening the bowel. Swelling was hard and exquisitely tender, particularly laterally and in the neighbourhood of the seminal vesicles, where there were distinct lateral bulgings: no pain or swelling in scrotum. Under treatment by hot hip baths, leeches to perineum, hot water enemata and morphine suppositories, the symptoms rapidly subsided, and three weeks later he was well enough to go to the sea.

There are many other interesting facts relating to inflammations of the deep urethra and its connections, to which I should like to call attention, but the time allowed for my paper is already exceeded. The conclusions I am led to by my present experience are as follows:—

1. That inflammatory disorders of the seminal vesicles and their ducts are not uncommon.
2. That they are in many respects analogous to inflammatory diseases of the Fallopian tubes in women.
3. That while occurring sometimes primarily, they are as a rule secondary to inflammation of the urethra.

4. That the ejaculatory ducts may become obstructed, and the seminal vesicles consequently hyper-distended.

5. That termination by suppuration is exceptional.

6. That when suppuration occurs it should be dealt with by incision from the perineum rather than from the rectum.

7. That gonorrhœa is by far their most common originator.

8. That they are frequently concomitant with gonorrhœal epididymitis.

9. That they are usually diagnosed as inflammation of the prostate or neck of the bladder.

10. That while certain subjective phenomena are suggestive of these disorders, their diagnosis can only be made by objective examination from the rectum and bladder.

Since my attention has been particularly directed to this class of diseases, I have been struck by the frequency of their occurrence, and although my paper falls short of what I could have wished, and will not contribute much to our stock of knowledge, I venture to hope that it may direct the attention of practitioners to a class of disorders which have hitherto received too little attention.

ANGULAR CURVATURE; PARAPLEGIA; RESECTION OF LAMINÆ; RECOVERY.

By W. ARBUTHNOT LANE, M.S.,

Assistant-Surgeon to Guy's Hospital, and to the Hospital for Sick Children,
Great Ormond Street.

A BRIEF description of the following case of paraplegia due to angular curvature, in which the laminæ of several vertebræ were removed, may be of some interest, both on account of the comparative rarity of this mode of treatment at the present time, and also because of the success which followed operative interference. I am indebted to my dresser, Mr. F. G. Swayne, for the report of the case.

H. S., aged 7½, was admitted into Job Ward under my care on February 4th, 1889. His father suffers from cough and hæmoptysis. He has lost three children with chest complaints. There is also a phthisical history in the mother's family. The boy had bronchitis every winter since birth. He had scarlatina when 4 years old, from which he recovered perfectly. Eight months afterwards he became weak in his legs, but nothing abnormal was observed in his back. This weakness gradually disappeared, and he was able to run about and attend school.

In January, 1888—thirteen months ago—his parents observed that his back was prominent between the shoulder-blades. He complained of no pain, either when the lump was pressed upon or on his moving about. By the advice of a medical man he was kept on his back. In March the parents observed that he was unable to move his legs; and in April, when his legs were extended they became suddenly and spasmodically fixed in that position, the feet being forcibly pointed. He was able to control his motions more or less perfectly, and he was conscious of a desire to micturate; it was, however, necessary to attend at once to his wants, otherwise he passed his water into the bed. He never complained at any time of pain, either in the trunk or lower extremities.

On admission, a very abrupt and prominent curve was observed to occupy the dorsal region immediately above its centre, the spinous process of the fifth vertebra apparently forming the angle. There was no pain on pressure, nor on causing the boy to flex or extend his column. The deformity did not occupy a median position, but lay slightly to the right. Consequent upon this asymmetry and upon the existence of this curve, the pelvis had undergone a marked rotation around an antero-posterior, and to a less degree around a vertical, axis, so that the vertical axis of the sacrum, if prolonged, passed through the lower angle of the right scapula, and the crest of the right ilium was an inch below the horizontal level of the left. He is unable to make any voluntary movement of his legs, but alters their position with his hand. They feel cold to the touch, and, though well covered in blankets the whole of the lower portion of his body seems of an abnormally low temperature. His legs can be flexed, and when in that position they can be moved through a considerable angle without any very marked resistance being offered to the movement. On gradually extending them, at a certain degree of extension the muscles of the whole leg become suddenly violently contracted, and the leg and foot are

retained in a condition of extreme overtension. While this spasm is taking place, the patient experiences no discomfort, and the abdominal and intercostal muscles below the level of the deformity display no spasmodic movement. The abdominal and intercostal muscles act normally in respiration. I may say that the father states that the spasmodic contraction of the muscles of the leg while in the extended position is not so violent as it was two months ago. It is not possible to obtain any plantar reflex. The knee-reflex is exaggerated. There is no ankle-clonus. Though the patient can distinguish the point touched, it is apparent that he feels much less distinctly than normal, and that the sense of pain is diminished in the same proportion as that of touch. It is also apparent that this diminution of sensation is more marked in the feet than it is in the thighs and lower part of the trunk. Some resistance is offered to abduction as well as to adduction of the legs. The patient does not appear to experience any pain or discomfort while the legs are being moved, or while they are in a position of rigid extension.

The child coughs frequently. He does not fill the apices of his lungs during respiration, which is feeble, and there are a few *râles* to be heard in both bases. He is in the habit of wetting the bed; but whether this resulted from want of attention or not it was difficult to determine. The urine was acid, sp. gr. 1025, no deposit, no albumen, and no blood. The boy was kept quiet in bed for several days, and his temperature, which ran up for several days for a couple of degrees in the evening, varied but very slightly subsequently. His parents were very poor, and there appeared to be no sign of any improvement in the condition of his lower extremities, but, on the contrary, their state was becoming progressively worse. His family history was extremely bad; besides that, he was physically very feeble indeed. Considering these circumstances, I concluded that the only hope of preserving the boy's life was by putting him on his legs and placing him under more favourable circumstances, both as regarded hygiene and diet, than it was possible to do while he was in this helpless condition. I therefore determined to expose the child to an operation which, though comparatively trifling to a more vigorous subject, would be accompanied by some risk to one in his very enfeebled condition. The parents willingly accepted the risk.

Consequently, on February 15th, the operation was performed, every care being taken that the child was kept warm during it. The soft parts were rapidly removed from the spinous processes, laminæ, and inner portions of the transverse processes, and the spinous processes and laminæ of the fourth, fifth, and sixth dorsal vertebræ were removed with bone forceps. The spinal canal being opened, the dura mater was exposed, there being no granulation or cicatricial tissue obvious, nor was there any of the fat which normally exists in this position. The dura-matral sheath when first exposed was flaccid, and the cord felt thin and soft; at first it did not pulsate, but after a time the pulsations, though slight, were obvious. The body of the fifth dorsal vertebra appeared to be abnormally near the laminæ of the adjoining vertebræ, and it appeared to me that the cord had been forcibly compressed between these bony points. As there was nothing else to be felt there was no indication to proceed further. The wound was then closed, a small drainage-tube being fastened into the lower end of the wound, and a dressing of alembroth wool applied. The dressing was changed on the following day, when the drainage-tube was removed. The wound did not require another change of dressing.

On the evening of the 15th his temperature rose to 101.2°, and his pulse was 130, but on the following morning the temperature was normal. To prevent the dressings being wetted his urine was drawn off at intervals during the day. He knows when his bladder is full, but he cannot void it as he did before the operation. On February 18th he acquired control over his bladder, and it became no longer necessary to continue the use of the soft rubber catheter, which he objected to very much. On the second day after the operation the spasmodic contraction of the muscles of the leg in the extended posture had disappeared. Besides that, they felt warmer to the touch, as did the whole of his trunk.

On February 19th, four days after the operation, he was able to move his legs slightly in response to slight stimuli. He does not seem to be aware of the extent to which he moves. It is also obvious that he feels very much more distinctly than he did before the operation. For instance, it was possible to tickle the soles of his feet very vigorously without causing him the slightest discomfort, though he was conscious of the fact, but now tickling