visceral symptoms of that disease. It is, of course, degenerated secondarily to section or disease of posterior roots of that side, but the degeneration rarely extends far up, and therefore it is most likely that its fibres pass sooner or later into the posterior vesicular column or into the posterior median. The border zone then bears the same relation to the fine fibres as the posterolateral does to the large. Bechterew,²⁰ in 1885, had anticipated Lissauer somewhat by showing that this zone develops in the embryo at a time distinct from the rest of the posterior region.²¹

But it is probable that many visceral fibres, entering by the posterior roots, ascend the cord without communicating with the grey matter at all. The path taken by them is the posterior median column. In man this column is crowded with fine fibres. In the monkey, though the whole of the posterior median column is packed closely with them, there is a small part of the column set aside apparently for the special path of the fine fibres. This appears as a small wedge at the posterior part of the posterior septum (see Figs. 10, 11, and 12, p. v.); in it there are very few, if any, large fibres. This little tract runs right up to the medulla, and perhaps ends in the grey matter of the floor of the fourth ventricle.

We have now a complete visceral system in the cord on the same plan as the motor and sensory system of large fibres. The study of the visceral system of nerves is yet in its infancy, and its bearing upon clinical medicine is practically a new field. Is it not possible that we may by it throw some light on the pathogeny of such obscure diseases as diabetes, Addison's and Graves's disease, and perhaps even myxœdema?

My task, as far as these lectures are concerned, is ended, and yet it can scarcely be said to have begun. I have adhered strictly to my text, perhaps too strictly. I am quite aware that I have left untouched the important field of the physiology or functions of the various bundles of fibres in the cord. To deal with this subject at all satisfactorily on clinical grounds requires an experience to which I can lay no claim. To study it from the standpoint of physiological experiment would be practically impossible in the time allowed for the preparation of these lectures. It has taken fifty-three years to arrive at our present state of knowledge concerning the anatomy only of the spinal cord, from the point of view of degenerations; and I have spoken to little purpose if I have not made it evident that there is yet much to learn. If this be true of the anatomy, it is still more so of the physiology of the cord. If our art be so long that one life is too short to compass it, how much longer, then, must it be before we can pretend to understand the secrets locked up in a structure, the highest and most subtle result of evolution that Nature can show?

ERASMUS WILSON LECTURES

ON THE PATHOLOGY OF

GLAUCOMA.

Delivered at the Royal College of Surgeons of England, March, 1889

By PRIESTLEY SMITH, M.R.C.S.,

Ophthalmic Surgeon to the Queen's Hospital, Birmingham.

LECTURE III.

WE call a glaucoma primary when we are unable to point to a previous disease of the eye as its cause. The obscurity of its origin distinguishes the primary from the various forms of secondary glaucoma which we have already considered. It is important to observe, however, that, apart from their initial causes, primary and secondary glaucoma are, to a large extent, one and the same morbid process. Pressure, the essential factor, is common to both. The other leading symptoms of primary glaucoma—the progressive loss of vision, the contraction of the field, the cupping of the disc and, in the more acute forms, the pain, the injection, and the rapid abolition of sight—these are all to be found in one or other form of secondary glaucoma also. They are absent only in so far as they are hidden, or modified, or forestalled by the pre-existing disease of the eye. The special interest which belongs to the pathology of primary glaucoma lies at present, I think, not so much in the evil

consequences of high pressure—these are now fairly well understood—as in the mysterious nature of the causes which can give rise to a formidable disease to-morrow, in an eye which looks perfectly healthy to-day.

Clinical observation alone, however acute, fails to explain the malady. I will, therefore, in the first place, ask your attention to certain conditions which are found in eyes which have been blinded by primary glaucoma, and excised. Many minute descriptions of such specimens have been published, notably by Brailey in this country, by Knies, Weber, and Birnbacher and Czermak in Germany. I have myself examined about twenty-five specimens of primary glaucoma, and have placed many of them here for your inspection.

One word as to the method of examination. My own specimens were hardened gradually in Müller's fluid, and divided in the frozen state. I believe that it is not possible to open an eye in any other way without disturbing the relations of its internal parts; and even this method does not always enable us to see the parts exactly in the positions which they have occupied in the living eye. If the lens is elastic, it will become more globular when the tension of the suspensory ligament is relaxed; if it have been pushed forward by an excess of pressure in the vitreous chamber, it will tend to return to its normal position when the pressure falls after the removal of the eye. It is important to bear these points in mind in considering the changes which I have now to describe.

The filtration angle is closed, or at least compressed; I have seen no case in which it was normally patent. If the glaucoma have been of short duration, the iris-base may be merely pressed against the cornea, and may separate from it in the divided eye (see Fig. 27)¹. In older cases it is firmly adherent to it, and often much

Fig. 27.—Primary glaucoma—recent. The adhesion seems to form most rapidly

Fig. 28.—Primary glaucoma of long standing.

when there is much vascular disturbance; in the very chronic non-irritative form of glaucoma it occurs late, and sometimes not at all; but even in these cases there is always, so far as my observation goes, a well-marked compression of the filtration angle. Sometimes the iris-base adheres to the processes, and is drawn backwards when they retract, so that its adhesion with the cornea is more or less torn through; or, if it is not torn through, the iris-base may be stretched in the direction of its thickness, and present a peculiar bend or notch in its anterior surface just at the limit of the adhesion (Fig. 44).

The ciliary processes are usually altered, both in size and in position. If the glaucoma have been of recent date and congestive type, they are enlarged, their apices extending forwards far beyond the customary limit (Fig. 29).² In such cases they are in close contact anteriorly with the iris, internally with the margin of the lens; or, if no longer in contact, their wedge-like shape shows that

²⁰ Bechterew, *Neurol. Centralbl.*, 1885, p. 369.

²¹ See also Sherrington, *Brain*, vol. ix, p. 342

¹ *Ophth. Rev.*, 1882, p. 80.

² *H. and Ophth. Hosp. Reports*, vol. x, p. 25.

in the living eye they have been tightly pressed between these structures. Viewing them in transverse section (compare Figs. 30 and 31), we see that they are increased in thickness also, the

Fig. 29.—Primary glaucoma—acute.

Fig. 30.—Primary glaucoma. Transverse section through ciliary processes.

Fig. 31.—Healthy eye. Transverse section through ciliary processes.

spaces between the processes being narrowed or even obliterated by the swelling of the lateral convolutions. If, on the other hand, the glaucoma have been of long standing, the processes are sometimes much shrunken and retracted, being then far removed from the base of the iris; but even in such cases the iris-base often bears the impress of their former pressure. Between extreme hypertrophy on the one hand, and extreme atrophy on the other, any degree of swelling or shrinking may be present.

The ciliary muscle participates in these changes; at first it is drawn forwards with the processes; later it retracts and atrophies.³

The lens is sometimes in close contact with the processes and iris; more often it is a little separated from them, for being pushed forwards during the glaucomatous attack, it recedes when the pressure is removed. Its relations should be studied in the frozen globe immediately after its bisection.

A careful study of these changes in specimens of recent glaucoma leaves, I think, no uncertainty as to the manner in which the filtration angle becomes closed. The iris-base is pushed forwards by the ciliary processes, as Adolph Weber asserted thirteen years ago, and the processes are usually themselves pushed forwards by an advance of the lens and zonula, in other words by an excess of pressure in the vitreous chamber.

Now this point is of crucial importance, for it involves a true or a false conception of the whole morbid process, and I am bound to remind you that some investigators have attempted to explain the closure of the filtration angle in an entirely different way. Max Knies, the first writer on the subject, declared it to be due to a circumscribed inflammation of the ligamentum pectinatum and the adjacent tissues. Birnbacher and Czernak have recently maintained the same opinion, and have even gone so far as to assert that a closure of the filtration angle by pressure from the ciliary processes is entirely out of the question.⁴ In view of the conditions which are to be seen in many of the drawings and specimens which I am able to lay before you, this assertion is quite astonishing until one examines the evidence upon which it is based. Drs. Birnbacher and Czernak examined seven eyeballs: two of these were without clinical history. In the remaining five, the duration of the glaucoma had varied from seven months to twenty years. There was no very recent or acute case. The eyes were opened without freezing, and in one case without hardening. The material and method were obviously ill adapted to determine the point in question, and I feel sure that, with more favourable opportunities, these most painstaking observers would have come to a different conclusion.

We see, then, that in primary glaucoma the ciliary processes press against the base of the iris, and thereby close the filtration angle. We have next to inquire what are the conditions which bring the processes into this abnormal and mischievous position. It will be convenient, in looking for the initial causes of the disturbance, to consider separately the conditions which predispose to glaucoma and those which actually induce it. The predisposition may properly be considered first.

(To be continued.)

DIPHThERITIC GASTRITIS OR GASTRIC DIPHThERIA.

By TALFOURD JONES, M.B.Lond., Univ. Medical Scholar, Consulting Physician to the Breconshire Infirmary; formerly Medical Registrar and Pathologist to the Hospital for Sick Children, London.

I AM desirous of contributing the notes of a very unique case of diphtheritic gastritis or gastric diphtheria which came under my care many years ago, but which have never been published. Cases of true diphtheritic gastritis are very rare; a few have been described by Sir William Jenner, Dr. Wilson Fox, and Dr. Wilks; but in many works on medicine no mention is even made of such an affection; hence I am led to think that the following case is worthy of being recorded.

F. C., aged 2 years and 10 months, a rickety child, was taken ill with diphtheria on December 1st, 1862. On the third day there was difficulty in swallowing; on the sixth she vomited several times; on the seventh difficulty in breathing set in, and she vomited frequently, bringing up each time a little blood and some dark red pieces of a membranous character; on the eighth day the breathing became worse, and her doctor recommended her to be sent to the Hospital for Sick Children, in Great Ormond Street, London. On her way to the hospital she died in her mother's arms, but was left in the hospital, and I made an examination of the body.

The following are the chief points that need to be mentioned now. The tonsils were studded with shreds of a dirty grey coloured exudation about one-twelfth of an inch in thickness. A similar exudation, but more continuous, covered the arches of the palate, the whole of the uvula, and the recess between the base of the tongue and the epiglottis. The membrane readily peeled off, and the subjacent mucous membrane was seen red and congested. There were several irregular patches of false membrane on the posterior surface of the soft palate and on the upper part of the posterior wall of the pharynx. On removing the hard

³ Brailey, R. Lond. Ophth. Hosp. Reports, vol. x, p. 89.

⁴ Von Graefe's Archiv, vol. xxxii, Part ii, p. 1.