BRITISH MEDICAL JOURNAL.

BEING THE JOURNAL OF THE BRITISH MEDICAL ASSOCIATION.

EDITED FOR THE ASSOCIATION BY ERNEST HART.

LONDON: SATURDAY, JULY 6, 1889.

NOTES OF THIRTY-ONE CASES OF SUPRAPUBIC OPERATION.

PERFORMED BY

SIR HENRY THOMPSON, F.R.C.S.,

Consulting Surgeon to University College Hospital.

(Reported by H. T. HERRING, M.R.C.S., M.B.)

HAVING been engaged in assisting Sir Henry Thompson in completing the histories of all the cases in which he has operated on the bladder for calculus and for tumour up to the end of 1888, upwards of 1,000 in number, I have with his assent made a brief outline, comprising all the operations performed above the pubes in these circumstances by themselves. I may add that at four of the latter cases I have been present.

They are thirty-one in number, and include two performed by the old method, as generally adopted before the modification suggested by Garson, and carried out so successfully by Petersen, of Kiel. This latter operation Sir Henry Thompson introduced here, and first performed in this country, being Case I., second series. There are twenty-nine cases of the new operation; eleven of these were for tumour of the bladder, and eighteen for calculus, including one remarkable case¹ associated with calculus.

The chief points attended to in this operation were, to commence with a fully, yet prudently-distended rectum, followed by a fair distension of the bladder; a sparing use of the knife, so that a ligature was very rarely necessary; the opening of the bladder by a small incision sufficient to admit the finger, but enabling further enlargement to be made by dilatation only, since after this the opening contracts, and stitching is not necessary, and moreover appears to be generally not advisable.

TWO CASES OF SUPRAPUBIC OPERATION BY THE OLD METHOD.

CASE I.—E. F., aged 13, sent by Mr. Mason, of Burton-on-Trent, for admission to University College Hospital.

1865, January.—She stated she had passed a stout hairpin into the bladder; much pain followed; attempts in the country to remove it were unsuccessful. Under chloroform examined the bladder; the pin was embedded firmly, and no traction short of violence would remove it. The bladder was opened above the pubes upon a sound and the pin withdrawn. She made an excellent recovery in about a month's time, and was on the point of leaving, when she was suddenly attacked by severe pain and rigor, and died in two days. The autopsy showed that an abscess had formed in the track of the apparently healed wound, and had burst into the peritoneal cavity.

CASE II.—Capt. H., aged 58, 1877. A large uric acid calculus removed. Mr. Teale, of Leeds, present. The patient had been the subject of complete paraplegia for years, and on this account could not be placed in the position for lateral lithotomy. All the urine had, during that period, been drawn by catheter. A difficult operation from the presence of a long false passage. He sank exhausted ten days afterwards.

THE SERIES OF SUPRAPUBIC OPERATIONS BY THE NEW METHOD. (CASES OF TUMOUR IN THE BLADDER.)

CASE I.—Miss P., aged 73, sent by Dr. Edis, of Gloucester. Vesical bleeding about a year and a half, lately considerable, very frequent micturition, suffering extremely severe.

¹ Clin. Soc. Trans., vol. xxi, p. 46.

October 17th, 1885. Digital exploration, by dilating the urethra, revealed a large papillomatous growth springing from the upper extremity of the vesical wall; hence the suprapubic operation was necessary, and a tumour about the size of an orange removed. Recovering slowly but steadily, the symptoms greatly improved. She died nine months afterwards. The operation had undoubtedly prolonged and rendered her life more easy.

CASE II.—A. S., aged 55. Micturition frequent for four or five years, now every half-hour. Blood first seen one year and nine months ago, lately increasing. Patient very feeble.

November 16th, 1885. The bladder evidently filled by the tumour, and the high operation was performed. Dr. Lloyd and others present. A quantity of papillomatous growth removed, leaving a firm hard base. The patient returned home relieved at the end of a month, but died during the following year, recurrent masses (of round-celled sarcoma), having appeared in the situation of the wound.

CASE III.—J. A., aged 45. Seen with Mr. Maurice, of Reading. Symptoms of four years' duration, consisting chiefly of blood in the urine and increased frequency of micturition.

October 8th, 1886. Digital exploration revealed a soft growth, with wide base attached to the upper part of the bladder. Evidently was only to be dealt with through a suprapubic opening. The wound healed in a month. He was then able to hold his water four hours, which was passed free from blood, and without pain. It reappeared next year, and was similarly removed by Mr. Maurice.

CASE IV.—H. T. B., aged 54. Long-standing symptoms of vesical tumour, lately severe, and recognised as a large growth. He was seen with Dr. Boyd, of Wandsworth.

December 8th, 1886. A large pedunculated papillomatous tumour by high operation, and the whole removed. He made a rapid recovery. In November, 1887, he was in good health and actively employed, and there has been no sign of the reappearance of the growth.

CASE V.—Capt. B., aged 61. In April, 1885, a considerable papillomatous growth had been removed by a perineal opening with much relief. He returned February, 1887, his symptoms having reappeared.

February 26th, 1887. The high operation was performed in presence of Dr. Wilson, of Boston, and Dr. Mudd, of St. Louis, and a large papilloma removed. His symptoms were somewhat relieved, but he never regained his strength, and he died the following June.

CASE VI.—A. W. P. K.; sent by Dr. Travers, of St. Leonards. Attacks of hæmaturia for many years. The bleeding had lately become continuous, and papilloma was suspected. The pain and frequent micturition were very severe.

March 21st, 1887. Digital exploration verified the diagnosis. The suprapubic operation was done in the usual way, and a fibriated papilloma was removed, leaving a thick and hardened base. Much fever followed, from which he rallied, but gradually grew weaker, and ultimately died on April 1st.

CASE VII.—Major B., aged 62; for eighteen months has been subject to hæmaturia, lately becoming severe.

May 6th, 1887. Removed by the suprapubic operation a small, wide-based, papillomatous tumour. Mr. Maurice, of Reading, present. Recovery delayed owing to the deposit of phosphates in the bladder.

May, 1889. During the last year and a half he has had frequent trouble from the same cause, but when these have been removed by a lithotrite he lives in tolerable health and comfort.

CASE VIII.—G. L., aged 63, seen with Dr. Pickering, of Monte Carlo. Last fifteen months marked symptoms of vesical tumour.

July 7th, 1887. Digital exploration, finding a broad-based tumour, and the high operation at once performed, removing as much as possible. He recovered slowly, and returned to the

country relieved. Three months later he gradually sank from the extension of the growth, which proved to be epithelioma.

CASE IX.—E. I., aged 57. First seen with Dr. Montgomery, of Maidenhead. Had lost much blood during the last three months. Examination of the urine showed a fragment of papilloma.

July, 1887. Removed a broad-based growth, very like in form to that in the preceding case. The deposit of phosphates retarded his recovery, but eventually the wound healed soundly. He called the following October to say that he was free from symptoms, and following his very active business pursuits as usual.

CASE X.—Col. B., aged 59. Three years ago first saw blood, which has been gradually increasing since. Considerable masses of round and spindle-celled tissue were found in the urine. He was seen with Dr. Pavy and Sir James Paget.

April 11th, 1888. The bladder was examined with the cystoscope, and a tumour seen on the posterior wall.

May 7th, 1889. The tumour was removed in the presence of Dr. Pavy, and Mr. A. Parkes, Longton, Stafford, and a large mass separated close to its base. He left for Eastbourne five weeks afterwards, the wound not yet closed.

June 28. The growth had evidently developed in the cicatrix, and ultimately in the original site, and he sank a month or two later.

CASE XI.—T. C., aged 50. Sent by Dr. Dobie, of Chester. Four years ago first saw blood in the urine. Now persistent and severe. Microscopic examination of the *débris* showed the presence of papilloma.

June 22nd, 1888. An irregular tumour removed as close to the base as possible. July 18th he left town, the wound having healed and free from all symptoms. Heard, in the following November, that a mass of growth had appeared in the cicatrix of the suprapubic wound, and he subsequently died in the Liverpool Infirmary.

CALCULOUS CASES.

CASE XII.—C. H., aged 36. Symptoms of stone for some time, with an abundant deposit of cystine in the urine.

June 2nd, 1884. A cystine calculus weighing two ounces and three quarters was removed at the operation, at which Dr. Pinter, of Buda Pesth, and others were present. He made a slow but sound recovery, and is now in excellent health. He was seen conjointly with Sir William Roberts, of Manchester.

CASE XIII.—W. R. P., aged 12. The stone was evidently so large for a lad that there could be no question of adopting the high operation, which was performed on March 24th, 1885. An oval stone weighing one and a half ounce was removed. Dr. Pondrell, of Euston Road, present. He returned home quite well in a month.

CASE XIV.—W. H., aged 62. Symptoms of calculus for fifteen years. The last seven years he had passed mostly in bed and on the sofa. A very large stone found on sounding.

April 29th, 1885. Dr. Hickman, of Dorset Square, and Professor Rockwood, of Colombo, present, a calculus of pure uric acid weighing no less than fourteen ounces was extracted without difficulty. He made rather a slow but ultimately sound recovery. In the spring of 1889, he wrote that he was enjoying excellent health and leading a busy and active life.

CASE XV.—J. G., aged 73. Severe symptoms had existed more or less for five years.

May 2nd, 1885. Lithotripsy was attempted, but the stone being large and not easily grasped, the suprapubic operation was performed in the presence of Dr. Hickman and Professor Rockwood. The stone weighed rather more than one and a half ounce. A rather slow but sound recovery. In July, 1887, he called to say that his health and strength are remarkably good for his age.

CASE XVI.—J. B., aged 70. Seen in consultation with Mr. Adams, of Aldersgate Street. A large stone found on sounding.

April, 1885. The patient, who was very feeble, was submitted to the operation as a last resource; he never regained strength, and he died on August 13th.

CASE XVII.—J. B., aged 76. First seen in December, 1870, and a small stone, the size of a nut, discovered. He refused operation.

August 9th, 1885. Being fifteen years after the last visit, the calculus was now very large, and the suprapubic operation alone afforded him any chance of safe removal. This was done in the presence of Mr. R. Jones and Mr. A. A. Henley, of Woodbridge, and a uric acid stone, weighing six ounces and a half, removed. He made a good recovery without any troublesome symptoms, and lived in comfort for a year or two afterwards.

CASE XVIII.—J. M., aged 50. Sent from Sydney by Dr. Mackenzie, with uric acid calculus.

March 5th, 1886. Removed a stone weighing two ounces and a half in the usual manner, Dr. F. Keegan, Indore, India, being present. He had completely recovered in a month, and was then in good health.

CASE XIX.—E. H., aged 65. For many years had passed all his urine by catheter, with great frequency and pain, and had had two operations for stone. The prostate was of an enormous size, and to relieve the symptoms a suprapubic opening was made on April 29th, 1886. Mr. G. R. Ord was present at the operation. The wound was fitted with a curved tube passing through a shield on the surface, and thus to an ordinary urinal attached to the leg. The patient was much relieved by the operation, and is still in good health, without pain. He continues to pass all his urine through the tube in the suprapubic opening (April, 1889).

CASE XX.—E. G., aged 63. Sent by Dr. Whitley, of Belfast.

July 7th, 1886. A rough stone, weighing one ounce and a half, composed of oxalate of lime, removed with ease. He returned home in the beginning of August well and free from all symptoms.

CASE XXI.—J. J., aged 62. Sent by Mr. Atkinson, of Bennington.

October 15th, 1886. A uric acid calculus, weighing one ounce and a quarter. Mr. Rice Ord was present at the operation. The patient made a sound recovery.

CASE XXII.—T. T., aged 68, sent by Dr. Potter, of Collumpton. For some years had been compelled to pass all his urine by catheter.

October 25th, 1886. Crushing attempted, but the stone was found to be large and hard, and was therefore removed by the high operation. It weighed two ounces and three-quarters. Mr. Rice Ord was present. At the end of a month he returned home perfectly well; continued so to present time.

CASE XXIII.—J. A. P., aged 59.

May 13th, 1887. An oval uric acid calculus, one ounce and a half in weight, removed as usual. He was sent by Dr. Ensor, of Port Elizabeth, South Africa. He made a sound and rapid recovery.

CASE XXIV.—J. B., aged 67, from Toronto, Canada. First seen with Dr. De Woolfson, of Shrewsbury, and a large stone found on sounding.

August 1st, 1887. Removed in the presence of Dr. Vansart, of Philadelphia. The patient made a rather slow but ultimately sound recovery.

CASE XXV.—Col. A. A. B., aged 73. October, 1887. The high operation was undertaken in this case, as the stone was not easily seized with the lithotrite. The patient was very stout, and the subject of an unusually large prostate. For the first few days after the operation he did well, and then gradually lost power, and died exhausted on the eleventh day.

CASE XXVI.—R. L., aged 36. May 5th, 1888. Dr. Maclaren, of Harley Street, and Dr. Rowell, of San Francisco, present. Crushing attempted, but the stone was a large, rough oxalate of lime formation, and therefore the high operation was preferred. Thirty-six hours afterwards abdominal pain was complained of, and the tube was removed with relief. The urine then continued to flow freely by the wound, but vomiting and pain became severe, and he died on the fifth day. At the necropsy, signs of cellulitis were seen about the sides of the bladder under the peritoneum, and local peritonitis around these points. There was no perforation of the cavity. It is difficult to account for the condition; there had been no check or difficulty met with in the performance of the operation. The stone, from its large size and spiked character, was an awkward one to remove, and may have damaged the soft parts. No other circumstance happened likely to have done so.

CASE XXVII.—D. W. G., aged 68. October 5th, 1888. Operated upon at Kidderminster with Dr. Stretton, for an oxalate of lime calculus weighing 1 oz. The wound healed within a month, and the patient is now in good health.

CASE XXVIII.—R. C. H., aged 51. Had been the subject of entero-vesical fistula for upwards of six years, and lately had developed marked symptoms of stone. Lithotripsy was contraindicated in consequence of the condition described, and after consultation with Mr. Erichsen and Dr. Townshend, of Cork, it was decided to perform the high operation.

October 13th, 1888. In the presence of Mr. Erichsen and Dr. Cummings, of Cork South Infirmary, this was done with difficulty, owing to the impossibility of distending the bladder with fluid.

It was only with extreme care that the contracted bladder was safely opened on the point of a staff, and the calculus extracted.

All went well in regard to the operation, but on October 16th he had a severe attack of bronchitis and asthma, to which he had been subject for years. His asthma had long prevented sleep in a recumbent position, and an upright one in bed was necessary after the operation. He died exhausted on October 24th.

CASE XXIX.—C. R., aged 29. Had suffered for many years with symptoms of stone. He had been sounded frequently without result.

November 14th, 1888. Examined under ether, and a large and very flat oval calculus detected. The following day the suprapubic operation was performed. The calculus was composed of uric acid, and weighed one ounce and a third. His convalescence was somewhat prolonged, but ultimately he returned home perfectly well.

It will thus be seen that among the eleven cases of tumour only one was fatal, and that from pyemia. Among the eighteen calculous cases there were three deaths, if the fatal result in Case XXVIII is not included, in which death was clearly due to the severe attack of bronchitis and asthma, to which the patient had long been subject. The whole of these cases represent the most advanced and the most complicated examples of disease, and were on that account not amenable to lithotripsy, or to a perineal section of any kind.

NOTES ON ACQUIRED LEPROSY AS OBSERVED IN ENGLAND.

By JONATHAN HUTCHINSON, F.R.C.S., LL.D., F.R.S.,

Consulting Surgeon to the Hospital for Diseases of the Skin, to the Royal London Ophthalmic Hospital, and to the London Hospital.

II.

THE case with which I concluded my paper last week was of great importance as proving that the leprosy process (that is, the life and multiplication of the bacillus) may come to an end if the patient is removed to a place where the disease is not endemic. I have another yet to relate which is quite as definite, and I have, I believe, several patients under observation in whom the process of cure in England is slowly in progress.

CASE II.—The case to which I allude is that of a Methodist clergyman, of high endowments, who was born in Canada, and who, in 1850, at the age of 24, went on missionary service to reside in the Bermudas. Five years after going to the Bermudas he married, and, being then in excellent health, continued to reside there. In 1857, owing to some nervous failure, he became unable to write with his right hand, and in 1859 an eruption appeared. In 1863 he saw Dr. Brown-Séquard, who said that it was a confirmed case of leprosy with paralysis. During the next ten years he continued to suffer from the disease in an aggravated form. On one occasion, after an injury, he had an anæsthetic toe amputated. In 1875 his eyes began to suffer, and during the next five years he became quite blind after repeated attacks of inflammation. Several iridectomies were, I believe, done by a most skilful ophthalmic surgeon (Dr. Bullar, of Montreal), in the hope of arresting the disease, but without any permanent result. There are many interesting details in this case with which I forbear to trouble the reader. It is sufficient for my present purpose to say that the Rev. Mr. D. came under my observation about twenty-five years after the beginning of his leprosy, and that he was then, at the age of 55, an example of recovery. All leprous processes had come to an end. There had been no fresh manifestations for some years. He still enjoyed the use of a powerful intellect, but he was blind, much disfigured, and crippled in both hands and feet.

The case proves (1) that one not inheriting any taint may acquire leprosy by residing in a locality where it is endemic; (2) that such acquisition may be effected wholly without exposure to hardship or poverty; (3) that the malady may come to an end in the course of years during residence in a healthy district.

I do not see that it is possible to suggest more than two ways in which it is probable that the disease was acquired. Clearly Mr. D. in some way took in, while residing in the Bermudas, the specific germs of a specific malady. He must have taken them either by direct contagion, or he must have received them in food. A similar proposition may, I think, be made in reference to most of the cases with which I am at present dealing.

CASE III.—My third case is that of a young English lady who went to reside in Burmah in connection with a mission school. She went out at 19 and returned at 30 with her features so disfigured by "tubercular" leprosy that she looked like a woman of 60. She was for some months under my care in the London Hospital. It is now eight years since I saw her, and I do not know anything as to her present state.

CASE IV.—A woman, Mrs. C., was under the care of my colleague, Mr. Waren Tay, and presented a marked example of leprosy. Her age was 31 when I saw her. She had been taken to India at the age of 5, and had lived there twenty-two years. When she returned to England she was, as far as she knew, in sound health, and it was not till two years later that the signs of leprosy became definite. The first manifestation was the appearance of brown patches about the neck and shoulders, and when I saw her these brown areas covered almost the entire surface. I have not kept any record as to the precise part of India in which this woman had resided, but she told me that it was inland, and that there was abundance of fresh-water fish. She herself did not like the fish, but she believed that she had nevertheless partaken of it on an average two or three times a week.

CASE V.—A gentleman born in England, of English and Irish descent, named L. G. H., consulted me in 1880 for leprosy. He went to live at Bombay in 1856, and in 1875 his first patch was observed. It was a discoloration on his right leg. It was not until eighteen months later that his face was affected. The disease was in its seventh year when I saw him, and was well characterised. He said that in India he had eaten fish and prawns occasionally.

CASE VI.—A gentleman of Scotch descent consulted me on account of well-marked leprosy. He had been engaged for some years on the newspaper press of Calcutta, and there his disease had commenced. This patient had never been subjected to any kind of hardship, and he had not left England until he was 20 years of age.

CASE VII.—Colonel H. P. C. was born in England, of English parents. He went to India in 1849, and returned in 1869. In 1870 he consulted a London surgeon for an eruption which was diagnosed as syphilis, but in respect to which he denied that he had had any chancre, or been exposed to any risk of contagion. He was married, and had healthy children. In 1874 he consulted me, and he was then the subject of leprosy in an advanced form, his skin being almost universally discoloured, and of a brown tint. The lobes of his ears were thickened and dusky, and the soles of his feet painful. I saw Colonel C. several times during the next year, but do not know anything as to his subsequent history. He told me that whilst living in Burmah he had eaten fish moderately, and that he had been particularly fond of a condiment prepared from prawns, which he had special facilities for obtaining. This gentleman gave me interesting information as to the prevalence of leprosy in Burmah, with the facts concerning which he was officially well acquainted. He said that it was very common, and especially so among the coast population, and that the Government was at that time engaged in making extensive provision for the care of lepers. He drew my attention to a passage which had been recently published by Mr. Francis Day, and expressed his unbounded surprise that anyone who had travelled in Burmah could have written it. The passage to which I refer is as follows: "If, however, we look a little further to the east, we find the Burmese race, who are most assuredly a fish-consuming one, and who like their odoriferous nga-pee far better than fresh fish. Yet in my recent travels through Burmah I never saw an instance of leprosy amongst the indigenous population; neither did I see any cases of that congenital malformation of the fingers and toes so common among the natives of India born with a leprous taint. In fact, it has been frequently observed that this disease does not exist amongst the Burmese, and the same remark applies to the Andamanese, whilst it is as common in India amongst the inland classes who do not eat fish, as it is amongst those who consume it." I quote this passage as an example of the very misleading information

"On Fish as Food." By Surgeon Francis Day, F.L.S. and F.Z.S. *Indian Medical Gazette*, February 1st, 1871.