ing up the poisoning for some time. We were unsuccessful in the attempt, as the symptoms always passed off when the poison was eliminated, but still the conditions in our experiment were very different from those which would occur from the prolonged use of analgesics by a patient, inasmuch as the treatment of the latter might continue for weeks, months, or years, whereas in our experiment time could be counted by days. A circumstance that renders circumspection in the use of analgesics all the more necessary is that various nervous disorders in which the symptoms appear to depend either upon peripheral neuritis or an affection of the spinal cord have lately been described as arising from the use of some of the new explosives, whose chemical constitution approximates somewhat to that of the analgesics. Another action of analgesics which renders caution desirable in their use is their effect upon the blood. Thus exalgine may render the blood incapable of performing its respiratory functions, and thus induce marked lividity, with tendency to collapse, and thus endanger life, while others may destroy the blood corpuscles and produce marked anaemia.

Possible Relation of Leucomaines or Ptomaines to Spinal Disease.—In relation to this, another question suggests itself: whether diseases of the spinal cord may not owe their origin, at least in part, to poisons of the aromatic series, generated either in the intestinal canal or in the tissues of the cord itself.

Use of Suspension.—The remarkable benefit obtained from suspension in cases of locomotor ataxy is one of the most striking discoveries that have been made in therapeutics of recent years. It is very difficult to explain, but I should be inclined to think that suspension merely acted upon the cord in the same way that massage does upon the muscles, by removing the lymph and with it the products of nerve waste; while, at the same time, it may increase the processes of oxidation and repair by promoting a free circulation of the blood.

SUMMARY OF THREE LECTURES

ON

SOME AFFECTIONS OF THE GENITO-URINARY ORGANS.

Delivered at the Royal College of Surgeons of England.

By MATTHEW BERKELEY HILL, F.R.C.S.,

Senior Surgeon to University College Hospital.

LECTURE III. PROSTATITIS.

Symptoms.—In inflammation of the substance of the prostate besides the symptoms of granular prostatitis, there are aching or stabbing pains in the anus, sacrum, or perineum; pain at the suprapubic region is a common sign. There are also radiating lumbar or femoral pains after exercise or long journeys, general languor, *malaise*, or depressed spirits. Increased frequency of micturition is often absent; when present, it is felt by day, not by night. The flow of urine is altered by delay at the beginning, which may amount to retention. Dull pain after micturition, sometimes spasm at the end of the flow, with a sense of more to come are complained of. In cases of long standing, micturition during sleep is a symptom,—constipation is almost invariable. When the congestion is great, there is pain during defaecation and between the acts a sense of weight at the anus or in the perineum. Pain in the testes when it occurs is simply a neuralgia. There is constantly hypersecretion of mucus.

In some patients, the organ recovers its natural condition in a few weeks; but usually the disease drags on a course more or less wearisome; thus the termination is very uncertain. Relapses are almost sure to follow if the patient exceed in his diet or *régime*. In relapses small abscesses often collect in obstructed ducts, which usually empty themselves through a duct before accumulating much. Sometimes the inflammation spreads to the parenchyma, and the prostate then becomes unyielding to the touch, hard, and sensitive. The enlargement of the organ may last long. The increase in size may be symmetrical, or only affect one part, giving the sensation of irregularity or lobular form when felt by the finger. In course of time, the nodules may disappear, and the organ regain its natural size, or even shrink below it, and feel quite firm and smooth.

Treatment of Chronic Prostatitis: General.—The treatment of chronic prostatitis is very tedious. Of the many remedies in use, but few are trustworthy. General treatment only assists local remedies. In the first place, the dyspepsia must be cured. The tonics most beneficial are iron in non-astringent forms; nuxvomica or strychnine are usefully added to the iron. Belladonna also may be added to the tonics if micturition during sleep be a symptom. Ergotine is especially useful when the organ is large, and aching is caused by walking, or standing, or railway journeys.

Local Treatment.—When the organ is enlarged, soft, and not tender, a very good remedy is the cold sitz bath, taken at first for one or two minutes, once or twice daily, at a temperature of 50° F., gradually prolonged to ten minutes, and lowered to 35°. The cold douche on the perineum or anus is also useful. Cold enemata are more generally beneficial than baths, two to four ounces of cold water thrown into the rectum daily, and retained there; beginning at 45° F., and gradually lowered to 35°. Among recognised remedies is blistering the perineum, but the lecturer has no faith in its efficacy. Caustic injections are valuable if used after other means. The solution of 20 to 30 grains of nitrate of silver to the ounce, and ten minims injected at a time, is a good preparation. This has to be repeated several times.

Prostatitis from Masturbation or Excessive Venery.—Masturbation is a frequent provocator of prostatitis, sometimes in the acute form, more often in the subacute or chronic forms. In the mildest form of prostatitis through masturbation, the prostate bleeds easily, the urine is often tinged with blood, smoky, or even coffee-coloured. If felt per rectum, the organ is found little changed from the natural condition. General *malaise*, inappetency, and slight elevation of temperature usually accompany the bleeding. Bleeding from the congested prostate may be copious in young men when it follows prolonged venereal excitement and excessive copulation. Commonly very copious when thus originated, it may be profuse for an hour or two, and exhausting.

Treatment.—In these affections the liquid extract of salix nigra, a drachm three times daily, has often a marked effect in checking involuntary emissions, and thereby preventing the irritation and exhaustion and neuralgia which some persons feel after these attacks.

Sequels of Prostatitis.—Persons who have suffered from a congested prostate are liable to too speedy ejaculation during copulation. This may be so rapid, and erection so fugitive, that effective coitus is impossible. For such cases the liquid extract of damiana has sometimes a powerful effect. But cure is almost hopeless if continence is neglected. Some persons give way to the sexual appetite without limit. Then follows a condition of great helplessness. The intense neuralgia of the lower extremities which follows the shortest walk renders exercise impossible. Pain is rarely absent, even when at rest; it extends from the sacrum along the spinal column to the occiput. The digestion is deteriorated. Topical treatment, after the mucous catarrh and chronic inflammation of the prostate is quelled, must be laid aside. The best remedy is a life of quiet, absence of exertion of any kind; a long sea-voyage is one of the best remedies. In rare cases chronic parenchymatous prostatitis excites temporary mania, which has erotic irritation for an early or leading symptom. It disappears if the patient recovers from the physical affection.

TUBERCULOUS DISEASE OF THE PROSTATE.

Etiology.—This frequently follows gonorrhoea. In the early stages the symptoms, being those of catarrh, escape notice, and the tubercle in the prostate cannot be detected by the finger until it has become considerably developed.

Early Symptoms.—This catarrhal condition of the prostate, with slight irritation of the bladder, lasts an indefinite time; eventually attention is directed to the prostate by the occurrence of other symptoms, such as frequent seminal emissions or by weight in the perineum and pain in the sacrum. In other cases, the patient notices increased frequency of call to micturate, but a slowness in starting the stream of urine; then he feels a burning or cutting pain during the flow.

Later Symptoms.—As soon as the shape and consistence of the prostate are altered, the finger in the rectum finds the prostate irregular, perhaps larger than natural, tender, particularly at the posterior part near the trigone. The disposition of these irregular thickenings is variable. When cystitis is developed, copious viscid pus mixed with shreds and blood clot form the urinary sediment, and the urine contains albumen in considerable quantity. The

further progress of the case is accompanied by gradual participation of the ureters and kidneys in the septic inflammation, and the patient is gradually exhausted. Acute general tuberculosis may terminate his sufferings.

Age.—The ages of the patients usually range from 19 to 40 years.

Use of the Cystoscope.—The lecturer had inspected two cases of advanced disease by the cystoscope, and the surface was irregular, with pits at the uvula vesicæ or neighbouring part of the floor of the bladder, partly filled with shreds of tissue, muco-pus, and blood clot, though in those cases cystitis, ulceration, burrowing abscess in the floor and walls of the bladder are usually super-added before death.

Differential Diagnosis of Tubercle in the Prostate.—In early cases the presence of tubercle elsewhere may be wanting, as the prostate is often affected before other organs. The disease most commonly suspected when tubercle is irritating the prostate is stone in the bladder. The urine is at first only moderately turbid in both. In prostatic tubercle there is often a gleet, or history of such having continued since an attack of gonorrhœa; a little shreddy mucus may be washed out with the first portion of urine voided, the remainder being only slightly turbid. In stone, on the other hand, a drop of blood often comes at the end of the stream instead of pus at the beginning. When blood comes from the prostate it is usually washed out first as coagula mixed with curdy pus. If the patient be sounded and stone be there, it can be generally detected. Sounding does not always discover any alteration in the shape of the prostate or trigone unless the tuberculous disease be far advanced. The distinctions of vesical tumour from prostatic tubercle are so great that confusion is not likely to occur. Tuberculous pyelo-nephritis creates symptoms which often suggest disease of the neck of the bladder. In these cases the absence of change in the condition of the prostate and the presence of tenderness in the renal regions indicate the real seat of disease.

Treatment of Tuberculous Prostatitis.—In the early stages the catarrh must be cured, and the general treatment for tubercle applied. In the later stages the ragged cavities must be carefully washed out. A good antiseptic solution is two grains to the ounce of sulphate of quinine, two ounces being injected and left in the bladder after the pus and urine are well cleared out by repeated small injections of boric acid solution. Still more antiseptic is an emulsion of iodoform. In cases of chronic cystitis, if the bladder is well washed, and a couple of drachms of this emulsion injected, the most foetid ammoniacal urine is replaced by acid urine, fœtor disappears, and the pus rapidly diminishes. In cases of cystitis caused by neglect of prostatic retention, the urine in ten days becomes quite free from deposit, and even when calculus or tumour or malignant ulceration co-exist, the improvement is enormous. The lecturer exhibited two diagrams copied from drawings made of the floor of the bladder viewed by the cystoscope, which showed an early tuberculous ulcer of the mucous membrane before and after healing. This was cured by washing out the bladder with an injection of iodoform emulsion, so that in a fortnight the patient, a man with advancing pulmonary tubercle, was cured of his bladder trouble. When the pain of washing the bladder is severe, cocaine previously injected often renders the operation bearable. When things are too far gone for this, ether should be given while the bladder is thoroughly washed with a large silver catheter and Clover's extracting bottle, then a drachm of iodoform emulsion is injected and left in. Subsequent washings may then be carried out with very little suffering. The formula for the emulsion is: iodoform, 2 parts; mucilage, 4 parts; glycerine, 2 parts; water, 20 parts.

Perineal Section.—In former years the lecturer had found it necessary to do perineal section in order to obtain constant drainage when the urine could not be restored to a healthy state by other means. But the use of iodoform emulsion may prevent this necessity when perineal drainage is carried out. The daily washing is still as needful as when the urine is voided by the urethra. The effect of this thorough drainage is so great that patients feel strong and well while it is continued, and a permanent cure by this method may be attainable. In a patient under the lecturer's care in University College Hospital, the bladder trouble was kept in complete abeyance, if not quite cured, and the patient grew stout and strong, until he sank under rapid tuberculous meningitis.

Demonstration of the Endoscope and Cystoscope.—At the conclu-

sion of the lecture the endoscope and cystoscope were shown. Grünfeld's endoscope, illuminated with light from a kerosene lamp, was inserted into the urethra of an ox. Leiter's endoscope, illuminated with electric light, was used to explore the urethra in a patient with chronic urethritis. There were several cystoscopes: Two of Dr. Nitze's, and two of Leiter's (a modification of Nitze's). Nitze's is of better shape than Leiter's, but Leiter's has the most powerful illumination. One of Nitze's has been reconstructed by Mayer and Meltzer, so as to permit of a free flow of water through the bladder, by which blood and pus are cleared away and the contents of the bladder kept translucent during introspection. Introspection was made in "phantom" or dummy bladders, on the interior of which mock papillæ and orifices of ureters had been painted. These were borrowed from Mr. K. Schall, of Wigmore Street, Leiter's agent in England, who also superintended the batteries employed for illuminating the electric lights. Just too late for this exhibition, Mr. Schall received an irrigating cystoscope from Leiter, which is of extremely ingenious construction, and promises to be efficacious.

SOME CASES IN PRACTICE BEARING UPON THE FUNCTION OF THE PROSTATE.

By REGINALD HARRISON, F.R.C.S.,

Surgeon to the Liverpool Royal Infirmary, and Lecturer on Clinical Surgery in the Victoria University.

MANY examples will occur where both to the pathologist and surgeon the observation of processes of disease in organs of the body, as well as the operative procedures necessary for their relief, have proved of service in assisting to determine the purpose and function of the part in question. In my Lettsomian lectures (1888), I laid stress on the muscular action of the prostate relative to micturition, independently of its glandular association with the generative act, whatever the latter may be. In this muscular exercise I included (1) a sphincter action, such as we know of in connection with all outlets from the body, and (2) the provision of a muscular basis for supporting the bladder and its contents. The latter purpose seems to have been insufficiently recognised, not only as a normal and well defined function of the prostate, but relatively to that form of hypertrophy of the part which constitutes disease.

In the first place, it will be noted that all deep sections of the prostate radiating from the urethra outwards, deprive it for a time of its sphincter power, and thus render it incapable of controlling the retention of the urine. Of wounds of the prostate which may cause temporary or permanent incontinence, we have instances in the deep incision of lateral lithotomy. In the ordinary median operation, the prostate is not touched with the knife, hence, where the stone is not large and the amount of dilatation or prostatic tearing is not great, the patient may never entirely lose control over the urine. This is one of the objections to the median operation, where for some reason or other it is necessary to subject the bladder to a process of involuntary drainage. In the few instances which have come under my notice where, in the male, lateral lithotomy has been followed by permanent incontinence of urine, I have no doubt that the prostatic ring has been completely severed either by a too deep incision or by the tearing of the parts in the extraction of the stone. Under these circumstances it is hardly fair, on reflection, to urge such an occurrence as an objection against the operation of lateral lithotomy.

The power of the prostate in retaining urine is further evidenced in those cases where the urethra has either been extensively or completely ruptured in its membranous portion, just in front of the apex of the prostate. Where such an accident happens, retention of urine immediately occurs, and continues until the degree of tension on the outlet of the bladder becomes such as to paralyse it; in this way time is given for Nature to repair in, and for art to interfere. In numerous cases of this kind where I have opened the perineum in the median line within twenty-four hours of the accident, for the purpose of providing an escape for the urine, it has been noted that though the bladder may have been distended, there was no evidence that extravasation of urine had occurred. The contracted condition of the prostate as determined by the finger has, as I have frequently pointed out, been quite sufficient to explain this. In the case of the last patient I operated upon, that of a young man aged 22, who was