first to last would be startling could they be accurately set down.

The vast extension of the habit of novel reading amongst us is also demonstrated by an observation of journalistic literature. Newspapers that at one time confined themselves to news, political commentaries, and criticism, now form syndicates to supply their readers with a modicum of fiction, and journals entirely devoted to fiction have an enormous circulation. One such journal, a penny one, and by no means one of the best known, had to my knowledge a few years ago a circulation of a quarter of a million a week, and no doubt therefore reached millions of readers, for such journals pass from hand to hand; and journals which combine fiction with fashion, which is imagination applied to personal decoration, are sold, I am informed by a high authority, in hundreds of thousands in this country weekly, and are exported in tens of thousands weekly to our colonies, being purchased both here and at the Antipodes chiefly by servant maids and shop girls. And that the demand for cheap fiction will go on growing can scarcely be doubted, for the monotony of life which the division of labour has so greatly aggravated, the aspirations which even a humble education serves to implant, and the increased mental friction which arises from the aggregation of masses of our people in large towns, all tend to whet the appetite for an imaginative diet. To us as medical men it is interesting to remark that this appetite is most urgent in spring, when nervous erethism exists, and "a young man's fancy lightly turns to thoughts of love," and is least pressing in autumn, when the nervous system is comparatively quiescent. In the Birmingham lending libraries the issue of novels reaches its maximum in March—32,796 were issued in that month last year—and touches its minimum in August—27,140 were issued in that month in 1888—a difference of 5,656 in favour of March.

And now, gentlemen, having trespassed as far as I dare on your patience, I must close without having even mentioned several topics in connection with the hygienic use of the imagination which I had proposed to bring under your notice in some detail. As I voyaged forth on this address I trusted to be able to submit to you some observations on sex and age in relation to imagination, on types of imagination in individuals and races, on the conditions of imaginative production, on the operation of imagination, causative and corrective, in certain special nervous and mental diseases, and on the education of imagination; and now I find myself stranded on the sands of the hour-glass, when I have scarcely got beyond my introduction. Let me console myself by thinking that I shall on some future occasion—not, of course, in my present gratifying position, for that comes but once in a lifetime, but in some other place—disburthen myself of what I have to say on these matters. And let me console myself above all by believing that you will leniently judge my introduction to have been not altogether unworthy of your attention. Matthew Arnold said that two-thirds of life are conduct, and it may be truly alleged that three-fourths of thought are imagination. It is not merely the child who builds castles in the air. Each of us—the most matter of fact as well as the most romantic—has "cloud-capped palaces and gorgeous towers" known to himself alone, to which in hours of solitude and reverie or sadness and chagrin he retires with ever-renewed wonder and refreshment. And the great faculty that out of thin air constructs these cities of refuge for us, and enables us in a sense to shape our own environment, that is the hand-maid of science, the giver of insight, the harmoniser of discords, must always receive primary consideration in a study of mental hygiene.

BRADFORD.—*Heavy Mortality from Phthisis: Cases of Lead-poisoning.* The death-rate for the quarter ended March 31st, 1889, was 19 per 1,000 per annum, an increase on that of the quarter immediately preceding, and of the corresponding quarter of last year. The increase was of a general character, and does not call for special comment. From infectious diseases 101 deaths were recorded, made up as follows: small-pox, 1; measles, 14; scarlet fever, 17; diphtheria, 2; whooping-cough, 43; enteric fever, 2; diarrhoea, 20; and two of doubtful character. Phthisis caused 102 deaths, or one more than all the zymotic diseases put together. The question of lead-poisoning by water from the public supply attracted a good deal of attention during the early part of the quarter, and a few cases occurred in the borough, though none terminated fatally.

AN ADDRESS

DELIVERED AT THE OPENING OF

THE SECTION OF SURGERY.

At the Annual Meeting of the British Medical Association, held in Leeds, August, 1889.

By T. R. JESSOP, F.R.C.S.,

Senior Surgeon to the Leeds General Infirmary.

ON THE TREATMENT OF CANCER OF THE RECTUM.

GENTLEMEN,—You are probably aware that the Council of our Association have issued a request calling upon the chairmen of the various Sections to refrain from inflicting upon their audiences anything in the shape of a presidential address. In the judgment of the Council the Sections will best fulfil their purpose if the time allotted to them—admittedly too short for the adequate discussion of more than a small proportion of the various subjects of interest offered for consideration—is devoted without deduction to the reading and criticising of the papers forming the agenda, which have been prepared by their authors at the cost of so much time and labour.

In this decision, as it affects us in the Surgical Section, I fully concur, knowing well how much more profitably your time will be spent in the thoughtful consideration of the important surgical questions to be raised, than in listening to mere generalities such as constitute the orthodox introductory discourse. With this request before us, my official colleagues were good enough to suggest that I should undertake to open a discussion upon some subject of my own choosing; and on looking at the length of the list of eminent surgeons who have promised to follow me, it is gratifying to me to find that I have succeeded in exciting so wide an interest by selecting as the initial subject of our deliberations the treatment of cancer of the rectum.

You will have observed that, in the exercise of that discretion which is granted to the officers of every Section, we have this year—extending an experiment which has in former years been tried on a smaller scale with, as I think, encouraging success—determined to commence each day's meeting with a discussion upon a set subject, for the more interesting and instructive treatment of which we have invited contributions from selected authorities. It is hardly necessary to say that, when each list of names has been exhausted, it will be our pleasure to listen to any remarks from other gentlemen desirous of speaking upon the questions under consideration; and I hope in this way the number of speakers will be largely augmented. Our aim is not only successfully to compete with those lighter attractions provided outside the sectional work, but is directed to the more worthy object of collating and of placing upon record the deliberate observations of men whose opinions are entitled to the respectful consideration of the profession. At the close of each discussion, such other papers as are on the list will be taken *seriatim*, so long as time shall permit.

I would here crave one moment for a pleasurable and obvious duty, that, namely, of expressing my sense of the high honour conferred upon me in being asked to lead and to regulate the discussions in this Section. The honour I feel is greatly enhanced by the fact that my predecessors have been chosen from amongst the most eminent surgeons of the day, and my responsibility is made heavier by the recollection that, when the Association last met in Leeds—exactly twenty years ago—the chair I now occupy was filled by one of the ablest and most cultured of the surgeons whose names are to be found upon the Leeds Infirmary roll—the late William Hey—at whose feet in 1869 I was proud to sit in the not less useful or less honourable capacity of Secretary to the Section. Fortunately for me, your Council have surrounded me with able colleagues, to whom I shall not look in vain for help, all of whom but one—himself a welcome accession to us, prominent even among London surgeons—have been closely associated with me in the past, and will, I trust, for years to come share with me that personal friendship which proverbially belongs to the surgeons of the General Infirmary at Leeds.

Without further encroachment upon your time, I will pass on to the consideration of the subject we are here to discuss, namely, the treatment of cancer of the rectum. And here let me say that my object shall be, whilst stating my own views when amply matured, to invite the opinions of subsequent speakers upon some of the debatable or imperfectly settled questions which present themselves to the surgeon during the course of cancer as it is met with in the upper, the middle, or the anal portions of the rectum, and again as it involves a portion only or the entire circumference of the bowel.

When making selection of this subject I had in my mind nothing more than the surgical, or, as it may perhaps more appropriately be termed, the operative treatment of the disease; but I may perhaps be permitted to make a very brief allusion to the help we may hope to obtain from medicines alone. Curative drugs are no more and no less applicable to cancer in this situation than to the same disease in other parts; upon them I would that we could profitably linger. My attitude towards the question of a cure for cancer is one of hope and even of faith in the future, but of absolute distrust in the past. That the discovery of an agent capable of selecting and of destroying the cancer microbe—if cancer microbe there be—may reward the lifelong labour of some future Pasteur is reasonably conceivable; that such an agent has been found in Chian turpentine or in any other known germicide rests upon the slenderest foundation; and after repeated, patient, hope-inspired (for who could turn a deaf ear to such statements as were put forth by a justly-respected member of our own profession?), and exhaustive trial, this drug, like all its predecessors, has been relegated to its proper place upon the back shelves of the chemist's shop, to be unearthed at some future time and curiously viewed as embodying, it may be, an honest attempt to accomplish an end which shall then happily have been successfully attained.

After lamenting his powerlessness to grapple with cancer, William Hey the first prayed that the time might come when the means of cure should be vouchsafed to man; and now, nearly a century later, Hey's prayer may still form part of the surgeon's missal; for although every district has for ages possessed and still supports its cancer curer, it is none the less true that cancer never yet was cured. Much, however, may be done medicinally to alleviate the great suffering and distress attendant more especially upon the later stages of the disease. The washing out two or three times a day with simple warm water, and the introduction into the emptied bowel of a suppository containing cocaine, belladonna, morphine, and iodoform have been attended in my hands with the most comforting results. Laxatives, again, seem called for chiefly in those cases in which the upper portions of the organ are involved, whilst the anal portion is free; and on the other hand, astringent measures and such as tend to solidify the evacuations I have found more suited to the cases in which, the anus being implicated, the sphincters have ceased to act, and the feces, liquefied by admixture with sanious mucus, are more or less constantly dribbling, to the patient's unspeakable distress.

Upon these points—the value of and the best method of securing cleanliness of the parts, the use of antiseptics and of anodynes, the means whereby we may procure for our patient the nearest approach to a single daily evacuation—I shall hope to elicit valuable information.

We pass on then to the operative treatment of cancer in the rectum, and I propose briefly to consider it under two leading heads: operations, namely, for the ablation of the disease—proctectomy, as they are termed; and secondly, operations for diverting the feces from their passage through the rectum, and thus far dispensing with the rectum as an active organ, and placing it as it were outside the alimentary canal, no longer required to perform its ordinary functions, and no longer subjected to the irritating presence of effete material. The former operations must almost of necessity be confined to the lower reaches of the bowel, whilst the latter have very generally been employed for relief from the disease wherever situated.

I pass by partial removal, whether by scraping or by scissors and knife, as being but rarely advisable, and barely scientific; and I will do no more than mention proctotomy as a possible occasional resource for the relief of partial obstruction when other measures are deemed inapplicable.

First, then, with regard to proctectomy—an operation which, though sporadically performed in this country for many years, has only of late been forced into prominence by the persistence and the success of our German colleagues—I will make this observation at the outset that, based upon the local origin of cancer,

it aims at the complete eradication of the disease, and therefore, if successful as an operation, not attended by an undue mortality on the one hand, or followed by detracting disadvantages on the other, it must take its stand as an established surgical procedure. I am not at all sure, indeed, that it may not be so considered already. The reports published from time to time—some of them by gentlemen to whom we shall hope to have the privilege of listening to-day—encourage us in the belief that, with the perfecting of procedure, it only remains to determine in what cases proctectomy is or is not to be recommended. The mode of operation, its every detail, I take it, is an all-important matter here; and I shall hope to have much light thrown upon various points of difficulty and doubt; such as what is the best method of obtaining free access to the parts to be removed; of avoiding and of arresting hæmorrhage during or after the operation; of dealing with accidentally opened cavities and wounded organs, the peritoneum, the bladder, the prostate, the urethra, the vagina; of adjusting the portions of rectum unremoved; of draining; of securing an aseptic condition; and of promoting the favourable healing of the wound; of retaining an active and efficient sphincter; and, finally, of preventing cicatricial contraction at the outlet. Then, again, what are the conditions favourable for this undertaking? When is it to be recommended, and under what circumstances is the patient to be denied the relief it offers? As yet my experience is small, hardly perhaps worth relating, my cases numbering no more than seven. Of these, one died as the immediate consequence of the operation. The patient was a poor curate, dejected by misfortune and poverty. The disease was extensive, occupying four fifths of the circumference of the gut just within the anus, and reaching so high that the finger could not touch healthy mucous membrane beyond. Although nothing occurred during the operation to intensify shock, he never fairly rallied, but lingering on a few days with the aid of stimulants, he died without actually emerging from the primary collapse. The remaining six obtained a varying amount of relief. In the three most favourable the disease was situated in the posterior or lateral walls of the rectum above the sphincter, and well within reach of the finger; and these, at the time of writing, are all alive: one apparently well at the end of twenty-one months, and showing no signs of return or of much inconvenience; the second with an equally favourable record at the end of seventeen months; whilst the third, operated on nine months ago, has not been able to leave his room owing to a speedy return of the disease, which has now invaded the pelvic tissues to such an extent as to threaten an early fatal termination. A fifth case, equally favourable with the former three, was that of a lady, upon the anterior surface of whose rectum was a deep hard-based carcinomatous ulcer, of the size of a florin. The vaginal mucous membrane could be moved upon the growth, so that I was able to dissect away the infiltrated structures and leave the vaginal canal intact. An untoward circumstance marred and interrupted the course of recovery I naturally looked for. The wound was dressed with undiluted salufer, and to the escharotic action of this powerful and useful antiseptic I attribute the subsequent perforation of the recto-vaginal wall, by the occurrence of which this patient's convalescence and ultimate good recovery was delayed. The present condition of this lady—thirteen months after the operation—is fairly satisfactory; for, whilst there is no return of the disease, the small vaginal opening, now completely cicatrised, in no way interferes with the action of the sphincter, and permits only of an escape of feces when from any cause the motion is not solid.

The two remaining operations were undertaken for the removal of disease situated at a higher level. In each there was a hard nodular ring completely encircling the bowel, through which the finger-tip could be made to pass, whilst the anal portion of the rectum to the extent of from two to three inches was unaffected. In both, without much difficulty, the whole of the new growth was removed. The first was an elderly male with a capacious pelvis, whose tissues were spare and relaxed. After removal of the ring I succeeded in approximating and suturing the divided edges of the mucous membrane. In the second, a stout middle-aged woman, Douglas's peritoneal pouch was extensively wounded, and through the opening the uterus and the small intestines were readily felt. Here I determined not to attempt any suturing, but to trust to unaided adaptation, free drainage being secured by the introduction of a large celluloid tube. Both cases recovered so as to leave for home, and both remained well when last heard of, five months in the one case and twenty-six weeks in the other having elapsed since the operation.

Of the seven cases, one, I have already said, died within a few days; in another the disease rapidly returned; and in a third, except that a small recto-vaginal opening permits of the occasional involuntary passage of flatus and fæces, a cure, at least of temporary duration, has been brought about; whilst in the remaining four excellent immediate results have been obtained, and the conditions in each as regards defæcation and local suffering are almost all that can be desired, and are incomparably better than the most exceptional colotomy can effect. In addition to these advantages is the hope of a more or less prolonged immunity from the disease. Results even such as these encourage us to hope that, with increased experience and a more accurate appreciation of the causes of danger and of failure, we shall find in proctectomy a valuable means of prolonging life and of mitigating suffering applicable to a considerable number of cases for which other measures offer at the best but imperfect palliation. We shall look to this discussion for the fruits of collective experience, and especially for guidance in the methods of procedure and in the perfecting of the details of the operation. I have not myself attempted hitherto the extirpation of cancer when beyond the reach of the finger as felt in an ordinary examination, and I shall, therefore, feel much interested in the experience of those who have removed neoplasms from the higher portions of the rectum. Neither have I in the male advised removal where the disease has been seated on the anterior wall of the rectum, as I have had reason to believe the operation might involve injury to the urethra, prostate, or bladder. The opening of the peritoneum in proctectomy, though by no means free from risk, is not, as we have seen, of necessity followed by serious consequences. On the other hand, a communication with the urethra or bladder could hardly fail to result in such inconveniences as would more than neutralise any benefit derived from the extirpation of the disease.

Before leaving this part of my subject, let me briefly allude to one or two points of interest and importance in the operation itself. With the patient placed in the lithotomy position, access to the disease may be obtained in one of two ways—either by widely expanding the sphincter by means of a modified Weiss's three-branched dilator, or by cutting from the rectum deeply down to the coccyx. The former plan may suffice in thin, wasted subjects, whilst in stout persons a full view of the interior of the rectum can only be brought about by a deep and extensive incision. With the ulcer or growth exposed to sight, an outline incision with the knife may be made to encircle it well outside its margin, and the removal may be completed with scissors, care being taken to include in the parts removed a considerable margin of healthy tissue. The arrest of hæmorrhage is not usually attended with difficulty; but few vessels call for ligature, and venous bleeding is best put a stop to by the temporary insertion of a well-squeezed sponge. In some I have found it practicable to adjust the cut edges of the mucous membrane, but I have not observed that the care bestowed upon this point has been attended with compensatory advantages. In the greater number the wound has been left unclosed, care being taken to keep it cleansed by the free use of iodoform or other antiseptic, and patent by means of celluloid drainage tubes, so as to avoid accumulated secretions or concealed hæmorrhage. Much of the success of the operation, I am convinced, depends upon the thorough cleansing of the wound not only at the time of operation, but during many subsequent days. I have usually had the rectum washed out with some antiseptic solution every eight hours, and have followed this each time by the insertion of an iodoform suppository. By attention to details such as these, dangers are averted and the healing process is much accelerated; and thus the operation may be expected to take rank with those for the removal of cancer in other situations. And yet it will be generally conceded, I think, that in the existing state of our knowledge and experience, proctectomy is an operation of limited application, unsuited to more than a minority of the cases presented to us. When the disease is well within reach of the fingers, and confined to the posterior or lateral portions of the rectum, or if, situated anteriorly, it has not invaded the tissues outside the muscular wall of the bowel, and when also the external sphincter can be preserved in functional activity, I have no hesitation in recommending its adoption; but in those cases in which the disease has originated beyond the reach of the finger, or, having commenced lower down, has crept up to a higher level, we must still, I think, trust to those other less radical measures which till recently constituted our only resource.

In none of my cases have I seen fit to precede the removal of the disease by an opening in the colon, and yet I cannot but think

a complete diversion of the fæces as a preliminary measure would prove an advantage in proctectomy. The passage of fæces over the wound, besides complicating and delaying the healing, materially adds to the dangers and miseries of the patient. It is a question for the future to determine how far by combining the two operations of colotomy and proctectomy we may succeed in still further prolonging the duration of life.

We proceed now to the consideration of the second of the operative measures, namely, colotomy, and as bearing upon this part of my subject it will be interesting and instructive to note a few points in what may be termed the natural history of cancer in the rectum. Before considering how we can best postpone death—for, apart from possible eradication of the disease by proctectomy, we can hope for no more than a prolongation of life—let us inquire in what way death usually comes in these cases. Of 102 patients, seen during a period of ten years in consultation with their medical attendants or at my own rooms, for whose histories I am much indebted to my medical friends, I find that seventeen died of complete obstruction, unrelieved by any operative interference, whilst fifteen others, similarly obstructed, underwent the operation of left lumbar colotomy. From this it would appear that about 31.4 per cent of all cases of cancer of the rectum would die of obstruction if left to Nature. Of the remainder, seventy in number, three died from hæmorrhage, five are said to have died of acute septicæmia, two from implication of other organs—the peritoneum and abdominal lymphatics in one, the vagina, uterus, and bladder in the other. In nine the operation of left lumbar colotomy was performed, in the absence of complete obstruction, for the relief of local distress; whilst, in the remaining fifty-one, exactly one half of the whole number, death is attributed to “exhaustion,” and in several of these such complications are mentioned as “œdema of legs,” “ascites,” “urinary difficulties,” implying doubtless extension of the disease to neighbouring organs and tissues.

At first it seems a little surprising to find that less than one in three, if unrelieved by operation of any kind, will die of obstruction. It must be borne in mind, however, that it is complete obstruction which is here referred to. More or less interference with the passage of the intestinal contents is met with in a much larger proportion of cases, in nearly every case indeed at some period of the disease; but it is an incomplete stoppage which may be got over by ordinary measures, such as injections, the introduction of the finger, or of bougies, the use of laxatives, and the like, and which, though painful and exhausting, is not by any means the sole factor in causing death. It is a pregnant fact, and one worth dwelling upon, that in not one of the thirty-two obstructed cases above referred to, in which colotomy was imperatively called for, was the disease situated within easy reach of the finger; and in the majority it was quite beyond the touch, its position being diagnosed by other agencies, or inferred from the past and present history of the patient. I have for some time held, and have often given expression to the opinion, that in cancer of the lower half of the rectum we have not much fear of the occurrence of complete stoppage. I may now add with equal confidence that where the upper portions of the rectum are involved complete blockage is almost certain sooner or later to appear. This latter assertion would seem warranted by the results of the thirty-two cases I have quoted; and of the truth of the former I have several times met with most striking confirmation. Let me mention two cases. In August, 1883, Mr. Griesbach, of Garforth, sent me an elderly man, who had been suffering since the previous January, and of whom, after examination, I wrote to his doctor:—“An immediate colotomy is desirable to avoid complete obstruction, which seems imminent. The hard ring of growth has almost obliterated the lumen of the rectum.” He declined operation, and yet his death did not take place till January, 1885, seventeen months after I saw him, and at no time was he threatened with anything like complete obstruction of the passage. In February, 1884, I saw, with Dr. Hall, of Leeds, a patient whose rectum, at a distance of three inches from the anus, was encircled by a thick nodular ring of carcinoma, through which it would have required force to pass one's little finger. Colotomy was pressed upon him as needed to avert an early death from obstruction. He declined; and yet he lived on till February, 1887, a period of three years, without realising the agonies I had predicted. The reasons for this difference in the effect of cancer as a cause of occlusion are doubtless to be found in the anatomical relations of the upper and lower portions; for whereas the rectum as it approaches the outlet becomes more closely applied to the sacrum and pelvic wall, in its superior por-

tion it is comparatively free, and thus the contractile action of the colon above is exerted with effect in forcing the contents through a contracted ring when that ring is fixed and immovable; whereas when the narrowed portion is free, movable, and not attached as it is when seated in the upper portions of the rectum, the efforts of the bowel above succeed only in invaginating or otherwise displacing the growth, often so as to enable us to make a complete diagnosis by bringing the disease within reach of the finger, and fail altogether in effecting any onward movements of the contents.

If, then, we are satisfied that we have to deal with a case of cancer situated in the upper movable part of the rectum, knowing that complete obstruction may of a certainty be looked for, we shall probably all agree that it is desirable to make an artificial anus in the colon above as soon at least as the first symptoms of impending blockage appear, and in time to anticipate those further changes upon which the mortality of colotomy so much depends. And I will here state my opinion that in these circumstances the ordinary left lumbar colotomy is to be preferred, not only because of the facility with which the distended bowel is reached, but chiefly because the opening is made at a sufficient distance from the disease to ensure its being found in a healthy state, free alike from cancerous infiltration and from those ulcerative changes which are apt to occur on the distal side of a stricture canal. That an early operation is called for is abundantly shown by a reference to the mortality in the cases just mentioned. Of the fifteen operated on for complete obstruction, five died within the first week, two within three months, and the remaining eight lived an average of rather more than twelve months, including one who survived the operation twenty-two months. On the other hand, of the nine operated on when no symptoms of obstruction were present, the earliest death did not take place for upwards of two months, and the average of life throughout was a little over eleven months. And, again, reviewing a larger series, I find that out of fifty-four cases of colotomy taken from both my hospital and private notebook, performed for the relief of distress only, and not for obstruction, three only died during the first month, and in each of these there were some special circumstances, such as advanced age or great emaciation, to account for the early death. It is not in cases of complete obstruction or of threatened obstruction that any difficulty arises in the course to be pursued. When death is imminent and other means have failed to bring about relief, an opening somewhere and of some kind in the bowel for the exit of the pent-up excreta is obviously called for; and, as we have seen, the earlier that opening is made the more certainly will a lasting relief be obtained. And, following out the result of my observations, I would say that, given a diagnosis of cancer high up in the rectum—in the earlier stages, it is true, by no means always an easy matter—the duty of the surgeon would seem clear, namely, to recommend without delay the formation of an artificial anus.

There remain, however, still to be considered those cases of disease affecting the middle and lower portions of the rectum, to which excision is deemed inapplicable, and in which obstruction to any serious extent is not likely to happen. How are these to be dealt with? The propriety of recommending colotomy here must turn upon the answers to two questions—is the life of the patient prolonged by its adoption? are his remaining days made more tolerable? In order to answer these questions I have made inquiry into the life history of eighty-six patients, of whom fifty-two ran their course without operation, and thirty-four submitted at some stage or other of the disease to left lumbar colotomy. In this inquiry I have taken care to exclude all doubtful cases, and I have limited it to such in my private practice as have within recent years come under my own personal observation, in order, as far as is possible in such an inquiry, to exclude elements of uncertainty. The average duration of life in those who were not operated on, reckoned from the date of the earliest symptoms complained of, was a little over seventeen months; whilst that of the thirty-four who underwent colotomy was twenty-two and a half months. This gives an average extension of life in each case operated on of five and a half months. Again, the average duration of life in the thirty-four after the time of operation was fifty-three weeks, whilst in thirty-seven out of the fifty-two cases in which I have put on record in my notes the fact that I have recommended colotomy, and in which my advice was not taken, I find the average of life, reckoned from the time of my seeing them, reached to a little over thirty-four weeks, thus giving an additional nineteen weeks of life to those who submitted to colo-

tomy. It is not unfair, perhaps, to assume that the patients who consented to operation were suffering more severely, as a rule, than those who for some reason declined; and, if so, the duration of life in them had they refused colotomy would probably have shown a shorter average than in those who, suffering less severely, declined the surgeon's aid. Be that as it may, there is ample evidence in the foregoing facts to show that in recommending the formation of an artificial anus in the cases under consideration we are justified by the knowledge that by its means a substantial addition to our patients' lives may be procured.

It is, perhaps, worth remarking that the proportion of patients who have declined operation has increased during the last few years. I believe the cause of this is to be found in the fact that for some indefinable reason my faith in the value of colotomy had gradually waned; and we all know how much patients are influenced by the mental attitude of their advisers. On commencing this review of my own work and experience, I am free to confess it would have given me no surprise had the results turned out upon the whole unfavourable to colotomy. The actual results have had the effect of re-establishing, nay, of materially raising, in my mind the value of colotomy as a means of prolonging life in all sufferers from cancer in the rectum.

In the next place we have to consider how far the patient's physical condition is improved, to what extent his sufferings are alleviated by the operation. Bear in mind I am dealing only with cases in which there is no material obstruction to the passage of fæces, and which, on the other hand, are characterised by the distress and worry of frequent ineffectual evacuations, and by unceasing pain and discomfort, due in great measure to the presence of excreta in the rectum, and of more or less loss of control over its expulsion. The amount of relief to be obtained is to be measured by the extent to which suffering depends upon the presence of fæces and of their passage over the diseased surface. Complete relief is never obtained. A variable amount of pain and distress is inseparable from the condition itself. To this fact must be attributed the not infrequent expressions of disappointment which escape the lips of our colotomised patients. We have not here the comforting assurance of a respite from the inroads of the cancer which comes of excision, or the obvious escape from actually impending death from complete obstruction. The mental relief obtained in the one case, the rescue from death, and the palpable physical improvement in the other, are absent here. However real the actual relief, it is never complete; and there is still the prescience of a lingering death. These militate, except in the most sanguine, against realisation of benefit received. The colotomised patient has not experienced the miseries of him who has declined operation. His own sufferings are not small; to speak to him of an escape from greater seems a mockery. The attendance upon dying patients who have willingly accepted every surgical aid at our command is not calculated to impress the surgeon even with an overwhelming sense of benefits conferred. It is probably from frequently witnessing the last weeks of these patients, who remain under observation more persistently after operation, that I had gradually come to question the advisability of opening the colon where life was not immediately imperilled. A careful and impartial comparison of their sufferings, however, in a series of patients, including some who had declined and some who had undergone operation, has convinced me of the great utility of colotomy, and has completely dispelled my doubt as to its alleviating power. The conclusions I have been able to draw are that, by its means, the continuous pain is lessened in severity; the desire to evacuate, almost constant, and very distressing where no operation is performed, disappears in some and is diminished in others; incontinence of fæces, a frequent source of anguish in rectal cases, though occasionally met with, is usually absent in the colotomised; and, finally, the motions are not only discharged with comparative ease, but, in the majority, are reduced to at the most two or three in the twenty-four hours. These are substantial advantages, generally, I am aware, though perhaps vaguely, claimed as pertaining to colotomy, which it may be well for this meeting to acknowledge, to qualify, or to repudiate.

Finally, there remains for consideration to determine what kind of artificial anus best supplies the needs of the patient, and obtains for him the nearest approach to a state of Nature. Till recently, one operation, that, namely, of left lumbar colotomy, practically held the field. Inguinal colotomy was here and there undertaken, and now and again there appeared a paper founded upon one or two examples, setting forth its advantages over the other proceeding; but it was not until the last two years, since Mr. Allingham,

jun., read his paper on this subject in the Surgical Section at the Dublin Meeting, that hospital surgeons in general have taken up the question in earnest, and have put to the test the advantages claimed for this over the usual operation. I have not myself yet performed the inguinal operation in any case of cancer; but there are doubtless many gentlemen present who will give us their experiences, and so will enable us to form a judgment upon its merits.

I am bound to confess that, having opened the descending colon in the left loin 103 times, I find it in some respects an unsatisfactory operation. Now and again it is not unattended with difficulties; but I have never found these insurmountable. In stout persons it will sometimes be found that the bowel is in front of the anatomical line; in the thin and wasted, the empty gut may have dropped down, as it were, so as to form an attenuated cord by the side of the spine. In these cases the peritoneum may require to be opened; and in others an opening may be unintentionally made. With cleanliness and care in avoiding the introduction of putrescible material, there need be no fear of inserting the finger freely within the cavity of the abdomen in search of the truant bowel, and the accidental wound need not so much interrupt the even tenour of the operation. It has happened to me more than once to have opened a tightly distended colon and the peritoneal cavity by the same incision, and to have become suddenly conscious of the accident only during a copious flow of liquid feces over the entire wound; and yet I am not aware that any disastrous effects have ensued. The tightly distended bowels secure an accurate closure of the wound, and I have always taken care to second this by seizing the cut edge of the colon with forceps and dragging it to the surface. It is not on account of the difficulties we sometimes encounter, or of the accidents which cannot always be avoided, that I am dissatisfied with this long established operation; neither is it because of the high mortality which undoubtedly attends it when undertaken for relief of complete obstruction, a mortality which I am prepared to show arises more from the circumstances preceding the operation than from the operation itself; but rather because of its failure in a number of instances to secure that complete diversion of the feces which it is intended to bring about, and because of a tendency to a gradual stenosis and even closure of the opening, on the one hand, or of unmanageable prolapse of the bowel on the other, which in a not inconsiderable number of cases detracts materially from the comfort to be derived from a perfect operation.

It is claimed for inguinal colotomy that, when rightly performed, there is no possibility of feces finding their way beyond the opening in the groin, that the removal of a portion of the free border and the kinking of the bowel secure a continued patency of the artificial anus, and that prolapse may be effectually prevented by the use of a properly constructed truss. If these claims should prove to be well founded—and the examples I have seen in London, in Leeds, and in Newcastle would seem to support them—the chief drawbacks of colotomy will have been removed. The lumbar opening will continue to be preferred when the disease is so high up as to render it uncertain if an inguinal opening will enable the bowel to be reached, and probably also in cases of great distension where an immediate opening for relief of obstruction is imperative. Not long ago a man, whose colon I had some months previously opened, returned in a condition but little, if at all, improved by the operation, which had wholly failed to prevent the dribbling of feces through the rectum, although the artificial opening had remained patent as at first. To his complete relief I performed the operation of colectomy. The colon, having been cleared from its attachment to the lumbar opening, was dragged outside and completely divided. Having then temporarily secured the upper end with an encircling ligature of silk, so as to prevent the escape of feces, and given it in charge of an assistant, I first dealt with the lower portion, and finding it empty, I was able at once securely to close it by stitching with fine silk its invaginated wall in such manner as to oppose two serous surfaces throughout the entire circumference. Thus safeguarded against any back flow of its contents, the distal end of the bowel was dropped into the abdominal cavity. The upper open end was then stitched all round to the edges of the lumbar wound, and in this way the whole of the contents of the bowel were perforce discharged through the lumbar opening.

How far this operation may be safely undertaken as a primary proceeding we have not sufficient evidence to show. Should it prove on trial to be not more fatal than the ordinary method, it is clear that by its means we shall more completely accomplish

the chief end we seek to attain—namely, the diversion of the intestinal contents through the loin, and the detaching of the diseased rectum from the alimentary canal. It has been proposed, indeed, and even, I believe, essayed to accomplish, the removal of the cancerous bowel through the lumbar or inguinal opening, and that such a procedure may, in exceptional instances, be warranted, I can well conceive; but for the present our choice of operation in cancer of the rectum would seem to lie between proctectomy as performed from the anal extremity on the one hand, and some form of artificial anus on the other.

AN ADDRESS

DELIVERED AT THE OPENING OF

THE SECTION OF PATHOLOGY AND PHYSIOLOGY.

At the Annual Meeting of the British Medical Association, held in Leeds, August, 1889.

By JOSEPH COATS, M.D. GLASGOW.

Lecturer in Pathology and Pathologist, Glasgow Western Infirmary.

ON THE NATURE OF CONSTITUTIONAL SUSCEPTIBILITY TO DISEASE.

GENTLEMEN,—In opening the Section of Physiology and Pathology it is fitting that I should recognise the close relationship between these two branches of science. The methods and leading principles of physiology have, with great advantage, been extended to pathology, and through pathology to practical medicine and surgery. We are apt to think that this is a special characteristic of recent times, but we must not forget that in the work of our great master, John Hunter, this combination was exemplified in the highest degree. Not only is his work largely physiological, but in his marvellously full and suggestive lectures on the Principles of Surgery, it is only after a prolonged discussion of the fundamental principles of physiology that he goes on to the principles of disease. It is, I think, matter for congratulation that, following John Hunter's example, modern pathology does not confine itself to pathological anatomy, but recognises that the processes of disease occur in living animals, and are to be investigated from the physiological as well as from the anatomical standpoint.

MISUSES OF PATHOLOGICAL TERMS.

It must be recognised that pathology has made great advances in modern times. It follows from this that the meanings and uses of terms must have undergone considerable changes in the course of years, and that confusion is liable to occur between the older and more recent uses of these terms. This applies especially to terms which imply theories or views, because as the views change the terms either become obsolete or else linger in some indefinite sense, to the confusion of the minds of men. On the other hand, a term which expresses some great fact of fundamental importance must remain, although the attitude towards the fact and the interpretation of it may undergo alteration. I propose to-day to take up in this aspect one of the expressions belonging to the latter class, but before I do so let me give two examples of the former.

The term "lymph" in its pathological use is, I believe, a Hunterian expression. Hunter's "coagulable lymph" was primarily the fibrin which forms when the blood coagulates. He says that the blood separates into the coagulable lymph, the serum, and the corpuscles. It is well known that Hunter ascribed very special powers to the coagulable lymph, regarding it as the vehicle of the living principle of the blood. What we now regard as dead fibrin was to him the most active and vital matter in the body. Hunter regarded the new formation of tissue as brought about by this lymph, so that even the extension of a scirrhus cancer to the axillary glands, for example, was by the passage of the coagulable lymph from the tumour in the mamma. Our views of all this have entirely changed, but the term unfortunately remains, and is used, especially by surgeons, in what I cannot but regard as a loose and misleading way. Thus, not only is the fibrinous exudation of acute inflammation called lymph, but the connective tissue