amount to be removed in one bone section simply depends upon the extent of the disease. We try to avoid any encroachment upon the epiphysary line, but we must do it if the disease brings us there. Our object is to save the patient's limb; and if we have to do that at the expense of much shortening, we cannot help it. But my own experience has not been of this kind; and I believe it will be found that this misfortune is ideal in the great majority of cases.

I believe that once we determine upon cutting open a knee-joint in general tuberculosis of the synovial membrane, we had better go the whole way, and perform the ordinary excision. The operation is, I maintain, safer and more satisfactory in its general results than erosion. My own hospital experience, and as far as it extends to the practice of my colleagues, is a mortality of one in over fifty successive cases. That leaves little to be desired; and I am satisfied that the patients who have been submitted to excision of the joint have on the whole been better treated than if they had undergone exploratory incisions and tentative scrapings at the hands of a surgeon.

ON DISEASE PREVENTION.¹

By STANLEY HAYNES, M.D.,

President of the Worcestershire and Herefordshire Branch.

I HAVE thought a few notes on disease prevention as influenced by medical politics might be acceptable for your kind consideration, and it is with this hope that I venture to solicit your patient attention and forbearance for a few minutes.

County Councils.—My chief reason for bringing this subject forward now is because I believe that if we combine, as by means of this and other Branches of the British Medical Association, to endeavour to influence the County Councils, created by the new Act, in the direction of public health there seems, now and in the near future, to be a very favourable opportunity of exerting our knowledge and strength to some beneficial purpose. For instance, the Branches might request County Councils to recommend the District Councils to amalgamate in the appointment of medical officers of health; and the continuance of the faulty system of each local board and sanitary authority appointing its own separate health officer should be discontinued, as by a County Council intimating that it would instruct auditors of district accounts to disallow the salaries except for sufficiently combined areas: this would soon cause combination of districts and the appointment of skilled medical officers of health for large districts; they should receive good incomes (beginning at £600 per annum, with allowances for expenses) and be prohibited from private or other practice. It is gratifying to observe that many medical men have been nominated for election on the County Councils, on which they will do much to advance State medicine and the interests of our profession in public work. Let us wish them all success in their candidatures. The medical members of County Councils could bring forward, among many other measures, the desirability of having a Minister of Health in the Cabinet, and the fixity of tenure for medical officers of health.

Disease Notification.—The compulsory notification of diseases has, with few exceptions, been so satisfactory that this seems one of the matters we should endeavour to extend. I am one of those who consider this notification, when made by medical men, should be rewarded by fees paid by local authorities, and that the penal clauses adopted by some towns (and very curiously acquiesced in by their practitioners) are uncalled for, are derogatory to the dignity of our profession, and calculated to arouse opposition against such publication of illnesses. Such notification should be made by the householder.

Inherited Disorders.—Hereditary tendencies to disease and crime and alcoholism should be repressed as far as possible. The strict fulfilment of the Inebriates Act would do something to lessen drunkenness, idiocy, insanity, epilepsy, and other neuroses in the next generation. So far as we can we should discourage the marriage of persons with histories of cancer, consumption, syphilis, and alcoholism. Convicts should be detained in penal servitude during the full terms of their sentences, in order that they might have fewer opportunities of transmitting criminal propensities: the cost of this to the taxpayers would be minimised by the adoption of the beneficent industrial system in successful operation in the Indian and Burmese convict service.

Vaccination.—Vaccination and revaccination should be compulsory and enforced. The recent memorandum issued by the Local Government Board on revaccination is an important authoritative indication of the value of this procedure, and it is to be hoped both the public and vaccinators will avail themselves of the facilities rendered by this salutary action of our Government.

Inebriety.—One of the pressing wants of the community in the prevention of disease is compulsory power to place inebriates, of all classes, in homes provided for their reception and superintended by physicians. These homes, for the middle and lower classes at least, should be industrial. The Inebriates Act lately passed is a slight improvement on the Habitual Drunkards Act (chiefly, in name) but, while thankful for small mercies, we should regard it as only an instalment towards wiser legislation, because it does not go far enough. Some of our colonies and the United States Government are far ahead of us in regard to this legislation. Power is wanted here for the isolation of inebriates for twelve months as a minimum: no appearance before a magistrate should be necessary, but notices of admission, in writing, should be posted, within twenty-four hours, to the Clerk of the Peace and to the chairman of a visiting committee of magistrates, three of whom should visit each inebriate so segregated. Grocers' licences to sell intoxicating drinks should be revoked, and it should be penal for any grocer or unlicensed tradesman to give such drinks to any customer or servant of a customer.

Leprosy.—Leprosy has attacked some of the populations of Queensland and Natal. Those and any other countries under British rule should have power to separate those afflicted with the disease, some interesting notes on the spread of which in Spain appeared in the JOURNAL of January 12th, 1889. Norway and Hawaii have set us brilliant examples in this respect with great success. As an example of the ignorance prevailing concerning the contagiousness of leprosy, it may be mentioned that last year an action was brought—happily it failed—against the head master of one of the largest schools in Scotland, because he had refused to receive a boy suffering from that disease.

Veneral Diseases.—Those conversant with the state of our ports and of our troops in India can, unfortunately, give too ample evidences concerning the alarming increase of venereal disorders resultant on the suppression of the Contagious Diseases Act, the medical men testifying to the type as well as the frequency of such maladies, while the police know how prostitution has increased, and how young girls—that is, from 12 to 15—who, while the Act was in operation, were kept off the streets, are now found on them, as before the enactment.

Coroners.—Coroners should be selected from medical men only, and be prohibited from practice, their districts being made sufficiently large to furnish adequate salaries. This is one of the topics on which our profession could appeal to the County Councils.

Cremation.—Cremation (on which Sir Henry Thompson wrote in the *Contemporary Review* in 1874, and the first lecture on which by a medical man was, I believe, delivered by myself—in 1876) is gaining ground rapidly amongst the educated and thinking portion of the community; but we are far behind some of our European neighbours and transatlantic cousins in resorting to this healthful mode of disposing of the dead. This is a reproach under which we should not be satisfied to rest. Some very necessary attention to cremation has lately been given at Glasgow, while an urgent need for the process has been revealed in Edinburgh. At Glasgow it is calculated the actual incineration would be completed in half an hour, at a cost of 2s. only each body; at present the ratepayers are paying 13s. 11d. for each pauper interment. A hundred persons have joined a Cremation Society in Glasgow; it is to be expected this enlightened undertaking will be speedily followed by other centres of population, in consequence of the horrible revelations recently made in connection with burial. Incineration of corpses is as necessary for the lessening of human disease as for the prevention of maladies in our cattle; its extension and general adoption would, besides diminishing the spread of many kinds of illness, prevent the serious risks to health now incurred by mourners, ministers, and others assisting at funerals. Burial in brick or stone graves should be absolutely prohibited by the Home Office, while medical practitioners should do all they can to cause the substitution of quickly perishable coffins in place of the massive ones now too frequently used, when cremation is impracticable.

Isolation of Infectious Diseases.—The JOURNAL of April 30th,

Abstract of a Presidential Address read before the Worcestershire and Herefordshire Branch of the British Medical Association, at Malvern.

1870, p. 439, prominently called attention, by a leading article, to the importance of establishing "isolation homes" for patients suffering from infectious fevers. As I deem this paper would be incomplete without some slight mention of epidemic diseases, I propose to conclude, with your permission, with the reading a letter of mine published in the JOURNAL of May 28th, 1870, p. 564. [This letter was then read.]

CLINICAL MEMORANDA.

TREATMENT OF SCARLET FEVER BY THE BINIODIDE OF MERCURY.

R. K., aged 9, began on September 30th with sore throat and fever. Summoned on the following day, in the evening, I found him suffering from scarlatina anginosa, and the rash well marked. I gave the following mixture: Sol. hyd. bichlor $\mathfrak{z}$ vi; pot. iod. gr. xv; sp. am. co. $\mathfrak{z}$ j; syrup. $\mathfrak{z}$ ss; aq. ad $\mathfrak{z}$ vi; $\mathfrak{z}$ ss secundis horis. Throat to be sprayed every four hours with 1 in 2,000 biniodide in sodic iodide.

October 2nd. Rash fully out; restless, sleepless, and very feverish.

October 6th. The following note made: The rash became less and less marked each day, until now there is no trace of it. Healthy granulation of the ulcerated tonsils. Pharyngeal glandular swelling externally decreasing. Bowels moved once each day. Temperature normal. The rest of the family occupied the same room.

October 9th. Medicine changed to iron and chlorate of potash with biniodide in one-sixteenth grain powders three times a day.

S. A., aged 3, began with scarlatinal throat but no rash, furred tongue, vomiting, and feverishness, and was treated as for quinsy.

October 10th. S., aged 5, began with high fever (102°) and rash all over the body. Ordered the following: Sol. hyd. bichlor. $\mathfrak{z}$ ss; pot. iod. gr. viii; sp. am. co. $\mathfrak{z}$ ss; syrup. $\mathfrak{z}$ iii; aq. ad $\mathfrak{z}$ iss; $\mathfrak{z}$ j secundis horis.

October 11th. S. passed a bad night. Objects to the spray. Temperature 102.7°. Throat much inflamed and covered with exudation. The application of a 1 in 500 solution of biniodide by laryngeal wire-handled camel's hair brush, to the throat, ordered three times a day, the father undertaking to see to it. S. A.'s case, now one of well-marked fever with rash, and treated similarly.

October 13th. J., aged 10, began with high fever, sore throat and rash. Had same mixture as R. Tonsils and pharynx much inflamed and covered with exuded lymph.

October 14th. All doing well except J., who began with purging in the night. The mixture was at once stopped, and the liquor ferri perchlor. fort. in $\mathfrak{z}$ ss minim doses ordered, in glycerine and water, every $\mathfrak{z}$ or 2 hours, and the biniodide in one-sixteenth grain powders three times a day.

October 15th. J.'s diarrhoea was rapidly cured by the iron. Each case had throat-painting, with the 1 in 500 every four hours. S.'s skin quite free from rash. Temperature 99°. Sleeps and eats well.

October 16th. S.'s temperature 99°. J. improving. S. A. to get up.

On the 11th, M., aged 2 years, had inflamed throat, and took five or six doses of the biniodide mixture; but no rash ever showed itself, so little notice was taken of the child.

October 20th. J.'s temperature normal.

October 24th. R., having passed the twenty-second day without kidney symptoms, may be considered free from the danger of any sequelæ. The slightest scurfy desquamation noticeable, and that chiefly affecting the fingers and toes.

October 28th. S., after severe earache and swelling under the ear, had a discharge of matter from the left ear.

On the 30th I was called in, and prescribed the 1 in 2,000 lotion to be dropped in, and the mixture with ammonio-citrate of iron to be taken three times a day for the pain remaining. M., who had no throat-painting, was now suffering from a large abscess in the neck. Poultices were applied and it was incised on November 6th, a large quantity of pus being evacuated. It healed in ten days with carbolio ointment.

REMARKS.—All four cases had the temperature reduced to the normal in six days, and were convalescent on the seventh. None had dropsy. In one, not at first "throat painted," the disease caused otitis by extension along the Eustachian tube. The only case of cervical abscess was one in which there had been no local appli-

cation made. In one, J.'s, urgent purging came on within twenty-four hours, and the throat lesion was noticed to be pharyngeal rather than tonsillar. This was immediately checked by iron frequently given. As regards infection, I have since found that by painting the throat with the 1 in 500 solution every four hours, the disease is prevented from spreading. Desquamation in all was merely scurfy and chiefly affecting the fingers and toes.

Accrington.

C. R. ILLINGWORTH, M.D., M.R.C.S.

SURGICAL MEMORANDA.

SYNOSTOSIS OF THE RIBS IN EMPYEMA.

SYNOSTOSIS of the ribs in empyema is not, I think, so uncommon as Mr. Pye believes, nor is it a mere coincidence. In my experience, I should go so far as to say it is not uncommon to find bridges of bone uniting adjacent ribs in cases where a drainage tube has been long worn. When resection of one or more ribs has been performed without removal of the periosteum and surrounding soft parts, the bone being merely shelled out, I have found subsequently that the gap has been filled up by a definite bony shield, a more extreme but similar condition.

St. John Street, Manchester.

G. A. WRIGHT.

THERAPEUTIC MEMORANDA.

EUCALYPTUS IN CANCER.

I LATELY had a case of cancer of the breast in which the ulcerating surface was about 8 inches by 6. The odour from the discharge was most sickening and offensive, and it was made worse by the odour of the carbolic acid which they were using. No one could remain in the room long with the patient on account of the intolerable smell. The carbolic acid was stopped, and the eucalyptus disinfectant used. The wound was sprayed with it, as also the lint with which it was dressed; tenax was wetted with it and placed under the wound to catch the discharge, and, by the use of a spray diffuser, the air of the room was filled with the fragrant odour of the eucalyptus. All offensive smell at once ceased, and there was only the smell of an ordinary suppurating surface.

In a case of cancer of the sigmoid flexure where the discharge was frequent and very offensive, the injection of a small quantity of eucalyptus emulsion twice a day stopped the offensive odour, and checked the frequency of the discharges. And in a case of cancer of the neck of the uterus, the offensive smell was stopped, the discharge lessened in quantity, and the tendency to hæmorrhage diminished.

J. BRENDON CURGENVEN.

Bayswater.

REPORTS

ON

MEDICAL & SURGICAL PRACTICE IN THE HOSPITALS AND ASYLUMS OF GREAT BRITAIN, IRELAND, AND THE COLONIES.

STAMFORD INFIRMARY.

SHORT NOTES ON RINGWORM: TRANSMISSION FROM COWS.¹

(Under the care of Dr. W. NEWMAN.)

F. F., a male, aged 56, was admitted on February 22nd, 1881. He had on the chin a hard swelling extending on each side of the symphysis for fully two inches in length and about an inch and a half in depth. It was raised about a quarter of an inch above the level of the skin, somewhat reddened, boggy to the touch, and discharging pus from several small points. About the middle there was an island of perfectly sound skin. On the right cheek also there was a small hard tumour, over which the skin was red and tender to the touch. There was no enlargement of glands.

He was a farm labourer, in good health; he had had charge of cattle lately; among the herd several had ringworm severely. He was quite certain that sixteen days ago (day named) he had no swelling on or beneath the chin, for on that day he was able to shave as usual. Ten days ago he was unable to shave from the

¹ Read before the Midland Branch of the British Medical Association.