of the pleura was an expression often used in a somewhat vague way, and the idea that seemed to be in most people's mind was that the pleura was thickened by the formation of a mass of new material between the parietal and visceral pleural surfaces. He pointed out that the lymph so deposited had very little to do with the subsequent formation of cartilaginous tissue, which was, for the most part, due to collapse of the superficial layers of air cells, and their fibroid degeneration. He agreed with Dr. Powell in his estimate of the value of so-called antiseptic treatment of tubercle. Until surgeons could show that they could successfully treat a tubercular affection of a synovial membrane, which was accessible to every kind of local treatment, there was no reasonable ground for thinking that these means, given by the mouth or inhaled air, in vapour or spray, could be effective. He protested against the suggestion of Dr. Churton, that the chest should be incised in simple pleurisy, and denied that there was any analogy for thinking that such an operation would prevent the occurrence of tuberculosis. With reference to the remarks of Dr. Drysdale, he wished to recall the words of Virchow, who, in his address at the Medical Congress in 1881, said that the recent discoveries of specific micro-organisms brought into greater prominence than ever the old doctrine of predisposition.

IX.—F. IMLACH, M.D., Liverpool.

DR. IMLACH had been much interested in Dr. Powell's observations on cases of tuberculous peritonitis in which no primary lesion of any abdominal or pelvic organ could be found. Either the primary lesion had become obsolescent, and was without clinical importance, or there was none. In such cases abdominal section with washing out and drainage resulted in an apparent cure, as he had found in five instances. In these the tuberculous nodules of the peritoneum were previously unsuspected. But when there was tuberculous ulcer of the intestines surgery seemed to be of little service, and when the lungs or liver were affected, of none at all. He had watched the progress of one of his five cases for four years, and there had been no return of the disease.

X.—JULIUS DRESCHFELD, M.D., F.R.C.P., Professor of Pathology, Victoria University, Manchester; Physician to the Manchester Royal Infirmary.

DR. DRESCHFELD said he was led to draw attention only to one or two points of pathology which might explain the peculiar behaviour of cases of tuberculous peritonitis as regards prognosis. When a tuberculous peritoneum was examined tubercle bacilli were often found to be absent. In such cases the spores were present but no bacilli. Such a condition was found also in so-called healed phthisis, where no bacilli were found, but where inoculation in animals with particles of the caseous nodule produced typical tuberculosis, etc., and it might be assumed that a very similar condition occurred in peritoneal tuberculosis, and that many of these cases might recover when the spores did not develop further, or they might, when a particular predisposing cause arose, spring into activity and produce acute tuberculosis or general tubercular disease.

THE PRESIDENT (Dr. J. E. Eddison) commented on various points that had been raised during the discussion, and Dr. POWELL replied.

NOTES OF A CASE OF TREPHINING FOR OLD DEPRESSED FRACTURE OF THE SKULL: RECOVERY.

By GEORGE BUCHANAN, A.M., M.D., LL.D.,
Professor of Clinical Surgery in the University of Glasgow.

THE following case is a good example of the beneficial effects which may follow removal of pressure from a localised spot of the brain, even long after the injury received. The notes of the case are by Dr. Dunlop, of Thornhill, who sent the patient to me and took charge of him after the operation.

P. M., aged 33, a railway surface man, received a compound fracture of the skull by his head coming in contact with a bolt

projecting from a bridge under which a train was passing, he being at the time standing in one of the wagons. The seat of injury was the occipital bone, near its upper angle, about an inch or more to one side. He was stunned by the blow, and was taken in an unconscious state to Dumfries Infirmary, and remained so for three days. Eight days afterwards he was conveyed home, and was, as far as can be ascertained, fairly well for six weeks, when he again became unconscious and was confined to bed for three months. At that time he complained of great pain in the top of his head. The headaches were sometimes so severe that they were accompanied by fever and sometimes delirium. Blisters were frequently applied to the nape of the neck. He recovered sufficiently to allow of his going about his ordinary occupation; married, and lived comfortably in a little cottage some miles from Thornhill.

Six years ago, the pains in the head again became worse, and symptoms of gastric derangement showed themselves. The pain was not uninterrupted, but came on occasionally with great severity, completely unfitting him for all duty, and causing indigestion and inability to take food.

In April, 1889, he called on me, and I found at the seat of the old injury a manifest but very slight depression in the skull. From this point he declared that the pain always arose, and felt as if shooting all round his head. I told him that the only hope of recovery was the removal of a piece of bone from the site of the old injury; but that the result could only be problematical, and the operation not devoid of danger.

During the summer he went for change of air to the sea coast, but not finding any improvement, but rather an increase of pain, he determined to have the operation performed. I did this on July 29th at his house near Thornhill, with the assistance of Dr. Dunlop and my son.

The head being shaved, the cicatrix of the old wound on the scalp was seen at a point about an inch and a half from the superior angle of the occipital bone to the left side. A crucial incision having been made over this spot, and the flaps retracted, I could see and feel the edge of the old fracture, with a depression about an eighth of an inch. I applied a trephine, and removed a disc of bone the size of a shilling, and had the satisfaction of finding I had got the bit which had been pressing inwards. There were no starred or rough edges. Whatever had been smoothed away; but there was a projection inwards, something like a bone button—making the skull in that part two or three times thicker than normal, and the thickness pressing in on the membranes. There was no abnormal appearance on the membranes. I stitched the points of the flaps together, and applied a simple dressing of alembroth gauze and cotton. The wound healed by first intention. Dr. Dunlop only changed the dressing two or three times, and the patient made an excellent recovery.

The headaches have entirely disappeared (December, 1889), his appetite and digestion are both good, and he feels himself "a new man," as he expresses it.

NOTES OF A LECTURE ON SOME FORMS OF RETINAL PIGMENTATION.

Delivered at the Royal Westminster Ophthalmic Hospital,

By W. ADAMS FROST, F.R.C.S.,
Surgeon to the Hospital.

I do not propose here to deal at all exhaustively with the subject of pigmentation of the retina, but merely to illustrate some forms in which it occurs, which appear to me to be in some degree typical of their kind. It is, I believe, generally admitted that in all cases in which pigment is found in the anterior layers of the retina, its source is the layer of hexagonal pigmented cells which is now considered to form the deepest layer of that membrane. The most common cause of pigmentation of the retina is undoubtedly choroiditis, and although it must be considered an established fact that the pigment layer is histologically a part of the retina, yet clinically it is much more intimately associated with the choroid, participating, as it does, very readily in affections of this membrane,