REFERENCES

- 1 Moore, H. E., Watson-Jones, R., Bristow, W. R., and others (1933). *Lpool. med.-chir. J.*, **8**, 19.
- 2 Report of B.M.A. Fractures Committee, *British Medical Journal Supplement*, 1935, **1**, 53.
- 3 Kessler, H. H. (1935). *The Crippled and Disabled*, Columbia University Press.
- 4 Middleton, D. S. (1936). *Edinb. med. J.*, **43**, 641.
- 5 Rehabilitation and Industrial Injuries. Evidence of Trades Union Congress and B.M.A. *British Medical Journal Supplement*, 1937, **2**, 367.
- 6 Watson-Jones Rehabilitation Centre for Industrial Injuries, Liverpool, 1938, **6**, No. 2, April.
- 7 Cunningham, J. (1938). *Edinb. med. J.*, **45**, 712.
- 8 Griffiths, H. E. (1935). *Injury and Incapacity, with Special Reference to Industrial Injuries*.
- 9 Rehabilitation after Accidents. *British Medical Journal*, 1939, **2**, 417.
- 10 Final Report of Interdepartmental Committee on the Rehabilitation of Persons Injured by Accidents, 1939. H.M.S.O., London.
- 11 Civilian and War Injuries: Rehabilitation, Vocational Training, and Employment. Central Council Care of Cripples, 1940.
- 12 Hermann Levy War Effort and Industrial Injuries. Fabian Society, Series 253, July, 1940.
- 13 Nicoll, E. A. (1941). *British Medical Journal*, **1**, 501, 831.
- 14 Houlding, R. N. (1941). *Ibid.*, **2**, 429.
- 15 Malkin, S. A. S. (1941). Rehabilitation of Service Men. News Letter. Central Council Care of Cripples, October 10.
- 16 Medical Supervision of Industrial Workers, Report of B.M.A. Committee. *British Medical Journal*, 1941, **2**, 783.

A CASE OF STAPHYLOCOCCUS ALBUS SEPTICAEMIA

WITH REACTIONS OF 150 STRAINS OF STAPHYLOCOCCI

BY

MARY BARBER, M.D.Lond.

(From the Department of Pathology, British Postgraduate Medical School, Hammersmith Hospital)

It has long been realized that the pigment production of a particular strain of staphylococcus is not wholly reliable as a guide to its pathogenicity. Nevertheless, most laboratories are still content to report a staphylococcus as being albus, aureus, or citreus without further comment. The presence of a zone of haemolysis on a blood-agar plate has little or no significance. Soluble haemolysin production is fairly closely related to pathogenicity, but a reliable test entails growing the organism in an atmosphere containing about 20% carbon dioxide on a semi-solid medium and testing a filtrate of the resultant growth with rabbits' red cells. It is therefore too laborious to be used as a routine laboratory test. Loeb (1903-4) observed that certain strains of staphylococci were capable of clotting citrated plasma. This was also described by Much (1908), who emphasized the correlation of coagulase production and pathogenicity. More recently Cruickshank (1937) correlated coagulase production with fermentation reactions and haemolysin production, and found that the coagulase test was the one most closely related to pathogenicity for the rabbit. Chapman *et al.* (1934) compared haemolysin and coagulase tests as guides to pathogenicity in a large series of strains, and found that the coagulase-positive strains were usually pathogenic. Unfortunately their observations on haemolysis were confined to results obtained on blood-agar plates. In the following case of *Staph. albus* septicaemia no primary focus of infection was discovered, and the first positive finding was the blood culture. It therefore emphasized the need for some means of differentiating between a pathogenic *Staph. albus* and a saprophytic strain from the skin with which the blood culture might have been contaminated. In view of this I have thought it worth while to add to the description of this case a summary of the reactions of 150 strains of staphylococci derived from a variety of sources.

Case Report

The patient, a woman aged 52, was admitted to hospital on September 17, 1941, with a three-weeks history of febrile illness but no special symptoms. Six months previously her legs, up to the knees, had started to swell. The swelling persisted for five months and then passed off in a week, and on her admission was completely absent. On examination her temperature was

98.6° F., pulse 130, and respirations 22. She was flushed and sweating. The most conspicuous feature was skin lesions of an uncommon type. Nodules were present over the right deltoid, on the left upper arm, in the region of the left elbow, over the left hamstrings, in the right popliteal fossa, and on the buttocks. They ranged from 0.5 to 10 cm. in diameter, were dusky red to purple in colour, and a little warmer than the surrounding skin; some were slightly tender. They appeared to be involving the subcutaneous tissues. The teeth were carious, and there was gross pyorrhoea. The knee-jerks and ankle-jerks were absent; owing to the patient's mental confusion it was difficult to test for loss of sensation. Retention of urine was present, and this persisted. There was apparent neck rigidity, which at first suggested meningitis, but, as it later became clear, it was merely part of a general muscular tenderness and rigidity. No other significant physical signs were discovered. The urine contained albumin but was otherwise normal.

The patient ran a swinging temperature between 98° and 102° F. On September 18 a blood count showed: red cells, 5,000,000 per c.mm.; Hb, 80%; colour index, 0.8; white cells, 8,000—polymorphs 78%, lymphocytes 12%, monocytes 10%. The cerebrospinal fluid was under a pressure of 300 mm. but was otherwise normal. Blood cultures taken on September 18, 22, and 23 gave a pure growth of *Staph. albus*. On September 20 a biopsy of one of the cutaneous nodules showed it to consist of pyaemic abscesses of a somewhat chronic type, which contained colonies of staphylococci.

On September 20 oedema was observed over the sacrum, and there was ulceration of the cutaneous nodule over the right buttock. A blood count on the 25th showed that the haemoglobin had fallen to 66%, with 4,300,000 red cells and a colour index of 0.76; the white cells had risen to 24,000—neutrophils 89%, eosinophils 2%, lymphocytes 9%. On the 27th a sinus in the right buttock discharged a large quantity of pus. A culture of this gave a pure growth of *Staph. aureus*. The supuration in this area increased very much, and on the 30th the abscess was widely incised and explored. A large loculated abscess cavity full of pus was found in the subcutaneous tissues, but no evidence of a more deep-rooted lesion, such as osteomyelitis of the sacrum, was detected.

The patient was given a course of sulphathiazole, a total dose of 50 gm. being taken between September 19 and October 4; but her general condition steadily deteriorated, the abscess over the right buttock and sacrum increased in size, and on October 23 she died.

Post-mortem Findings

In the skin and subcutaneous tissues were found multiple abscesses, many of which were healed. A very extensive and foul-smelling abscess was present over the sacrum. It communicated with sinuses in the buttocks and with the pelvic soft tissues, which were inflamed. Thick yellow pus was seen in the extradural space, with collections round the cauda equina. There was no intrathecal inflammation or any evidence of disease in the sacrum or coccyx.

The lungs showed a few small abscesses on both sides and bronchopneumonic consolidation of both bases. The heart was rather small and there was a single circumscribed abscess in the myocardium, but otherwise no gross abnormalities. The kidneys had numerous abscesses in the parenchyma and there was severe bilateral pyelitis. The ureters were double on both sides, the left arising from a double and the right from a single pelvis. They united on each side before entering the bladder, which showed a severe purulent cystitis. The spleen was not enlarged, but appeared rather firm and fleshy. The genitalia were normal except for a chronic cervicitis. No abnormality was seen in the liver or the brain.

Histological section showed that the abscesses were of a rather chronic type, and there was little or no evidence of inflammatory reaction in the surrounding tissues. They contained clumps of staphylococci, but not in very large numbers. The spleen was not grossly abnormal, but the granulocytes were increased. There was a slight excess of polymorphonuclear cells in the sinuses of the liver.

Bacteriological Findings

A pure culture of *Staph. albus* was isolated on three occasions from the blood; *Staph. aureus* was grown from the abscess over

the sacrum; and from the lung abscesses post mortem an intermediate type was obtained.

Table I summarizes the reactions of the strains from the blood and the lung. The strain from the blood remained porcelain-white if grown on plain agar at 22° C. for a week or left at room temperature for many days, and even after passage through a rabbit. The strain from the lung produced a golden pigment when cultivated at 22° C. It will be seen that the strain from the blood, like that from the lung, was haemolytic to rabbits' red cells, was lethal to rabbits, and produced coagulase, so that the evidence of its being a pathogenic organism is fairly conclusive.

TABLE I.—Reactions of Strains from Blood and Lung

	Blood	Lung
Pigment production :		
Original culture at 37° C.	Albus	Albus
Later culture at 22° C.	"	Aureus
Coagulase production	+	+
α-lysin production	+	+
Pathogenicity to rabbit	+	+
Fermentation of { lactose	+	+
mannite	+	+
glucose	+	+
sucrose	+	+
Litmus milk	Acid (no clot)	Acid (no clot)
Liquefaction of gelatin	+	+
V.P.	+	+
M.R.	+	+

Alpha-lysin production was tested for by growing the organisms on 0.3% agar in an atmosphere containing 20% carbon dioxide for forty-four hours. The cultures were then centrifuged and the supernatant fluid was passed through a Seitz filter. The bacterium-free filtrate was set up in serial dilutions with equal volumes of a 2% suspension of rabbits' red cells in saline. The organism from the blood culture produced complete lysis in a dilution of 1 in 64, and that from the lung in a dilution of 1 in 32. The coagulase test was carried out as described by Fisk (1940).

Analysis of 150 Strains of Staphylococci from Various Sources

Table II summarizes the reactions of 150 strains of staphylococci, all isolated from material sent to this laboratory during the last few months; 68 produced coagulase

TABLE II

Group	Pigment Production		Fermentation of		Coagulase Production	No. of Strains
	Original Cultures at 37° C.	Later Cultures at 22° C.	Lactose	Mannite		
I	a Aureus	Aureus	+	+	+	43
	b " "	" "	+	+	+	3
	c Albus	" "	+	+	+	19
	d " "	Albus	+	+	+	3
II	a Albus	Albus	+	—	—	51
	b " "	" "	—	—	—	7
	c " "	" "	—	+	—	7
	d " "	" "	+	+	—	11
	e " "	Aureus	—	+	—	2
	f Aureus	" "	+	—	—	2
III	Citreus	Citreus	—	—	—	2

and 82 did not. Of the 68 coagulase-positive strains (Group I) 46 were typical aureus, and these 46 all came from obvious sources of infection such as osteomyelitis, boils, carbuncles, infected wounds, etc., except one which came from a nasal swab; 22 of the coagulase-positive strains were quite definitely porcelain-white on primary isolation at 37° C. even after leaving the plates at room temperature for one or two hours, but 19 of these produced an aureus pigment when grown at 22° C. or left at room temperature for several days. These 19 came from suppurative infections with two exceptions—one from a nasal swab and one from the vagina. Of the 3 true albus strains which produced coagulase, one came from the blood of the patient here reported; one was obviously an albus variant of an aureus strain as it was isolated from a septic finger, the culture from which produced a large number

of *Staph. aureus* colonies and a few albus; and the third came from the urine of a patient suffering from *Staph. aureus* septicaemia who had previously had *Staph. aureus* in the urine.

With regard to the 82 coagulase-negative strains 76 were typical albus (Group II) and 2 were citreus (Group III). Of these 78 strains 63 were quite definitely not causing any infections, and 14 of these apparently saprophytic organisms fermented mannite; 8 of the albus coagulase-negative strains were found in association with *Staph. aureus*, and one of these fermented mannite; 7 appeared to be the cause of a urinary infection and 3 of these fermented mannite. Two strains which did not produce coagulase appeared to be aureus after cultivation at 22° C. and fermented mannite (Group II). They both seemed to be the cause of a urinary infection. Two coagulase-negative strains were aureus on primary isolation. Neither fermented mannite and neither appeared to be the cause of any infection.

Discussion

This case of staphylococcal septicaemia is unusual in that no primary focus of infection was discovered and that the organism isolated from the blood was *Staph. albus*. On admission the only evidences of infection were the temperature and the cutaneous nodules. Later, obvious suppuration developed in the region of the sacrum and right buttock and there was a leucocytosis. Previously the white cell count had been normal. In view of the neurological signs it is possible that the collections of pus round the cauda equina were present on admission. The primary source of infection may have been a small area of osteomyelitis in the sacrum or coccyx, but no evidence of this was found on x-ray examination or at necropsy. The cutaneous nodules are interesting, and the biopsy of one of them was a valuable means of confirming the diagnosis. McCrea (1940) describes a case of septicaemia and pyaemia following a carbuncle in which "multiple painful red papules about the size of a shilling" developed over the right tibia and one at the insertion of the deltoid muscle. Mendell (1939), in an article dealing with 35 cases of staphylococcal septicaemia from various causes, described 10 cases which developed multiple metastatic abscesses in the skin; but I have not found a single recorded case in which, apart from pyrexia, cutaneous nodules were the sole evidence of infection. A man aged 48 was recently admitted to this hospital with *Staph. aureus* septicaemia, the primary focus probably being in the urinary tract. He developed a maculo-papular rash after admission, and the papules gradually increased in size to become slightly tender swellings in the skin and subcutaneous tissue. One of these was removed for histological section and showed a picture similar to that of the case described.

In the small series of staphylococci described not a single coagulase-negative strain was giving rise to infection outside the urinary tract, whereas 65 out of 68 coagulase-positive strains came from suppurative lesions, and 20 of these would have been reported as *Staph. albus* if only incubated at 37° C. All the pathogenic strains in this series fermented mannite, but many apparently saprophytic strains did too. The coagulase test is easily and quickly performed, and a four- to five-hour culture gives reliable results, so that if a single colony be subcultured into a tube of broth first thing in the morning, a few drops of the broth culture can be added to diluted plasma early the same afternoon. The test might therefore be used with advantage as a routine laboratory procedure.

Summary

A case of *Staph. albus* septicaemia in which no primary focus of infection was found is described.

A summary of the reactions of 150 strains of staphylococci is given, illustrating the value of the coagulase test as a means of differentiating between pathogenic and saprophytic strains.

My thanks are due to Prof. J. H. Dible and Dr. R. Cruickshank for advice and criticism; to Dr. P. Sharpey-Schafer, under whose care the patient was admitted to hospital; and to the Chief Medical Officer, London County Council, for permission to publish the case.

REFERENCES

- Chapman, G. H., Berens, C., Peters, A., and Curcio, L. (1934). *J. Bact.*, **28**, 343.
Cruickshank, R. (1937). *J. Path. Bact.*, **45**, 295.
Fisk, A. (1940). *Brit. J. exp. Path.*, **21**, 311.
Loeb, L. (1903-4). *J. med. Res.*, **10**, 407.
McCrea, H. M. (1940). *Proc. roy. Soc. Med.*, **33**, 679.
Mendell, T. H. (1939). *Arch. Intern. Med.*, **63**, 1068.
Much, H. (1908). *Biochem. Z.*, **14**, 143.

THE PRESENCE OF DIFFUSING FACTOR AMONG THE METABOLITES OF AEROBIC MICRO-ORGANISMS

BY

J. UNGAR, M.D.

AND

A. L. BACHARACH, M.A., F.I.C.*

(From the Bacteriological and Biochemical Departments,
Glaxo Laboratories Ltd.)

The presence of "diffusing factor" has been reported in: (i) mammalian testis extract (Hoffman and Duran-Reynals, 1931; McClean, 1930, 1931), (ii) snake and spider venom (Duran-Reynals, 1936), (iii) leech extract (Claude, 1937), (iv) malignant tumours (Boyland and McClean, 1935), as well as in cultures of the following micro-organisms: (v) staphylococcus (Duran-Reynals, 1933), (vi) streptococcus (Duran-Reynals, 1933), (vii) *Clostridium welchii* (McClean, 1936), (viii) *Cl. oedematis maligni* (*Vibrio septique*) (McClean, 1936), (ix) pneumococcus (McClean, 1936). Whether or not the diffusing activities evinced by these organisms and tissues are due to the same substance, or are even physiologically identical, cannot be stated with certainty, if only because the methods of test—to which we refer again below—have varied somewhat widely. We wish at the outset to make it clear that our references to "diffusing factor" and "diffusing activity" are to the substance that, in the simple but lucid definition of McClean and Morgan (1933), "dramatically increases the permeability of the skin to injected fluids and particles." All our tests, qualitative or quantitative, have been carried out by the procedure described elsewhere (Bacharach, Chance, and Middleton, 1940) for testicular diffusing factor (t.d.f.): the results of these tests are not necessarily the same as would have been obtained by different methods.

The presence of diffusing factor in bacterial cultures is of interest in itself to those investigating the metabolic processes of micro-organisms or applying the results of their work to medical practice, and has possible applications in therapy. Pharmacologically, diffusing factor appears to be inert: it has no effect on blood pressure (Christensen, 1938) and its action is not antagonized by histamine or adrenaline (Madinaveitia, 1939). Its injection into the cerebrospinal canal, however, markedly hastens the periaxial spread of cerebrospinal fluid (Cohen and Davies, 1939). Apart from the possibility, to which considerable attention has been given by McClean (1941c), that diffusing factor produced by micro-organisms might affect their pathogenicity, there is also the question whether non-toxic preparations of the active substance might not be used as

an adjunct in therapy—along with, for example, anaesthetics and narcotics, or in electrophoretic treatment—to reduce the quantities required for a given effect or the time required for an effect to be manifested.

Bound up with such considerations are the questions of the chemical nature and the specificity of diffusing factor. In some respects its properties are those of an enzyme system. Chain and Duthie (1939) found purified preparations of diffusing factor to possess marked mucolytic properties and suggested its identity with mucinase; but correlation between diffusing and mucolytic activity, whether this is determined by measuring the reduction in viscosity or in other ways, has not always been established. East, Madinaveitia, and Todd (1941) have recently shown that liberation of reducing substances does not go *pari passu* with reduction in viscosity, and suggest that two different enzymes may be involved: they have also demonstrated, by its action on β -phenyl-N-acetyl glucosaminide, the presence in testis extracts of an enzyme they call " β -glucosaminase," although it is admitted (Todd, personal communication) that " β -glucosaminidase" would be the more correct term. This enzyme is distinct from the substance (or complex) in the extract responsible for reducing the viscosity of hyaluronic acid. Robertson, Ropes, and Bauer (1940) examined preparations of *Cl. perfringens* and state that they hydrolysed mucoid (synovial fluid) in two stages: first there was a rapid decrease in viscosity, accompanied by a change in precipitability with aluminium hydroxide; then there was a gradual liberation of amino-sugars and reducing substances, reaching a maximum at twenty-four to twenty-eight hours. The work of Meyer and his colleagues (1941) suggests, on balance, a close correlation between diffusing activity and hyaluronidase content, as does the finding of McClean and Hale (1941) that the production of both hyaluronidase and diffusing activity by *Cl. welchii* or by streptococci is increased when these organisms are cultured in presence of hyaluronic acid. On the basis of existing knowledge it is not possible to say whether or not the increase in reducing power shown during the action of diffusing factor on mucoid is due wholly, or even in part, to the action of β -glucosaminidase. In any event, the relationship is not a simple one, and the matter must be considered still *sub judice*.

Unfortunately, lack of agreement as to the identity of diffusing factor even with a mucinase—or hyaluronidase, as most authors prefer to call it—and its differentiation from β -glucosaminidase have complicated the position, because different workers have been led to use different tests for the presence of diffusion-promoting substances. Apart from viscosimetric determinations and estimations of the amount of reducing substances liberated, a clot method has been described by McClean (1941b). In this the enzymic activity is measured by the extent to which, in the presence of acetic acid, the formation of a clot from mucoid is inhibited: there are also variants of the test on the skins of rabbits. Our method is based on the measurement of "blebs" produced in the skin of shaved rabbits after intradermal injection of the test solution. Full details have been published elsewhere (Bacharach, Chance, and Middleton, 1940), and it is here necessary only to indicate the quantitative limitations of the restricted tests reported below.

Between bleb areas of 4 and 7 sq. cm. the relationship between area and log dose is a linear one, these limiting values corresponding with about 0.4 and 2 of our units, respectively. Over this range the mean variance of an individual bleb was found to be 0.83. From this it follows that the means of three blebs differ from each other significantly at the 5% level when the difference exceeds 2 sq. cm.; with six blebs significance is established at the same level for differences of more than

* In this investigation Dr. Ungar had the assistance of N. McLeod and E. V. Welch; and Mr. Bacharach that of T. R. Middleton.