A PROSTHETIC ULCER OF THE MOUTH

BY

H. T. SIMMONS, M.D., Ch.M., F.R.C.S.

*Honorary Surgeon, Christie Hospital and Holt Radium
Institute, Manchester*

(WITH SPECIAL PLATE)

Benign tumours of the buccal cavity are uncommon. In two recent years 660 cases of disease of the mouth were examined at the Holt Radium Institute, and only 80 of them were non-malignant. Of these non-malignant cases 20 proved to be benign tumours and the remainder were inflammatory. Malignant growths therefore formed 97% of tumours in the buccal cavity, and 3% were innocent. These innocent tumours are important, however, because unless recognized they may receive unnecessarily drastic treatment, and if radiotherapy is used there will be little effect upon the growth.

One tumour, running true to type and possessing well-defined characteristics, has occurred often enough to merit description, especially as it has been invariably suspected of malignancy. Current surgical textbooks and literature do not mention this swelling, but a search through dental writings has revealed a description of a case by Thoma (1934).

The swelling lies in the sulcus between the cheek or lip and the gum, and is situated anteriorly to one or other side of the midline. It is much more common in the lower than in the upper jaw, and in women than in men. A series of ridges are formed which run parallel to the line of the alveolus and are slightly deeper in colour than the surrounding mucosa (Plate, Figs. 1 and 2). There are always two or more ridges, about an eighth of an inch in thickness and as much as a quarter of an inch in height, and after running for an inch or more along the gum they gradually slope away into the normal tissues. The ridges are covered with mucous membrane, but in the furrows ulceration exists in the deeper parts. On palpation the tumour is surprisingly hard, and this undoubtedly gives rise to the suspicion of malignancy. Very little movement on the deeper tissues can be detected because the swelling is anchored to the alveolus by the innermost furrow. There is little tenderness except during attacks of inflammation, which occur at infrequent intervals. It is only during these attacks that pain is experienced, and the main complaint is inconvenience, with a feeling that "this swelling ought to be attended to because it might lead to cancer." The patient is past middle life, and the gum is always edentulous. Advanced atrophy of the alveolus makes the denture very loose-fitting, and it will slip into any one of the furrows without inconvenience to the patient. The denture is an old one and has never been remodelled. The atrophy of the edentulous alveolus and the wearing of the loose-fitting denture are the cause of this variety of prosthetic ulcer, the "hit-or-miss" fit explaining the number of the ridges.

On microscopical examination the epithelial surface is irregular, but, although it extends fairly deeply into the underlying stroma, it is everywhere limited by a definite membrana propria. The underlying stroma shows scattered blood sinuses with occasional foci of lymphoid cells indicative of chronic inflammation.

The diagnosis from carcinoma is not difficult, but the patient usually presents himself when there is an acute attack of inflammation. The tumour is then reddened and angry-looking, with ulceration between the ridges. He has discarded his denture, and therefore the mechanical cause of the condition is not at once evident. The extreme

hardness of the swelling and the induration of the base due to the ulceration in the furrows still further suggest the diagnosis of malignancy. On the other hand, the growth has been present many months—often one or two years. The hypertrophied portions are not friable, and bleeding does not occur. The ulceration is confined to the deepest portions of the furrows, and if the denture is obtained it will be found to fit accurately in any furrow. Finally there is no involvement of the regional lymphatic glands.

I have had 12 cases of this condition under my care in the past seven years, 9 of the patients being females and 3 males. Their ages ranged from 41 to 75 years, the mean being towards the end of the sixth decade. All were completely edentulous, and the original false teeth with which they had been supplied had been worn from five to twenty-six years. The atrophy of the alveoli was very advanced, the denture fitting loosely and slipping about whilst the patient was talking. Normally the lower denture in an edentulous mouth is much less secure than the upper, and this additional factor of instability would explain why in 10 cases the lower jaw was affected and in only 2 the upper. The tumours had been present from two months to five years, with the majority of patients seeking advice from six months to two years after noticing a swelling. In 2 cases the onset was painful owing to the pressure of the denture, but pain was infrequent and never severe. None of the patients suffered from gingivitis or stomatitis, nor was there any evidence of previous syphilitic infection.

The histories of the two cases illustrated are as follows:

Case 1.—A shopkeeper aged 62 had worn the same dentures for fifteen years. Two years before attending the hospital he developed a painful swelling "like a gumboil" on the inner aspect of the left lower lip. The swelling slowly increased in size, and was painless except when attacks of inflammation occurred. It had always been hard; there was no discharge and no bleeding. The patient was a teetotaler, did not smoke, and the Wassermann reaction was negative. On examination there was a typical prosthetic ulcer in the left lower alveolar sulcus 2.5 by 1.5 cm. in size (Plate, Figs. 1 and 2). The lower denture was very loose-fitting and slipped in and out of the furrows of the ulcer. There were no palpable lymphatic glands. The ulcer was excised with the diathermy loop and microscopical examination confirmed the clinical diagnosis. The wound healed in six weeks and the wearing of the remodelled dentures was resumed six months afterwards. It is now seven years since the operation, and there has been no recurrence.

Case 2.—A housewife aged 45 had had all her teeth extracted nine years previously and the dentures she then obtained had not been altered. Twelve months before coming to hospital she noticed a small hard painless swelling on the inner side of the left upper lip (Fig. 3). This swelling had doubled in size in the twelvemonth and was painless except when the teeth were pressed into it. There was no discharge or bleeding. She did not smoke or take any form of alcohol, and there was no suggestion of syphilis. On examination a typical prosthetic ulcer was found in the left upper alveolar sulcus 2.5 cm. in length. The upper denture was loose and slipped down whilst the patient was talking. There was no enlargement of the neighbouring lymphatic glands. The ulcer was removed by diathermy in November, 1934, and the wound healed in just over a month. Histological examination confirmed the clinical diagnosis and there has been no recurrence. New dentures were obtained and worn six months after operation.

The best treatment proved to be excision of the ridges with the diathermy loop, and the base of the tumour was very lightly coagulated with the diathermy button (evipan anaesthesia was used). Healing took place in about a month, with very little scarring, the only treatment

required after operation being bland mouth washes. All the patients were warned before operation that they would require a new denture or remodelling of the old one, and wearing of the teeth has been deferred until six months after the healing of the wound. One case had been previously treated elsewhere as malignant disease, and received an efficient implant of radium element giving a dosage of 6,000 r. It was interesting to find that the tumour had proved completely resistant, although a certain amount of troublesome necrosis had occurred in surrounding normal tissue.

Summary

A type of benign prosthetic ulcer of the mouth is described.

It is suggested that its relative frequency, trueness to type, and well-defined characteristics warrant its recognition as a definite clinical entity.

The causation, pathology, and treatment are discussed.

REFERENCE

Thoma, K. H. (1934). *Clinical Pathology of the Jaws*, Baillière, Tindall and Cox, London.

Clinical Memoranda

Irreducible Intussusception in a Baby, treated by Resection: Recovery

(WITH SPECIAL PLATE)

The problem of irreducible intussusception has evoked a variety of surgical methods, all more or less unsuccessful. In small children the mortality is very high. Perrin and Lindsay (1921) in a series of 400 consecutive cases of intussusception in the London Hospital give these mortality figures for irreducible cases: all cases treated by resection, 70%; cases under 3 years old treated by resection, 100%; cases treated by other methods, 100%. The conclusion one draws is that resection, though very unlikely to succeed, is the only method worth attempting. The following case was successfully treated by resection in Addenbrooke's Hospital in May, 1940.

CASE RECORD

The patient was a boy of 18 months, previously quite well apart from bronchitis. He was admitted with a history of abdominal pain, severe vomiting, and absolute constipation for four days in spite of enemas. Castor oil had been given at the onset. He had been passing small quantities of blood for two days. On admission he was dehydrated, the abdomen was distended, with a step-ladder pattern on it, and a mass could be felt in the right iliac fossa. The pulse was 160.

Operation was carried out under ether anaesthesia. An ileo-colic intussusception was found, beginning about twelve inches above the ileo-caecal valve and extending to the hepatic flexure. As only a few inches of this could be reduced and the part was gangrenous and was considered quite unviable the whole mass was resected between clamps. Anastomosis was done between the side wall of the ileum above the closed end and the open end of the colon a short distance below the hepatic flexure. The child's condition was not alarming at the end of this operation.

Intravenous saline and glucose were given continuously for six days, the child's leg being immobilized in plaster for this purpose. A nasal tube was passed and the stomach contents were removed. The fluid withdrawn was at first offensive and dark brown, and it remained so for three days. On the second day he passed an offensive dark-coloured stool. On the third day the abdomen was less distended and he had a mild attack of diarrhoea. On the fourth day he was taking half an ounce of fluid hourly by the mouth. Convalescence was complicated by a mild attack of measles, with bronchitis and otitis media, but he was in good health when he left the hospital four weeks after the operation.

Anterior and posterior views of the resected piece of bowel are shown in Figs. 1 and 2 of the Plate.

H. T. LAYCOCK, M.A., M.B., F.R.C.S.,
Resident Surgical Officer, Addenbrooke's
Hospital, Cambridge.

REFERENCE

Perrin, W. S., and Lindsay, E. C. (1921). *Brit. J. Surg.*, 9, 46.

Multiple Foetal Abnormalities

(WITH SPECIAL PLATE)

In view of the unusual and extensive type of abnormalities in the case described, I consider that it merits record. Careful search in the literature through the good offices of the librarian of the B.M.A. has failed to reveal a similar case with such extensive deformity.

CASE REPORT

Mrs. —, aged 33, has an only child of 4 years, who is apparently normal in every way. Since this child was born she has had a normal menstrual history, and used no contraceptives. She can recall no personal accident or untoward incident during the second pregnancy such as might give rise to a monster.

The confinement, at a maternity home, was a breech presentation with extended legs. No foetal heart sounds were heard. Under general anaesthesia a foot was brought down, the breech delivered, and the first abnormality displayed itself as a hernia of the liver covered by peritoneum through the abdominal wall (see Plate). There was a deficiency in the chest wall, involving the third, fourth, and fifth ribs, through which the heart and pericardial sac protruded. Finally careful delivery of the head showed a condition of hernia cerebri. The vertex of the foetal skull was absent and the free skull border appeared continuous and apparently fused with the amniotic membrane, which enclosed the brain substance. The brain was normal on section. Beyond this was the placenta, firmly attached. The face showed double harelip and cleft palate. The foetal skin was clean and intact and of normal appearance. The foetus showed no sign of life. There was at least fourteen to eighteen pints of liquor amnii.

Ante-natal care had shown a consistently normal urine and blood pressure, with breech presentation, but this was not interfered with.

I beg to thank Dr. Burn, medical officer of health, Barnsley, for the photograph and for his general interest in the case; also Dr. Rose.

Barnsley.

A. B. SLACK, M.B., Ch.B.

A Case of Unilateral Twin Ectopic Pregnancy

(WITH SPECIAL PLATE)

Unilateral twin ectopic pregnancy is a comparatively rare condition. Falk and Blinick (1938), having reviewed the literature and described two cases of their own, state that 65 authentic cases had been reported. Anderson (1939) describes a case of his own.

CASE REPORT

A woman aged 24 was admitted to the North Riding Infirmary, Middlesbrough, on August 20, 1940, with the following history. She vomited in the morning and had pain in both sides of the lower abdomen which sometimes passed round to the back. This pain was severe and persistent, and on admission rose in the abdomen almost to the costal margin. There was a history of two months' amenorrhoea. She had had two children, the youngest being 18 months old.

On examination the patient was markedly exsanguinated, cold, and sweating, and the pulse was rapid and weak. There was a slight blood-stained discharge per vaginam. The diagnosis of a ruptured tubal pregnancy was made and the patient was taken to the theatre immediately. Under general anaesthesia the abdomen was opened and the diagnosis confirmed. The peritoneal cavity was filled with blood; the right Fallopian tube was ruptured and the bag of membranes was found intact. Right salpingectomy was performed. The ovaries and left Fallopian tube were normal. An autotransfusion of blood was carried out. When the unruptured sac was opened clear

DOUGLAS HUBBLE AND G. R. OSBORN: ACUTE BRONCHIOLITIS IN CHILDREN

FIG. 1.—A bronchiole 0.7 mm. in diameter. Complete destruction of lining epithelium and plug of almost pure mucus, though there are no glands in the wall. General emphysema and localized alveolar changes well shown.

FIG. 2.—Respiratory bronchiole blocked by purulent plug. Peribronchiolitis, interstitial alveolar changes, and general emphysema.

FIG. 3.—Small bronchus blocked by a mucous plug. Damage to lining epithelium and active secretion by mucous glands. Mucicarmine stain.

A. B. SLACK: MULTIPLE FOETAL ABNORMALITIES

Showing herniation of liver, heart, and brain, and cleft palate and bilateral harelip. Placenta adherent to brain.

WILLIAM C. SMITH: UNILATERAL TWIN ECTOPIC
PREGNANCY

H. T. LAYCOCK: IRREDUCIBLE INTUSSUSCEPTION
IN A BABY, TREATED BY RESECTION

FIG. 1.—Anterior view.

FIG. 2.—Posterior view.

H. T. SIMMONS: PROSTHETIC ULCER OF THE
MOUTH

FIG. 1.—Prosthetic ulcer of mouth. Note atrophy of alveolus.

FIG. 2.—Prosthetic ulcer of mouth. With denture in situ.

FIG. 3.—Prosthetic ulcer of mouth. Upper lip type.