was not haemophilia. Conditions which are strictly comparable to this phase of the experiment occur spontaneously in man, and we hope to discuss them on another occasion.

Discussion

These data are presented in the first place to show that a venom-lecithin reagent gives results in the accelerated clotting test which are in all respects comparable to those of thromboplastins such as brain extract; and in the second place to give figures for normal values and range of variation. The most potent accelerator of coagulation which we have used is a mixture of Russell-viper venom with 10% emulsions of brain. When this is added to normal plasma, clotting occurs in 5 seconds after recalcification, which is faster than with any other reagent recorded in the literature. Such a reagent demands for its study instrumental methods such as Nygaard's, which are unfortunately hard to procure at the present time. Without such instrumental aids excessive potency of the clot-accelerating reagent may, as we have shown, be a source of error. We hold no brief for Quick's test in any of its modifications as a means of estimating prothrombin. It is affected not only by the rate of conversion of prothrombin into thrombin but also by the presence of inhibitory or accelerator principles. It is nevertheless a valuable clinical test when due heed is paid to possible fallacies. Russell-viper venom and lecithin are relatively cheap and stable reagents for the test. They allow us to decompose the action of thromboplastin into its water-soluble and lipid-soluble moieties and thus gain more insight into disturbances of coagulation than is possible with the more complex reagent. They have the additional advantage that they are effective in widely different animal species. Results with dicoumarin suggest that this is a point of great importance, and we shall deal with it in a subsequent paper.

Summary

Quick's prothrombin test is based on the hypothesis that if the coagulation of plasma is accelerated by excess of thromboplastin the speed of coagulation will depend mainly on the amount of prothrombin present. Other things being equal, the speed of coagulation is inversely proportional to the concentration of prothrombin.

When reagents composed of mixtures of Russell-viper venom and crude lecithin are used as thromboplastins the coagulation time of recalcified plasma can be reduced to 5 seconds. Unless instrumental methods of recording are used, such reagents are too swift for accurate work.

The venom-lecithin reagent has been standardized to give an average coagulation time of 10 seconds. The normal range of variation has been determined, and methods of scoring the results in terms of prothrombin are discussed.

We are indebted to Dr. R. H. Mole for much help in standardizing the technique of the test, and to Dr. R. B. Fisher for explaining the mathematical treatment of the data.

REFERENCES

- Butt, H. R., Allen, E. V., and Bollman, J. L. (1941). *Proc. Mayo Clin.*, **16**, 388.
 Ferguson, J. H. (1939). *Proc. Soc. exp. Biol.*, N.Y., **42**, 33.
 Fullerton, H. W. (1940). *Lancet*, **2**, 195.
 — (1941). *British Medical Journal*, **1**, 98.
 Hobson, F. C. G., and Witts, L. J. (1941). *J. Path. Bact.*, **52**, 367.
 Holmboe, H. S., and Holmboe, R. W. (1940). *J. Lab. clin. Med.*, **26**, 408.
 Illingworth, C. F. W. (1939). *Edinb. med. J.*, **46**, 762.
 Jaques, L. B. (1941). *J. Physiol.*, **100**, 275.
 Lozner, E. L., Kark, R., and Taylor, F. H. L. (1939). *J. clin. Invest.*, **18**, 603.
 Macfarlane, R. G., and Mole, R. H. (1942). Personal communication.
 Nygaard, K. K. (1941). *Hemorrhagic Diseases*, St. Louis.
 Owen, T. K., and Toohey, M. (1941). *Lancet*, **1**, 724.
 Pohle, F. J., and Stewart, J. K. (1939). *Amer. J. med. Sci.*, **198**, 622.
 Quick, A. J. (1936). *Amer. J. Physiol.*, **115**, 317.
 — (1937). *Ibid.*, **118**, 260.
 — (1939). *Pennsylvania med. J.*, **43**, 125.
 Witts, L. J., and Hobson, F. C. G. (1940). *British Medical Journal*, **2**, 862.
 — (1941). *Ibid.*, **1**, 170.

PERINEURITIS

BY

JAMES CYRIAX, M.D.

Assistant Medical Officer, Physiotherapeutic Department,
St. Thomas's Hospital

In the diagnostic sphere the territories of the neurologist and the physiotherapist overlap at one point—in the elucidation of some of the diffuse pains so often spoken of by the patient as "neuritis." Many, probably most, patients complaining of "neuritis" are suffering from referred pain clearly demonstrable as taking origin from a deeply situated somatic tissue. Other patients with this complaint are found to have some disease of the nervous system. Evidence has, however, been accumulating that cases occur in which, despite the presence of unimpaired conduction along a nerve, the pain originates in a nerve. Such pain is not the lancinating stab of neuralgia but a continuous burning ache closely resembling that caused by severe inflammation of such structures as muscle, tendon, joint capsule, etc. In these cases the following criteria are satisfied: (1) no pain can be elicited from the muscles, joints, etc., comprising the same group of segments as the painful part; (2) conduction along the nervous system is normal; (3) the pain is referred to a region obviously corresponding accurately to the area of cutaneous supply of a peripheral nerve. The conclusion is then inescapable that the pain arises in a nerve but not in its conducting elements. In other words, it is the nerve sheath that is inflamed—perineuritis.

Theoretical Considerations

Since the inflamed structure in perineuritis is a fibrous sheath, the quality of the pain resembles that of fibrositis of muscles and similar tissues. It is in its mode of reference that it differs: for not only does the region in which the pain is felt correspond with a known cutaneous area, but the patient can tell whether the front or the back, say, of the finger is affected. In fibrositis of other structures the pain is usually felt right inside the part. Though at first sight it may appear unlikely, symptoms due to perineuritis are often brought on by using the painful member. This can be demonstrated by a simple experiment. The subject sits on the edge of a chair in the manner known to cause pressure on the sciatic nerve in the upper thigh. When paraesthesiae can be felt in the foot, the subject, keeping otherwise perfectly still, voluntarily flexes his toes. This at once increases the pain, though the movement in no way alters the tension on the affected part of the nerve trunk. Hence, though the cause of this phenomenon is obscure, a complaint of pain brought on or aggravated by exercise does not preclude perineuritis.

Secondary parenchymatous change may occur in perineuritis, presumably because of the pressure exerted by the swollen sheath on the nerve fibres. In sciatic perineuritis this occurred in 10 cases out of 18, and showed itself by wasting of the gluteal muscles and by incomplete ability to contract them voluntarily, wasting of the hamstrings, and loss of the ankle-jerk. When these neurological signs appear the label "sciatic neuritis" would seem warranted, but this name leads to an error in emphasis, focusing attention upon the secondary rather than the primary organic process.

In these figures of proved cases of sciatic perineuritis lies the answer to the school of thought that would confine the use of the term "sciatica" to sciatic neuritis. Since roughly half of all cases of sciatic perineuritis show no evidence of loss of conduction along the nerve, this limitation results in the indefensible position in which some, but not other, cases of sciatic perineuritis are called "sciatica." Moreover, a few patients with perineuritis complain only of backache: this raises the question whether that symptom, if of sciatic periradicular origin, can be called sciatica. The alternative, and to my mind better, course is to retain the word sciatica—well known to the public—as the name of a common symptom, like headache. A medical definition compatible with this use of the word would then be: "A

Sixty per cent. of India's medical stores are now made in India as compared with 25% at the beginning of the war. Instrument manufacturers have expanded their works and increased their production, and only such instruments as cystoscopes, microscopes, etc., need be imported. A number of medicinal items have recently been taken off the list of importations, also certain bacteriological stains which are now prepared by the distillation of ordinary commercial and crude dyes. The rubber industry has widened its range of products so that it is now possible to obtain a large percentage of the appliances necessary in hospital.

diffuse pain characterizing some lesions situated in the proximal parts of the fourth lumbar to second sacral segments." For the layman the definition would be: "A pain felt to run down the lower limb from above."

Confirmation by Local Anaesthesia.—All diagnoses of perineuritis are speculative, and no case can be considered such until confirmation by local anaesthesia has been carried out successfully. The affected part of the nerve sheath is infiltrated with a procaine solution, whereupon the pain disappears at once for the duration of anaesthesia. Only cases in which this test showed the diagnosis to be correct are cited here. Direct injection about the nerve sheath was employed in each instance except that of the sciatic nerve, whose roots were anaesthetized epidurally via the sacral hiatus.

Types of Perineuritis.—Perineuritis may be set up by friction; this variety is well recognized. It is the perineuritis due to inflammation of unknown origin, and therefore called "rheumatic," that causes difficulty in diagnosis.

Friction Perineuritis

Little need be said about this familiar variety. Whereas prolonged or severe pressure on a nerve results in traumatic palsy, intermittent pressure—i.e., friction—sets up perineuritis at first. The common cause of this type of irritation of a nerve sheath is pressure against a skeletal projection. This may be a cervical rib, a protruded lumbar intervertebral disk, an osteoma (e.g., of the head of the fibula), or a cubitus valgus deformity. The lateral cutaneous nerve of the thigh may be pressed upon by the edge of the foramen by which it pierces the fascia lata (meralgia). Patients may unwittingly subject a nerve to intermittent pressure of postural origin (e.g., by sitting cross-legged or by keeping the elbow in full flexion). Tumours may stretch or invade a nerve, but do not fall within the scope of the physiotherapist. If perineuritis due to friction is not relieved, progress to a true neuritis in course of time is certain.

Rheumatic Perineuritis

This group comprises those cases in which no cause for the inflammation is detectable. The nerves most often affected are the facial and the sciatic, but ulnar, median, musculospiral, and external popliteal perineuritis also occur.

Facial Perineuritis.—The swelling of the nerve sheath in its bony canal leads to a secondary pressure palsy and thus to paralysis of the facial musculature. Since the nerve is purely motor, premonitory pain is absent.

Rheumatic Perineuritis in the Upper Limb

Ulnar Perineuritis.—The pain is felt chiefly at the ulnar border of the hand and in the little, ring, and middle fingers. There may be some discomfort at the elbow. In 2 particularly severe cases the pain was felt to extend from the base of the neck to the fingers; yet the whole symptom was completely abolished by infiltration with a procaine solution of a stretch of nerve sheath at the elbow one inch long, no peripheral anaesthesia being thus produced. Previous examination by a neurologist had shown conduction along the nerve to be normal. The sheath was inflamed at the elbow in all the cases that I have seen. Postures involving prolonged flexion at the elbow (e.g., lying supine with the hands under the occiput) may provoke perineuritis here, as may the altered stress on the nerve caused by malunion of a condylar fracture of the humerus. Recurrent dislocation of the ulnar nerve often leads at first to perineuritis; in addition to pain, the patient then complains of something going "out and in" at the elbow. In other cases no reason can be found.

Median Perineuritis.—The chief pain is in the thumb, but the fact that it also extends to the palmar surface of the two adjacent fingers should immediately arouse suspicion. In a case of 4 years' duration two operations had been performed on the thumb in the vain search for a foreign body. Contrary to expectation simultaneous extension of the wrist and elbow failed to evoke the pain. Search was made along the course of the median nerve in the forearm for a spot pressure on which set up the pain in the thumb. In 3 proved cases these spots were found at the wrist, and 2 inches and 5 inches above it, respectively.

Musculospiral Perineuritis.—Of this condition only 2 examples have come my way. There was severe and constant pain stretching from the base of the neck to the radial side of the hand. The only localizing sign was a marked paresis of the triceps muscle. This finding led to an examination for tenderness along the course of the musculospiral trunk in the arm; it was found where the nerve winds round the shaft of the humerus, and local anaesthesia confirmed this tentative diagnosis.

Brachial Neuritis

The preceding condition is one of the few in my experience that merit the label "brachial neuritis"; for in addition to the pain originating in the nerve sheath there was evidence of loss of conduction. Even so, a diagnosis of musculospiral perineuritis would be more accurately descriptive. I have seen 3 other cases in which pain arose, apparently spontaneously, at the root of the neck and in the arm; a week later paresis of the deltoid, supraspinatus, and biceps muscles was apparent. To these cases the label "brachial neuritis" might also be applied, but again the localization of the lesion to the fifth and sixth cervical roots suggests that "brachial" is an unnecessarily wide term. These cases are mentioned in order to suggest that increased accuracy in diagnosis may render "brachial neuritis" a term of very rare and restricted application.

Rheumatic Perineuritis in the Lower Limb

When perineuritis affects the upper limb, stretching its nerves does not always evoke the pain. On the other hand, stretching the nerve regularly gives rise to pain in the two perineuritic conditions familiar to me as affecting the lower limb. These are sciatic and external popliteal perineuritis, for each of which a pathognomonic test exists.

External Popliteal Perineuritis.—The pain is usually felt at the outer side of the shin and foot, but in severe cases it may extend from the region of the greater trochanter to the toes. The test for the nerve is as follows. While the knee is kept extended the foot is fully dorsiflexed: this causes no pain. The foot is then fully inverted: this hurts. While the position of the foot is maintained the knee is allowed to flex a little. If the pain disappears the nerve is at fault. In every case the sheath proved to be affected where it winds about the head and neck of the fibula.

Sciatic Perineuritis.—The history is of unilateral pain first felt in the lower back, then spreading to the buttock, thigh, leg, and foot. Exceptionally, only a complaint of backache is made. In the acute stage the pain is continuous and severe; in the subacute stage there is a constant ache aggravated by all movements that stretch the nerve (e.g., standing trunk-flexion); in the chronic stage there is pain only on such movements. An asymmetrical posture designed to take all possible tension off the nerve roots is adopted in all but mild cases; sciatic perineuritis is a common cause of "sciatic scoliosis." Secondary parenchymatous change occurs in about half the cases; this shows itself by wasting of the gluteal and hamstring muscles, and by loss of the ankle-jerk. This reflex, once thus lost, does not return even after the patient has been well for some years. Taken alone, loss of the ankle-jerk, even on the affected side only, is not much to go by; for not only may it be present in sciatic perineuritis, but it may be unaccountably absent in cases in which the pain can be shown not to originate in a nerve.

Head-and-Knee Test for Sciatic Perineuritis.—This is carried out as follows: The patient sits; his trunk and thigh remain motionless throughout. The leg is passively extended at the knee until the pain just starts. It is then allowed to flex a very short distance until the pain just goes. The patient now drops his chin on to his chest. This brings the pain on again. The knee is then allowed to flex a little more. Disappearance of the pain shows the sciatic nerve to be at fault. The principle behind this test is that a tension created by a flexion movement of the neck is released by a flexion movement at the knee. This would be possible only for a structure continuous from neck to leg: there exists but one such—the spinal cord and its continuation, the sciatic nerve. Pain on straight-leg raising, which stretches every posterior structure in the thigh, buttock, and lumbar region, does not indicate that the lesion necessarily lies in the sciatic nerve.

Signs Suggesting Frictional Sciatic Perineuritis

At times it is possible to be fairly certain when the patient is first seen that protrusion of a lower lumbar intervertebral disk has occurred. Clearly, the factors set out below suggest this *only when they supplement the signs of sciatic perineuritis*.)

1. *The Length of History.*—Non-frictional sciatic perineuritis usually gets well of itself in 8 to 12 months. Hence a history lasting for over a year is suggestive.

2. *Signs of Nervous Disease in the Leg.*—When to the signs of sciatic perineuritis are added weakness of the dorsiflexor muscles of the foot, particularly of the extensor longus hallucis, and/or impairment of sensation in the skin of the outer side of the leg or foot, the presence of a protruded disk is highly probable. Absence of ankle-jerk, gluteal wasting, and postural deformity are common to both idiopathic and frictional perineuritis and thus throw no light on the problem. So far I have come across one case in which pressure from a protruded disk had led to a marked paresis of the dorsiflexor muscles in the leg without affecting the amplitude of the ankle-jerk, although the time interval between the tap and the response was considerably increased.

3. *Great Pain on Epidural Infiltration.*—This sign was pointed out by Love in 1936. If the pain is such that the patient has to ask for the discontinuance of the injection, the presence of a spinal block is likely. The converse does not hold; for patients found later to have a protrusion have tolerated the injection well.

4. *Failure to Improve after Epidural Infiltration.*—Whether idiopathic or frictional in origin, the pain of sciatic perineuritis and the range of straight-leg raising improve markedly during the hour or so after the injection. Occasionally the result is dramatic, and the standing patient may bend and touch his toes painlessly after having been unable to reach beyond his knees for months. If there is no immediate improvement the diagnosis of sciatic perineuritis is probably wrong. If when seen a week later he is no better, it is my practice to repeat the injection. If when seen at the end of another week the improvement is again found to have been transitory, the inference is drawn that a protrusion exists.

5. *Sudden Disappearance of the Whole Syndrome.*—Patients suffering from protrusion of a disk sometimes give a history of having had and having fully recovered from the same condition in the past. Spontaneous reduction of a protruded disk may occur, therefore. One such case—as yet unproved—has come my way. The patient's history was of five months' severe sciatica, which disappeared overnight the day before he attended. Full straight-leg raising caused no discomfort, but unilateral absence of the ankle-jerk and gluteal wasting were evident. In my experience untreated idiopathic sciatic perineuritis has always got better very slowly; hence reduction of a protruded disk is the probable explanation.

Signs of Perineuritis

A summary of the facts outlined above shows that the signs fall into five groups, but are seldom all present in any one case. It is noticeable that the nerve sheath is nearly always affected where it lies in close contact with bone. This raises the question whether there is not some slight frictional element in even "rheumatic" perineuritis.

Pain on Stretching the Nerve.—This is the outstanding sign when perineuritis affects the lower limb. The whole brachial plexus can be stretched by abducting the arm to 60 degrees and applying traction to the hand while the neck is bent fully to the opposite side. Pain can usually be elicited in brachial perineuritis by this means, but the vertebro-scapular muscles are also tightened thereby; they too are not uncommon sources of pain felt to radiate along the whole upper limb. Thus no certain test can be devised for the upper limb.

Tenderness and Swelling of the Nerve Sheath.—Both these signs are presumably present in every case. Except for the sciatic nerve roots, tenderness of the nerve trunk is readily found, and pressure at this point reproduces the distant pain. Swelling of the nerve has been palpable only of the ulnar nerve at the humeral condyle.

Postural Deformity.—In sciatic perineuritis that is at all severe, lateral curvature of the lumbar spine associated with little or no rotation develops from the adoption of the posture best calculated to take all tension off the nerve. Together with this, the ability to bend the trunk forwards from the upright position all but disappears. Cubitus valgus, whether developmental or the result of malunion of a fracture near the elbow, is often associated with ulnar perineuritis. Gross osteophyte

formation may cause pressure on the external popliteal nerve at the knee.

Evidence of Secondary Parenchymatous Change.—Perineuritis may progress to neuritis if the swelling of the nerve sheath exerts pressure on the conducting fibres. In the case of the facial nerve it is only when palsy supervenes that the condition becomes apparent at all. Correct attribution of the early symptoms to perineuritis may be assumed when signs of neuritis appear later and are diagnostic of pressure on a part of the peripheral nervous system.

Cessation of Symptoms for the Duration of Local Anaesthesia.—This is the only sure method of confirming what can only be a more or less tentative diagnosis. If this is correct, and the anaesthetic solution reaches the nerve sheath, relief is achieved within a minute or two, and lasts for about an hour and a half.

Treatment of Perineuritis

First employed as a method of corroborating a diagnosis, local anaesthesia was soon found by experience to be an excellent method of treatment. When the pain due to the return of perception to the nerve sheath had worn off, marked symptomatic improvement was reported in almost every case. Two, or at the most three, infiltrations spaced at weekly or fortnightly intervals have afforded full relief. So reliable is this treatment that if primary cessation of pain is not followed by secondary alleviation the presence of an unsuspected frictional cause is suggested.

Direct action of the procaine on the nerve sheath appears to be the way relief is obtained. Alternatively, distension with fluid of the space between the nerve and the tissues round it might be responsible. If this were so, the effect would presumably be obtained by separation of adhesions about the sheath by hydraulic pressure. In the case of the sciatic nerve this object would also be attained by stretching under general anaesthesia; but this procedure makes the patient much worse in acute or subacute sciatic perineuritis. Since, then, mobilization of the nerve is not responsible, the view that procaine initiates a lasting effect on nerve sheaths is the more probable hypothesis. In my experience no treatment other than the induction of local anaesthesia of the nerve sheath is effective. Heat has proved valueless and massage harmful.

Illustrative Case

A girl cut the inner side of her left elbow at the age of 10; the wound healed and was forgotten. At the age of 20 pain began to be felt in the inner side of the hand, and anterior transposition of the ulnar nerve was performed. Relief lasted 4 years; the symptoms then returned, getting gradually worse until severe continuous pain extended from the base of the neck to the ulnar fingers. Infiltration of the transposed ulnar nerve sheath at the elbow led on several occasions to complete relief, lasting as long as a fortnight. Then, however, the pain returned and was soon as bad as ever. The conclusion reached was that recurrence was due to the nerve's being bound down mechanically, and open mobilization finally became necessary. At the operation the ulnar nerve was found bound down by a thick scar to the front of the medial humeral condyle, and by the next day all the pain had gone. There can be no doubt that the pain arose in the ulnar nerve, yet at no time did investigation in the neurological department reveal any loss of conduction, in spite of the severity of the symptoms and the length of history. I believe that it is to this type of case, when the perineuritis is acute enough to cause pain referred both up and down the limb, that the term "brachial neuritis" or even "causalgia" is sometimes applied, and that it is then an unnecessarily wide term.

G. Wiberg (*Nord. Med.*, 1941, 10, 1929), who records his observations on twenty-three cases of de Quervain's tenosynovitis stenosis, discusses the possibility of a connexion between trauma and this condition. In individual cases such a connexion seems to be forthcoming, but a clear proof of a former injury to the wrist-joint is required in the form of damage to the tendons of the extensor pollicis brevis or abductor pollicis longus as occurs in an open wound, or the discovery at operation of a definite sign of an old tendon lesion, or a scar corresponding to the tendons of extensor pollicis brevis or abductor pollicis longus in their course over the styloid process of the radius.