BRITISH MEDICAL JOURNAL

LONDON SATURDAY MAY 9 1942

MEDICINE AND THE PUBLIC WELFARE*

BY

VISCOUNT DAWSON OF PENN, M.D.

When the war, as now, is passing through an anxious phase we are apt to say, "A plague on reconstruction talk—first win the war." This is anxiety entangling the mind in one-track thinking. On the contrary, we should at times occupy our minds with fancies free and especially with the future, in order to re-create our minds for their war duties; for in so doing we indirectly assure ourselves there will be a future. And here, in passing, I may remind you that the best psychological treatment is often that which is implicit rather than explicit.

War does not so much produce social changes as hasten the fruition of those already in seed or bud, or again bring back to the light of day projects born years previously but later lost in the mists of inertia or political exigency. For 25 years or more the idea has been spreading that the practice of medicine should be occupied not only with those disabled by sickness or accident but also, if not primarily, with the building up of health, and that for this extended purpose our medical services have become too haphazard and incoordinate in their arrangement. In other words, organization has not kept pace with ideas. If the community is to have in its care the health of its citizens as it has their education, the location of health centres and hospitals must, like that of schools, correspond to the distribution and needs of the population. Indeed, it is surely apparent that the rearing and training of a healthy body and the education of its occupant are two processes so interdependent that a system of health centres is as necessary as a system of schools, and when it is recalled that the integration of health is in many of its aspects a matter of education the parallelism becomes closer.

Health Services in Retrospect

A brief retrospect will clarify this contention. Although the care of public health came into being many years before public elementary education was instituted in 1870, it was in those early years, for the most part, impersonal and environmental, and concerned itself with such matters as drainage, sewerage, disposal of refuse, pure water supplies, air, and light. It was not until 1915 that there opened up the chapter of communal care for the health of the human body in the shape of special services for expectant mothers, maternity, infants, children in schools, dental care, and for certain diseases, of which tuberculosis was the forerunner (1912).

All credit to the founders of the communal health services; but greater fame would have been theirs had they secured the active participation of the whole profession. Instead they reared these services in bureaucratic isolationism. The responsibility lay in equal measure with the Royal Colleges and the teaching hospitals and their staffs; their minds were held within the ancient confines of curative medicine and were blind to the enlarging opportunities opening out before them. The result was a gulf fixed between preventive and curative medicine, which has now to be bridged. In these health clinics preventive and curative medicine meet, and who can doubt that general practice should embrace both and that hospital practice and the medical curriculum should include both?

Immediately after the war of 1914-18 the Ministry of Health was founded, and in its constitution there was provision for the

setting up of a "Consultative Council on Medical and Allied Services." This Council was intended as a link between the profession and the Ministry, and I was appointed its chairman. Its first reference was: "To consider and make recommendations as to the scheme or schemes requisite for the systematized provision of such forms of medical and allied services as should, in the opinion of the Council, be available for the inhabitants of a given area." Here was the notion of the region.

A Basis for Reconstruction

In 1920 there followed a unanimous report the recommendations of which might well form a basis of reconstruction to-day. Indeed, the movement of 1920-1 has been reborn and extended under the name of regionalization, which is being furthered with both skill and purpose by the Nuffield Provincial Hospitals Trust and by the Planning Commission of the British Medical Association.

The next landmark was the Local Government Act of 1929, which among its provisions included the setting up of rate-supported municipal and county hospitals, which are now making such progress in our midst.

Another trend, more and more apparent during the last 25 years, is the need for greater institutional provision. The increase of knowledge, and with that the development of teamwork, has made this inevitable. The doctor works best where equipment and ancillary services are available, and side by side with his fellows. Further, such institutional provision is economical of both time and cost and makes a health service available for all citizens. In short, the tendency is for the centre of gravity of practice to shift away from the home to organized surgeries, clinics, health centres, and hospitals, and these would be distributed in their various grades throughout the region so as to constitute a co-ordinated scheme of service—some deriving from voluntary and others from rate-supported institutions. The way in which the war has made clear the importance of industrial medicine illustrates this theme.

In the realms of both health and education there is organized provision for the welfare of the young from birth to the present school-leaving age. After this but little is ordered for either education or health. Yet youth in their teens and onwards are entering workshops, factories, and mines, and, let us hope, agriculture, and in these formative years their success or failure in life may well be determined.

An Ordered Service of Medicine

Here is opportunity for health centres attached to industrial establishments to concern themselves with healthy living. They will get to know physiques and aptitudes and their relation to working conditions; they will note defects of frame and tendencies to ill-health while in their early and curable stages, and provide corrective treatment for them; they will send cases of illness to their own or suitable hospitals, and on the positive side will direct physical training, encourage games, superintend the dietary of the canteens, and, above all, carry on the health education of the workers, impressing upon them that in the last resort their health depends upon themselves. To take the example of tuberculosis, it is not too much to say that, through the agency of such health centres, which would include examination of workers by x rays, the elimination of tuberculosis would in a few years make giant strides. Here would be medicine at

* Speech at a meeting of the Medical Society of a Military Hospital.

its best—prevention and cure hand in hand, scientifically and socially ordered and new knowledge, ever accruing. And here progress can be achieved during the war: no mass institution of health centres, but a few well-conceived experiments which would take cognizance of services already existing and prepare the way for future developments.

Is it not plain that for any national planning scheme to be successful it is necessary that voluntary hospitals and agencies on the one hand and rate-supported hospitals and local authorities on the other hand should work together in double harness with the guidance but not direct governance of the State, and that personal health services and curative medicine must be brought into close co-operation in general practice, in hospitals, health centres, and clinics, and, not least, in education.

Both the rate-supported and the voluntary hospitals have their values. Both have the welfare of the community at heart and are necessary one to the other. The former, though still needing adjustments and improvements, have made remarkable progress in the few years since their inception, and often show readiness to co-operate with the voluntary hospitals, notably in the County of London. For their highest posts clinical work is unfortunately at a disadvantage compared with administration. Their chief disadvantage is that they are not yet in close enough contact with the collective mind of academic medicine, which is illustrated by the defective methods for the appointment of their visiting staffs. On the other hand, the leading voluntary hospitals have centuries of tradition behind them; they have been the homes of the outstanding minds and thinkers in medicine, and to-day hold leaders of the profession. Their faults are that they think too much in terms of their individual selves and fail to appreciate they are members of a group. If in this changing world they were to yield some of their ingrained detachment and enter into this new fellowship they would make an incomparable contribution to the welfare of the community.

An Evolutionary Plan

If I have your agreement that an ordered service of medicine (just as an ordered service of education) is necessary for the welfare of this country, I submit that such service must be something which suits the habits of thought of our people and should evolve from what already exists—and it may be stated with confidence that what already exists is, speaking generally, very good.

General practice, the basis of any scheme of medical services, will require institutional provision at health centres and selected hospitals. In the 1920 scheme hospitals were grouped into primary, secondary, special, and teaching. Primary hospitals were to be staffed by general practitioners aided by visiting consultants, and the remaining groups by consultants and specialists.

Under regionalization such a scheme of graded hospitals (some hospitals rate-supported, others voluntary) and of personal health services would be implemented to meet the needs of the region and to eliminate overlapping and unnecessary units. A teaching hospital linked to a university would, wherever possible, be the focus or the directing centre of the region, and there would appear to be a general agreement that dividing the country into regions, although it may involve the combining of local government areas, for this purpose would ensure the best service.

If, then, we assume for the moment the country divided into regions, and a regional council for the ordering of each region, the question arises, What should be the powers of the regional council; should they be advisory or executive? Whether a regional council has executive or only advisory powers, the overriding desideratum is that the medical services should be guided by those who have knowledge and purpose and are not unduly influenced by political-mindedness. In either case a regional council will be faced by the difficulty of relations with multiple local authorities and of cutting across their boundaries. Again, in either case it will contain on the one hand representatives of voluntary medical and allied services, and on the other hand nominees of local authorities and probably a few co-opted members.

If it is to have executive powers, these powers would either be overriding and need to be established by statute or they would be delegated by local authorities themselves. The advantages

would be correlation of services by unity of command for the whole region, and a body possessed of the requisite knowledge and devoted with singleness of purpose to personal health and hospital services. But such a regional council would need to be supplied with funds and possessed of an administrative and financial establishment, which could not fail to rub shoulders with the major local authorities within the region, involving contacts so close as to be charged with the danger of friction. Thus might be imperilled that confidence and co-operation without which the regional council could not attain success.

On the whole I suggest it would be better in the first instance—on a five-year plan if you like—for the final executive power to be with the major local authorities through statutory committees (on the lines of the education committees), with an obligation placed on these committees to seek the advice of the regional council, which would have the right, in the event of its advice being rejected, of appealing to the central authority in Whitehall. If in its turn the regional council were assisted by a medical committee representative of the region on which medical officers of health had seats, I suggest it would as an advisory body have more weight than if it were itself possessed of full executive functions.

There is yet another, and bolder, plan worthy of consideration, which would be an advance in political science. Government to be efficient must be guided by scientific knowledge. Autocracies avail themselves of that knowledge with greater readiness because the dictator sets up the appropriate technical body and without delay adopts or refuses its advice, whereas democracies dilly-dally because so many departments that do not understand the technical problems must have a say. Innumerable examples come to mind—synthetic rubber; the synthesis of glycerin; which would have saved our fats, bread, milk, vegetable protein; protective measures against the spread of diphtheria and venereal disease—to mention a few examples at random.

To meet this difficulty democracy is setting up semi-autonomous executive bodies on which the Government is strongly represented and otherwise consisting of those who have the requisite knowledge. Radio-activity, broadcasting, electricity are examples of the subjects dealt with. Why should not health and hospitals be under such semi-autonomous bodies? In the 12 or more regions they would be based on the major local authorities, and in London there could be a central semi-autonomous authority with power delegated from the Ministry of Health. It is, I submit, worth thinking about.

Payment and Staffing of Services

Next the question of payment for services received. It is surely right in principle that patients should make a reasonable payment either directly or through contributory schemes or other forms of insurance. Here I submit that though contributory schemes are sound in idea and practice they are likely some day to evolve into a nation-wide scheme of comprehensive social insurance, and the planning of services would make this practicable. The assessment of payments by patients and of remuneration to staffs of hospitals would need to be uniform as between State-supported and voluntary institutions in the same region.

Let us pass to consider the staffing of these services in the homes, surgeries, health centres, and hospitals. Would the conversion of medical practice into a whole-time service be in the national interest? With the prospect of large social and economic changes following the war the answer might in the course of years be in the affirmative. Fortunately no answer is needed now. If agreement on planning is reached, it can only be implemented step by step, for a full pattern of services will not spring up in the night. The first stage of implementation will be the co-ordination of existing hospital services and the bringing of personal health services into the body of medical practice and the various people concerned in this implementation learning to work together.

These are the most urgent needs. Health centres and clinics will usefully appear here and there, but on no large scale until experience of their varying designs and functions has been gathered. Gradualness is inevitable.

With the payment of visiting staffs of hospitals, with the staffing of health services by suitably qualified general practitioners, with the upgrowth of industrial medicine, and the habit

of some local authorities to favour whole-time service, not to mention the proposed extension of panel practice, it is clear that the earnings of the profession will be derived in increasing proportion from salary and in decreasing proportion from fees.

So it should be with the staffing of hospitals. Apart from the professorial units, the larger hospitals will doubtless offer increasing opportunities to men who desire to undertake whole-time hospital service. The London County Council affords a good example of whole-time and part-time men working side by side.

Freedom and Elasticity

Five years will show us the way. We must innovate greatly but quietly. Meanwhile we can pause to consider a whole-time medical civil service on its merits. It would be a more compact and tidy administration, and would probably be run at less cost; but would it have the same value? We have an example of a whole-time service in the R.A.M.C. For war its organization is admirable, but for civil life would not its administration be too rigid, advancement too dependent on seniority, and its day-to-day work without a sharp enough spur of ambition? It is important not to confuse ambition with egoism. Ambition stimulates a man to do his best, and a healthy thought of self consorts well with a desire and endeavour to help others. A whole-time altruism is a futile philosophy.

There are few callings in which there is so big a gap between routine and the best work as in medicine, and no profession needs to be so elastic in its government if it is to be dynamic, not static. Its front line of knowledge is always mobile and often advancing; the men working there need freedom for initiative and should be unfettered by the formulae of administration, and, may be, on the other hand, by the demands of practice. And the same considerations hold in the sphere of clinical practice, for though we need ordered planning we must strive to avoid any cast-iron uniformity. Rather do we seek unity amidst diversity; for man, whatever his political colour, is individual, and in illness even more so than in health, and his doctor must remain an individual and not become an official. It is here that medical planning is up against its crucial difficulty, in that it requires collectivism for its fabric and individualism for human relationships. And individualism will not flourish easily within the rigid boundaries of a State service, but needs the freer atmosphere which belongs to the voluntary hospitals' tradition; for in the difficult conditions of a greatly changing world doctors will need not only knowledge but understanding if they are to guide bodies and minds along the straight road of health and content.

ESTIMATION OF PROTHROMBIN WITH VENOM AND LECITHIN

BY

L. J. WITTS, M.D., F.R.C.P.

Nuffield Professor of Clinical Medicine

AND

FELICITY C. G. HOBSON, B.A.

(From the Radcliffe Infirmary, Oxford)

In earlier papers (Witts and Hobson, 1940, 1941; Hobson and Witts, 1941) we described the use of solutions of Russell-viper venom and lecithin in place of the tissue extracts commonly used in the estimation of the prothrombin time. A brief account of our present technique and the normal range of variation may be of interest to workers in this field.

Technical Details

Glassware.—Tubes are checked with a gauge to ensure that they are all of the same internal diameter. Glassware must be scrupulously clean; tubes are left in potassium bichromate and sulphuric acid cleaning solution overnight and then washed in running water for at least 4 hours.

Lecithin.—This is ovolecithin B.P.C. as purchased. Specimens are variable and not all give such good effects as we have described. It is therefore wise to try another sample if first

results are disappointing. The active principle is not known, but it is probably an impurity or a breakdown product rather than pure lecithin. The strength of the stock solution of lecithin in alcohol should not be less than 10% to avoid an alcohol effect when the lecithin is added to the venom. As lecithin solutions become acid on standing the alcoholic solution of lecithin should be adjusted to a pH between 7 and 8 by means of a saturated solution of caustic soda in 95% alcohol; the reaction should be checked at fortnightly intervals. The optimal concentration of lecithin is 0.5%—i.e., 0.05 c.cm. of 10% alcoholic solution of lecithin in 1 c.cm. of the actual dilution of venom which is used.

Calcium.—We use M/40 CaCl₂ solution, which is the same as for Quick's test.

Temperature of Water-bath.—The optimal temperature is from 37° to 43° C. As prothrombin is very thermolabile, it is wise to work at the lower end of this range. Our observations have been carried out at 38.5° C.

Plasma.—Nine c.cm. of blood, withdrawn rapidly and with special precautions to avoid trauma, is promptly and thoroughly mixed with 1 c.cm. of M/10 potassium oxalate and centrifuged at 1,500 r.p.m. for 5 minutes. The duration and speed of centrifugalization should be maintained constant. If there is obvious haemolysis the plasma should be discarded, more particularly if the lecithin difference is to be determined. As oxalated plasma is not stable for much longer than an hour at room temperature the oxalated blood or plasma should be put into the refrigerator as soon as possible after venepuncture and removed only immediately before the estimation of the clotting time. It seems immaterial whether the blood is centrifuged soon after venepuncture or just before the estimation. In the refrigerator the plasma will keep its clotting time unchanged for at least 24 hours.

Estimation.—A dry clean test-tube (50 by 8 mm.) is placed in the water-bath and 0.1 c.cm. of plasma and 0.1 c.cm. of the venom-lecithin reagent are added. Mix and leave to attain the temperature of the water-bath; the right time is about 1½ to 2 minutes. Then add 0.1 c.cm. of the calcium solution, which is kept in the incubator alongside the water-bath, and start the stop-watch. The end-point is different when venom alone and venom-lecithin are used. In both cases the tube is held at a slant of 45 to 60 degrees and rotated round its longitudinal axis. The end-point with venom alone is the moment when fibrils of fibrin first appear. If the same criterion is used when the venom-lecithin reagent is employed the end-point is much more difficult to detect, because the opalescence obscures the fibrin fibrils, as Fullerton (1940) has pointed out. As the tube is slowly rotated, however, a fine film of opalescent liquid quite suddenly appears on the inside surface of the tube above the surface of the mixture. This film is actually coagulated plasma, and the end-point is the moment when it is first seen. It is a very sharp end-point, and is best seen by diffuse light. The tube should not be rotated round its horizontal transverse axis or shaken, because these manœuvres accelerate the end-point in a variable manner.

Lecithin Difference.—The lecithin difference is the difference in seconds between the clotting time with venom alone and with the venom-lecithin reagent.

Normal Values and Range of Variations

We have previously shown that whereas different batches of Russell-viper venom may vary in strength, samples from the same batch are constant. In the present investigation we have estimated the accelerated clotting time of the plasma from 56 normal men and women (medical students, laboratory workers, and nurses), using venom from a single batch. The venom was used in two dilutions—1 in 10,000, which is the maximum potency, and 1 in 80,000, which had seemed in our earlier experiments to provide a more convenient speed at which to work. Quick's test was also carried out on these same plasmas. For this we used two different preparations of dried rabbit brain. The first was prepared for us by Messrs. Boots Pure Drug Co., Ltd., who mixed 10 g. of dried rabbit brain so that it was homogeneous and then packed it in evacuated ampoules, 0.1 g. per ampoule. The second preparation was made in our own laboratory as required, and represents several batches. The results are given in Table I.