THE PROBLEM OF ABDOMINAL PAIN WITH SPECIAL REFERENCE TO THE LOCALIZATION OF VISCERAL PAIN

BY

FRANCIS R. BROWN, O.B.E., F.R.C.S.

Surgeon to Dundee Royal Infirmary

What is the problem? Why have so many theories been formulated? Morley (1931) has put the matter succinctly:

"The need for a theory of abdominal pain arose from the discovery that the gastro-intestinal tract and the other intra-peritoneal organs, when exposed to direct mechanical, chemical, or thermal stimulation, are incapable of producing painful sensations. This fact, when once realized, had to be reconciled with the universal experience that pain does result from various inflammatory or obstructive disorders of these organs. A further and more obvious difficulty was found in the peculiar distribution of the pain in many abdominal diseases. Sometimes it would be central, sometimes unilateral. In some cases it would be felt chiefly in the back, and occasionally it was most marked in so remote a region as the supra-acromial area of the shoulder. An explanation had to be found for this peculiar radiation of pain to regions more or less remote from the causal lesion. Further, in certain disorders of the abdominal viscera, it was found that the pain is associated with localized areas of tenderness on pressure, cutaneous hyperalgesia, and muscular rigidity, all in the abdominal wall, while in other and equally painful disorders these superficial manifestations are absent. A theory had to be elaborated that would explain all these highly confusing manifestations of abdominal pain."

In the present somewhat discursive paper there will be found much that is old and true, a good deal that is more recent, mainly speculative, but of great interest, and a little that is new. It is my hope that in the "new" something of interest may be found and perhaps even a grain or two of truth.

"Where do you feel the pain?" The purist may reply, "In my sensorium," but the ordinary patient clearly understands one's meaning, and will indicate, more or less exactly, where his pain is situated.

Before approaching the problem of abdominal pain it is of course essential to have a clear conception of the development, anatomy, and physiology of the tissues concerned, as well as of the morbid and pathological conditions which may excite the sensation of pain. Especially must we know the peculiar nerve pathways present and to what particular stimuli they respond. We know that each special sensory apparatus or system is so made or modified that it responds only to its own peculiar stimulus. Thus the visual apparatus responds only to light, the auditory system only to sound, and so on. The stimulus must be "adequate"—i.e., of the right kind and of sufficient strength—before it reaches the threshold of consciousness: there it is analysed and registered.

Visceral Pain

The abdominal viscera have no cerebrospinal (somatic) nerve supply, their innervation being by autonomic nerves only. That is why we can cut, pinch, or burn the intestine without producing pain. Why, then, does true visceral pain occur? Hurst (1911) has proved that increased tension in the muscular wall of the gut is the appropriate stimulus for the production of this pain. The cause of this increased tension is hyperperistalsis such as occurs in biliary colic, intestinal obstruction,

the pains of labour, and acute appendicular obstruction. In these conditions the exaggerated contractions—and the resulting pains—are intermittent. When the contraction is more prolonged, as in spasm of the stomach, the pain is more continuous—e.g., hunger pain of duodenal ulcer. Whether the pain impulse is excited by pressure on the afferent nerve endings or by ischaemia or by liberation of the pain-producing substance we do not know. Musser (1904), Nothnagel (1905), and Neumann (1910) all recognized that increased tension in the gut could cause visceral pain, but it was Hurst who proved conclusively that this is the only mechanism. It is often said that distension can cause visceral pain. It generally does, since distension stimulates contraction, but it is well known that in paralytic ileus and in the terminal stage of unrelieved intestinal obstruction no peristaltic action occurs. When the bowel, from fatigue, gives up the struggle, peristalsis and pain cease—"the silent abdomen"—but increasing distension goes on until the inevitable end. A similar sequence of events takes place in the atonic and over-distended urinary bladder. Thus, before visceral pain can occur, distension must be overcome or at least be resisted by active muscular contraction.

Morley (1931) has given us, chiefly through his observations and experiments on "shoulder-tip pain," his theory of peritoneo-cutaneous radiation and peritoneo-muscular reflex. This theory is an attractive one and of the greatest practical help as a working hypothesis. It has to a great extent supplanted Mackenzie's theory of a viscerocutaneous reflex and a visceromotor reflex. Mackenzie rejected Ross's splanchnic pain entirely, and explained all visceral pain as referred by a system of "reflexes" through the splanchnic afferent nerves to the cerebrospinal nerves of the abdominal parietes. With the aid of local and spinal anaesthetics it is now easy to prove Mackenzie's theory wrong and to show that Morley's conviction is right.

Morley writes: "I am convinced that true visceral pain exists . . . it is in no sense referred to the superficial structures of the abdominal wall, and is a deep-seated central pain, not accurately localized."

Some years ago an old and very thin woman was admitted to Dundee Royal Infirmary under my care with a diagnosis of acute obstruction due to a strangulated umbilical hernia. There was no doubt about the acute obstruction—complete constipation, vomiting, distension of the abdomen, and intermittent attacks of severe visceral pain in the umbilical region all occurring. On examination we found a large umbilical hernia, the diameter of the hernial opening being about 3 inches, but there was neither obstruction nor strangulation at this site, the hernial protrusion being very easily reducible. It was interesting and instructive to see and feel through the very thin coverings of the hernia successive waves of hyperperistalsis of the small intestine—each wave of tetanic contraction exactly coinciding with a bout of severe central pain. As each wave of peristalsis subsided so did the umbilical pain. No hyperaesthesia or hyperalgesia or muscular rigidity was found in the abdominal wall. Further examination revealed the presence of a small right-sided femoral hernia, of which the patient was completely unaware since it was producing no local spontaneous pain. Pressure on the femoral hernia elicited pain. This is an observation that is not always fully appreciated—i.e., that local tenderness, in somatic tissues, can be found, and should be sought for, before the onset of spontaneous pain; the simple

explanation being, of course, that pressure increases the pain-producing stimulus. I then, for the benefit of a confirmed "Mackenzieite," completely anaesthetized the skin and abdominal parietes in front of and for a considerable area around the umbilical hernia. This, as I anticipated, had no appreciable effect on the frequency or intensity of the visceral pain; thus proving—if further proof be necessary—that visceral pain is not referred to and "felt" in the abdominal parietes. At operation a small loop of strangulated small intestine was found in the femoral hernia sac as well as a little blood-stained fluid, the latter being a chemical rather than a physical irritant to the parietal peritoneum.

Practically all observations and experiments on visceral pain have been made on an anatomical and/or physiological basis in the fully developed subject, generally with due consideration of morbid and pathological states and operation findings. For a true understanding of this "deep-seated central pain" we

occupy the umbilical region, and the colon is situated in the hypogastrium. If these organs permanently occupied these respective positions we should naturally expect visceral pain produced in the stomach, duodenum, and gall-bladder to be situated in the epigastrium, and likewise small-intestine colic and appendicular colic to be localized in the umbilical region, and colon pain in the hypogastrium—exactly in the positions that clinical experience has taught us these pains are localized after full development of the intestine. (See Fig. 5.)

But the intestine does not remain a short straight tube: it elongates enormously, rotates, herniates into the umbilical cord, returns to the abdomen, and continues to lengthen. Finally it assumes its permanent position as shown in Fig. 4.

What happens to its nerve supply during this period of elongation, rotation, and migration? The nerves lengthen too, of course, but they are still the original nerves, and no new nerve

FIG. 1a

FIG. 1b

FIG. 2

FIG. 3

FIG. 4

FIG. 5

- FIG. 1a.—Showing alimentary tract in ventral view, a simple straight tube, at 4 weeks. Compare with Fig. 5.
 FIG. 1b.—Alimentary tract at 5 weeks, viewed from left side. Shows dorsal mesentery and formation of loop—caecum indicated.
 FIG. 2.—Rotation of intestinal loop round the superior mesenteric artery—caecum undescended. 6 weeks.
 FIG. 3.—Herniation of intestinal loops into cord. 7 weeks. Rotation not shown.
 FIG. 4.—The final disposition of the intestines. 10 weeks.
 FIG. 5.—Showing the localization of true visceral pain produced in the intestine. Compare with Fig. 1a. A, Epigastric region; B, Umbilical region; C, Hypogastric region.

must go far deeper. We must seek the help of embryology and morphology; we must go back to the study of the embryo before anatomical structure was completed or physiological functions were established, and follow its development step by step.

Figs. 1a, 1b, 2, 3, and 4 show diagrammatically various stages in the development of the gastro-intestinal tract in the human embryo. Figs. 1a and 1b show the alimentary tract as a simple straight tube, slung by a dorsal mesentery, lying in the midline of the body; nerve supply from fifth to twelfth thoracic ganglia through the three splanchnic nerves—a bilateral innervation from the autonomic nervous system. The gut has, in the abdomen, no somatic nerve supply. This is what one might term the original position of the intestine, and it is of the greatest importance for our consideration and understanding of the localization of visceral pain to keep this primary relative position ever in mind. It is clear from Fig. 1a that the stomach and the duodenum, with its diverticulum the gall-bladder, lie in the epigastrium, the small intestine and appendix

supply is acquired by the intestine during its growth and migration. And what of sensation? Nature has wisely decreed that the less we know, in health, about our "insides" the better, and so the intestine and peritoneum have neither tactile nor positional sensation. Nor have we any subjective knowledge of normal physiological processes, such as digestion and normal intestinal peristalsis. Life would be very unpleasant—indeed impossible—if we were continually conscious of our liver, hobnailed or smooth, rubbing up and down against our parietal peritoneum, or if we were aware of normal peristaltic action. For a similar reason the cornea has no tactile sensitivity. Yet it is common experience that certain abdominal diseases and disorders do give rise to pain, often of an intense character, and thus we are warned of impending danger—danger which may threaten health or life itself.

What conditions in a hollow tube are likely to endanger health or life? The presence of a harmful irritant or of something which may obstruct the lumen of the tube: one must be got rid of and the other overcome as soon as possible. For

that purpose peristaltic contractions become more powerful than normal, and if needs are desperate the muscle wall can quickly hypertrophy so that greater and greater force may be exerted. In case of necessity peristaltic action can even go into "reverse gear." When this hyperperistalsis occurs, then—as I have already described—the sensation of pain (visceral pain) is produced.

Where Visceral Pain is felt

Where will the pain be "felt"? Always, if the nerve afferents are purely autonomic, in the primary relative position of the affected organ. If we must have a word to describe this peculiar localization of visceral pain I suggest the term "retrojection" (to coin a word) but no reflex action is involved. We know that sight is *projected*, and that surgeons daily project touch by means of a probe. Growth and migration of the gut and its nerves have made no difference to sensation or sensitivity, and therefore pain produced, say, in the appendix—no matter where it lies—will be "felt" in the umbilical region, biliary colic will be "felt" in the epigastrium although the gall-bladder has migrated to the right hypochondrium, and so on. Thus we should know fairly accurately, by the position of the pain (visceral), which organ is affected. Incidentally, it cannot be too strongly emphasized that the most important question one can ask a patient suffering from abdominal pain is this: "Where did the pain first begin?" The answer to this question will in many cases indicate which organ is affected, and so will enable us to differentiate clearly such conditions as acute cholecystitis (early visceral pain in the epigastrium), acute appendicular obstruction (visceral pain in the umbilical region), renal colic (pain in the lumbar region—always unilateral, never central), rupture of a Fallopian-tube pregnancy (primary pain is "somatic" and always begins low in one or other side of the pelvis). Pain produced in a bilateral organ is always unilateral, because the nerve supply is unilateral.

And suppose the bowel perforates? We may not be able to tell which particular segment of the gut is affected, but the parietal peritoneum will soon indicate where to look for the trouble, and in time experience should enable us to make more and more accurate diagnoses. The parietal peritoneum is exquisitely sensitive to such irritants as inflammatory and infective exudates, bowel contents, extravasated bile, and even sterile blood. Such irritants stimulate the somatic pain-conducting afferents of the parietal peritoneum and abdominal parietes; and, clinically, local tenderness and muscular rigidity, and local pain, indicate when and where these irritants are present.

How, it may be asked, can the peritoneum, with neither tactile nor positional sense, give us this valuable information? The parietal peritoneum, or rather its inner serous surface, is endowed with a special sense, and one only—namely, sensitivity to a chemical stimulus (again like the cornea)—and thus the pain impulse is excited by the presence of any foreign chemical substance in contact with it.

Many of those interested in and puzzled by this problem have attempted by countless observations and ingenious experiments to find a satisfactory solution. Many theories have been formulated and propounded. They speak of visceral pain as being "radiated" or "referred," "reflected," "transferred," "deflected," etc., by means of what they call "reflexes" from the affected viscus to some distant tissue. May I remind them of this: where a particular sensation is essential for the welfare and survival of the organism it will be provided; if it is unnecessary or undesirable it will not be provided.

My view is that the completely developed intestine, long and rotated though it be, and however far parts of it have migrated, is still essentially a midline organ; its nerve supply is that of a short straight tube; its sensitivity is still that of a short straight tube situated in the middle of the abdomen; and it naturally and inevitably follows that pain produced in a part of that tube, whether short or long, twisted or straight, will be localized in the primary relative position of the affected part. (Figs. 1 and 5.)

Aristotle's Experiment

Let us now consider Aristotle's well-known experiment and see what significance it bears on the subject.

In his *Problemata* Aristotle (384-322 B.C.) asks this question: "Why is it that an object which is held between two crossed fingers appears to be two? Is it because we touch it at two

sentient points? For when we hold the hand in its natural position we cannot touch an object with the outer sides of the two fingers." From three separate sources I find that the *Problemata* of Aristotle in which the experiment is mentioned is not, strictly speaking, Aristotle's own work. It was the compilation of several hands, and belongs to the time after the Christian era—perhaps the fifth century. It would seem that nothing more than the question was asked, and no answer, except surmises, was given; and, besides, it may not possibly have been posited by Aristotle but by some later writer. It is clear that Aristotle knew nothing of the functions of the nerves and nervous system and so could not answer his question. However, for our purpose, that matters little. What does matter is the significance of this simple experiment quoted from *Starling's Principles of Human Physiology* (1941):

"If with the eyes shut we cross the first and middle fingers and bring them in this position in contact with a pea, we should at once say that two peas lay under the fingers. This is especially marked if the pea be rolled between the fingers. The two sides of the fingers which come in contact with the pea usually touch two different objects, and these parts of the skin would have to be re-educated—i.e., their local sign would have to be changed in accordance with the changed conditions—before the pea would be perceived in its true state as single."

I have repeated the experiment—other objects, such as the blunt end of a pencil, can be used instead of a pea—and of course found that the illusion, as such it is, does occur. The converse, as one would expect, also occurs—that is, two objects, one in contact with the now distal aspect of the index finger and the other in contact with the now distal aspect of the middle finger, are felt as one. I realized that the "two" objects in the original experiment were separated by a definite interval—two fingerbreadths: that was suggestive. Then, with eyes closed and forgetting that my fingers were crossed, I concentrated on the relative positions of the "two" objects (or points of contact) and found that the "object" in contact with my index finger undoubtedly lay—so far as my consciousness could discriminate—on the radial side of the "other object" in contact with the middle finger: in other words, I now believed my fingers were no longer crossed but were parallel—i.e., in their normal relation or primary relative positions. If pain is produced by pressure or by putting sharp projections on the object held between the crossed fingers the illusion becomes even more pronounced. I believe that the sensations of touch and pain produced in the crossed fingers are "retrojected" to the primary relative positions of the parts stimulated, and in my opinion this supplies the logical answer to the two-thousand-year-old question. Physiologists may say that the explanation is "the conditioned state of the cerebral cortex," but such an explanation could not apply to the localization of visceral pain, since we have neither consciousness nor memory of the position of our abdominal organs (unless impressions were registered very, very early in embryonic life?).

Tastevin (1937) writes: "The inversion of the sensations produced in two crossed fingers has been observed by several others, and, for the first time, by Czermak (1855), who at the same time drew attention to an analogous phenomenon in displacement of the lips." Tastevin, in an elaboration of Aristotle's experiment, found that these false localizations, or illusions, occur in any tissues mobile enough to be moved from their natural positions—e.g., the lips, the tongue, the fingers, the eyes, fractured limbs, etc. In a search of the literature I have found nothing to suggest that anyone has previously considered that the illusions produced by these experiments are similar to the illusion produced by pain excited in an organ which has been displaced from its primary relative position by normal processes of development and growth. It is my contention, and conviction, that the two conditions, artificial and developmental, are completely analogous, provided the migrated organ (or tissue) retains *only* its original afferent nerve supply. Thus, and thus only, can we explain the peculiar localization of visceral pain when situated at a distance from the affected organ. Here, indeed, is the key to the problem.

One suspects that some of those interested in this problem believe that if they knew "precisely" or "accurately" which tissues to anaesthetize in the painful area they could by induction of block anaesthesia prevent or abolish this so-called

"referred pain." They will never succeed: they might as well try to anaesthetize the original position of the foot of the amputee who complains of pain in his lost foot—always, be it noted, at a leg's length from the trunk.

The small child suffering from a slight attack of appendicular colic has no doubt about the locality of his pain. When asked, "Where is the pain?" he puts a finger in his umbilicus and replies, "In there." When the pain is more severe he places the whole hand over the umbilicus.

The Pain is Ill-defined.—This is a relative term, of course, but if one considers its mechanism it is just the type of pain one should expect. The common cause is a wave of hyperperistalsis in the bowel proximal to an irritant or obstruction. There is no abrupt transition from normal bowel to strongly contracting bowel: the action is wave-like (crescendo-diminuendo). Many patients describe their pain as "coming in waves." Who can tell where the wave begins or where the ripple ends?

Diaphragmatic Pain : Shoulder-tip Pain

Morley has emphasized the importance of shoulder-tip pain as a symptom in certain abdominal conditions. I have repeated his experiment under spinal anaesthesia, and can confirm the observation that even light irritation of the peritoneum covering the central portion of the diaphragm causes severe pain in the shoulder. A point of great interest and significance is that the conscious patient feels no pain or other sensation locally from the presence of the finger or swab in contact with his diaphragm.

Mainly from these experiments, Morley has formulated his theory of peritoneo-cutaneous radiation. He further states that if the shoulder-tip is rendered insensitive by injection of a local anaesthetic the pain is much less severe. He does not claim that the injection completely abolishes or prevents pain in the shoulder-tip when the under surface of the diaphragm is then stimulated, and writes:

"That shoulder-tip pain was not abolished completely by the cutaneous anaesthesia may be attributed to the fact that the descending cervical nerves from the third and fourth segments supply sensory fibres to the acromio-clavicular joint as well as to the periosteum of the acromion process and clavicle. Since care was taken to make the subcutaneous injection as superficial as possible, these fibres would escape the action of the novocain, and this would probably account for such pain as persisted after the shoulder-tip area had been anaesthetized. It can be stated with the greatest confidence that none of the novocain solution was injected anywhere near the trunk of the phrenic nerve, so that a possible fallacy in the experiment—viz., anaesthesia of the phrenic nerve—need not be considered."

Is there no other explanation why the relief from pain was only partial and not complete? Did suggestion—auto- and hetero—have any effect in relieving the pain? Did the local anaesthetic have any counter-irritant effect? Would complete anaesthesia of *all* the descending sensory nerves, including those to the clavicle, acromion, and acromio-clavicular joint, have entirely prevented this "peritoneo-cutaneous radiation"? Here it will suffice to say that in circumstances in which clinician and patient both expect a certain phenomenon to happen the more likely will their anticipation become realization.

My personal view is that when the central portion of the diaphragm is irritated shoulder-tip pain will occur (a) if all fibres of the descending cervical nerves mentioned are completely anaesthetized, and (b) if even a fore-quarter amputation were performed.

Here we have an organ—the central portion of the diaphragm—which has migrated from the neck, taking its original nerve supply (phrenic nerve from C3, C4, and C5) with it to its secondary or permanent position, where it has not acquired a new or additional innervation. This is analogous to what happens in the intestinal tract: therefore pain produced in this migrated organ will be localized in the primary relative position of the diaphragm. Such, at any rate, is my belief. We must of course remember that the diaphragm has an upper, thoracic surface, as well as an abdominal surface, and that medical conditions such as pleurisy, pulmonary embolism, and pericarditis can also give rise to shoulder-tip pain.

Conclusion

I anticipate the question, "Why is testicular pain localized in the testis, a migrated organ?" The answer is that the local pain is due to irritation of the tunica vaginalis, which is derived from a process of *parietal* peritoneum and supplied by the genito-femoral nerve—a somatic nerve from L1 and L2. Both parietal and visceral tunicae are sensitive, and so provide a protective mechanism for the testis in its very exposed position. The testis itself has a sympathetic innervation from the tenth or eleventh and twelfth thoracic segments (authorities differ slightly) through the renal and aortic plexuses. These nerves accompany the testicular vessels. True testicular pain will, like ovarian pain, be felt in the loin, since both testis and ovary have migrated from this position. It is said that two types of pain occur in injury to the testis: slight trauma causing local pain, and more severe trauma producing pain in the loin—a very perfect protective arrangement.

Here then, I submit, we have a rule without exception—in other words, a *law*, which may be stated thus: *Pain produced in an organ which has migrated from its primary relative position, and which has not acquired an additional nerve supply in its secondary or permanent position, is invariably localized in the primary relative position of that organ.* This law may be amplified to include artificially displaced tissues and other sensations besides pain, thus: Sensations produced in tissues which have been developmentally or artificially displaced from their primary relative positions and which retain only their original innervation are invariably localized in the respective primary relative positions of those tissues.

Summary

The development of the intestinal tract is briefly described.

The sensitivity of the intestine and parietal peritoneum is discussed, also the adequate stimuli which produce visceral and somatic pain.

Aristotle's experiment is described and analysed.

The view is advanced that the "false localizations" which occur in this experiment and elaborations of it are analogous to the "retrojected" pains produced in organs which have migrated from their primary relative positions.

I am indebted to Dr. Janet Conn for the diagrams.

BIBLIOGRAPHY

- Aristotle (384–322 B.C.). *Problemata* (The Works of Aristotle, English translation: editor, W. D. Ross: Vol. VII, *Problemata*, by E. S. Foster, 1927).
- Czermak, J. N. (1855). *S.B. Akad. Wiss. Wien.*, 15, 514.
- Head, H. (1893). *Brain*, 16, 1.
- Hilton, John (1863). *On Rest and Pain*, London.
- Hurst, A. F. (1911). *The Sensibility of the Alimentary Canal*, London.
- (1941). *British Medical Journal*, 1, 415.
- Mackenzie, J. (1920). *Symptoms and their Interpretation*, 4th ed., London.
- Morley, John (1931). *Abdominal Pain*, Edinburgh.
- Musser, J. H. (1904). *Amer. Med.*, 7, 503.
- Neumann, A. (1910). *Cbl. Genz. Med. Chir.*, 13, 401.
- Nothnagel, H. (1905). *Arch. Verdaukr.*, 11, 117.
- Ross, J. (1888). *Brain*, 10, 333.
- Ryle, J. A. (1926). *Lancet*, 1, 895.
- Starling's *Principles of Human Physiology*, 8th ed.; editor, C. Lovatt Evans (1941), London.
- Tastevin, J. (1937). *Encéphale*, 32, Pt. I, 57, 140.

The number of wartime nurseries established in Scotland for the young children of working mothers is growing steadily despite difficulties in securing suitable premises and building labour and equipment. Edinburgh now has seven nurseries in operation, Glasgow five, Aberdeen two, and Coatbridge, Falkirk, and Paisley one each. There is a keen demand for 63 other nurseries which were expected to be open throughout Scotland by the end of April or the beginning of May. One of the encouraging features of nursery development has been the rapidity with which staffing problems have been overcome. Many married women have volunteered as matrons, deputy matrons, and assistant nurses; Child Care Reserve courses are full; many applications for probationerships have been received from girls leaving or about to leave school. A booklet which will guide nursery superintendents—and be something in the nature of a textbook for younger workers—on the feeding of infants and pre-school children has been prepared by the Department of Health for Scotland; it deals particularly with Scottish requirements and is based on food supplies available in wartime.