but was abandoned because of the complications found during operative procedure.

A resection of the affected ileum, caecum, and part of the colon is obviously the correct method of treatment in all cases except those in which the infection is quiescent, and it now seems to be generally recognized that a two-stage operation is to be preferred. Mayo and Judd (1941) gave the mortality rate as 22% for cases subjected to a one-stage operation, and only 3% when a preliminary short circuit had been done. The delay between stages is open to variation in individual cases, but there is no object in unduly delaying the resection, and we feel that one month might be considered to be the optimum interval. Additional safety may be obtained in a single-stage operation by the exteriorization of the anastomosis or by a temporary enterostomy or colostomy. Mayo and Judd advised leaving out the proximal portion of the transverse colon and using it as a temporary colostomy.

There will be certain advanced types of lesion, complicated by fistula formation and adhesions, which cannot possibly be treated along any set lines. In these cases surgical treatment may be extremely difficult and hazardous, and will probably have to be carried out in multiple stages.

Summary

Two cases of regional ileitis (Crohn's disease) in women aged 22 and 18 are described, especially from the aspects of diagnosis and treatment. The importance of roentgenological examination of the intestinal tract after a barium meal as an aid to diagnosis and the advisability of a two-stage operation are indicated.

We are indebted to Dr. E. Rohan Williams for the radiological reports, and to Dr. G. E. Harre for the pathological reports.

REFERENCES

- Crohn, B. B. (1939). *Surg. Gynec. Obstet.*, **68**, 314.
 — Ginzburg, L., and Oppenheimer, G. D. (1932). *J. Amer. med. Ass.*, **99**, 1323.
 Donald, C. J., and Brown, P. W. (1941). *Proc. Mayo Clin.*, **16**, 561.
 Mayo, C. W., and Judd, E. S. (1941). *J. Amer. med. Ass.*, **117**, 836.
 Simpson, S. L. (1935). *Quart. J. Med.*, n.s., **4**, 191.

THE PATHOGENESIS OF PULMONARY TUBERCULOSIS OF THE ADULT TYPE

BY

R. C. WINGFIELD, M.B., F.R.C.P.

Much time and thought have been expended on the pathogenesis and exact method of implantation of the initial lesions of pulmonary tuberculosis of the adult type. I think we should be wiser, so long as our views are in the hypothetical stage, to avoid assumptions and dogmatic statements whenever possible, and to be careful, where assumptions are used, to base as little as possible upon them in the arguments that arise. Another mistake we make is in not taking into account the possibility that there can be more than one answer to the question. Such an attitude limits our endeavours and makes it very difficult to fit the facts at our disposal into our hypotheses without a deal of juggling.

I read in an annotation in the *British Medical Journal* of March 7 (p. 332): "There is evidence that . . . are responsible for the development of bronchogenic tuberculosis (*the common type of pulmonary tuberculosis in adults*). The italics are mine. Here is an assumption which I think is unwarranted. I have read many arguments and studied the data of much research that lead me to suppose that the common type of pulmonary tuberculosis may be bronchogenic in many or even the majority of cases, but I have never found anything approaching a proof that this is a fact, and therefore to build up an argument on the assumption that it is unreasonable and dangerous. Those who have studied the published work on the subject will be able to pick out other dangerously used assumptions like the instance I have given.

It is impossible to fix a truth in the scientific field of knowledge. A hypothesis that satisfies the facts passes into accepted

theory, and the theory holds good and must be accepted as truth just so long as no fresh facts come to light which prove it wrong. It is essential in trying to build a hypothesis to explain the pathogenesis of pulmonary tuberculosis that such a hypothesis must be asked to satisfy facts only, and although assumptions must be used in its structure it should not be based upon them.

Accepted Facts

The following are the facts which I believe are accepted by all students of the subject:

1. The majority of the urban population and a large proportion of the rural population are infected by tuberculosis by the age of 30.
2. Pulmonary tuberculosis of the adult type is not the direct result of a primary infection, although in some cases it may follow very closely on its heels.
3. The lesion resulting from a primary infection, if in the lung (Ghon's focus), is due to an inhalation infection. It may be situated in any part of the lung tissue, but the greater number are found in the right middle and lower and left lower lobes.
4. The early lesion of adult-type pulmonary tuberculosis may appear in any part of the lung, but the overwhelming majority are found in the upper thirds of the upper lobes and more often in the right lung than in the left.
5. When examining unselected samples of the population and unselected samples of contacts, adult-type pulmonary tuberculosis is found ten times more frequently among the contacts. The figures are roughly 0.5% and 5%.

It is tempting to add other "facts," but I have not, as they are facts only to me or to a restricted circle of students, and have not been generally accepted; they must therefore fall into the category of assumptions.

Questions to be Answered

The questions the hypothesis must answer are: (1) Why and how does a person get adult-type pulmonary tuberculosis? The possibilities are (a) by exogenous superinfection, (b) by endogenous exacerbation, (c) by a combination of (a) and (b). (2) By what route does the tubercle bacillus reach the site of the initial lesion of this type of pulmonary tuberculosis—by the respiratory tract, by the blood stream, or by the lymph stream?

Facts Nos. 1, 2, and 5 force the answer to the first part of the first question. Adult-type pulmonary tuberculosis is found in those persons who have been previously infected and is favoured by close contact with persons suffering from pulmonary tuberculosis with positive sputum—that is to say, by superinfection.

The three subsidiary questions to No. 1 and question No. 2 are more complicated, and are so bound up with one another that it is difficult to take them separately; they must be discussed together.

The simplest answer to the remaining question would be that in close contact with sputum-positive cases tubercle bacilli are inhaled in sputum droplets and dust and pass via the respiratory passage to the lung and there set up their characteristic lesions. But this simple answer does not satisfy the facts. Indeed, if it was so simple as that the new lesions would be multiple in most cases and have the same distribution as Ghon foci, and the incidence of the disease among contacts would probably be nearer 100% than 10%.

Ghon foci are definitely inhalation infections, and their distribution in the lungs must therefore give us information on the mechanism and results of inhalation infections. Ghon foci are found in either lung and distributed fairly impartially in all lobes, but the sites of election are the right middle and lower and the left lower lobes. This seems to contradict Bannen's (1942) assertion that an inhalation infection should pass most easily and frequently into the right upper lobe. The fact that an abnormal oily liquid (lipiodol) artificially introduced into the respiratory passages is usually coughed into the bronchus of the right upper lobe is not a conclusive argument. It is an experimental finding which it is dangerous to assume may be repeated under natural conditions. There is a great difference between a blob of lipiodol artificially introduced into the trachea and the 10 μ diameter droplet naturally inhaled (Jacobs, 1941). Bannen states that radiologically (presumably)

haematogenous pulmonary tuberculosis lesions are always easily recognizable. This is also an assumption which I, after many years' experience of x-ray interpretation, cannot corroborate. Jacobs in a well-reasoned argument shows the unlikelihood that inhalation superinfection is a cause of pulmonary tuberculosis of the adult type. Yet it seems that close contact and crowding, favouring frequent and massive inhalation infections, are outstanding in the aetiology of the disease.

Two Possible Causative Factors

I think that two possible factors have not been taken sufficiently into account in the many arguments on this subject.

1. What is the effect of frequent doses of inhaled or ingested tubercle bacilli from without on the immunity balance between a host and the tubercle bacilli he already harbours? If these inhaled or ingested bacilli lodge on a mucous surface—and those in close contact with freely talking and coughing open cases must inhale and ingest large quantities of bacilli daily over long periods of time—they must be taken into the system, and if they do not themselves actually cause a new tuberculous lesion they must be destroyed. In their destruction the protein moiety of the bacillus, to which all infected persons are sensitive, will be liberated and must have an effect on the host of exactly the same quality as that of a dose of tuberculin. If such doses of the protein moiety of the tubercle bacillus are frequent and large, some disturbance of the immunity balance in one direction or another is inevitable. An adverse disturbance, if of any great degree or of an appreciable duration of time, should influence the behaviour of the tubercle bacilli lying dormant and clinically inactive in the lymphatic system or elsewhere of all infected persons. I say "elsewhere" with reason, for all are familiar with the small tuberculous focus in the lung which remains unchanged in size for years and yet is seen to be the possible starting-point of a spreading granuloma at some later date.

2. The last barrier between the lymphatic system (which in all Mantoux-positive individuals harbours tubercle bacilli) and the blood stream is the bronchial gland system. Once past these glands the lymph flows unimpeded into the blood stream via the lymphatic duct. Krause's experiments in guinea-pigs show that tubercle bacilli, whatever the site of entry into the body, unless arrested on their travels will always percolate the lymphatic system centripetally, finishing with involvement of the bronchial glands; and, further, that it is not until the bronchial glands have been involved that dissemination to other organs occurs. If the site of entry is the muscles of the leg it will obviously take the bacilli longer to reach the bronchial glands than when the site of entry is the bronchial mucosa, and if the site of entry is the intestinal mucosa they will take an intermediate time. The size of dose and the virulence of the bacilli will also influence their rate of travel. Furthermore, the further they have to travel the greater the number of glands or filters they have to pass and the greater the chances of their being temporarily or permanently arrested on their journey.

In those persons whose bronchial glands contain tubercle bacilli it is not hard to suppose that an adverse disturbance of the immunity balance caused by repeated ingestions of non-lesion-producing tubercle bacilli from an outside source might result in increased activity of the resident bacilli and a seepage of them into the lymphatic duct and thence to the blood stream.

It is not necessary to suppose that tubercle bacilli in the blood stream will always produce widespread generalized military tuberculosis: we are all familiar with localized miliary-type infection. Generalized military tuberculosis, a comparatively rare event, surely presupposes a gross bacillaemia. The body, especially of the Mantoux-positive, is relatively tolerant of the tubercle bacillus, and in the event of a small entry into the blood stream it is quite possible that such bacilli, when deposited in small numbers in the lung tissues (the lung being the first capillary system they will encounter), may cause no lesion and be destroyed, or if not destroyed may only produce a minimal "secondary" and a non-progressive granuloma, in which they may ensconce themselves for hibernation. Others, again, may be deposited in larger numbers or be more vigorous and be the starting-point of the usual spreading granuloma.

Our knowledge of the rate of flow of the blood in the different parts of the lungs is scanty, but in view of the fact that in

normal breathing only the lower halves of the lower lobes are used it is quite possible that without the mechanical acceleration of lung inflation and deflation the blood flow is relatively more sluggish in the upper lobes. This would favour the deposit there of blood-borne bacilli and would fit in with the preponderance of initial lesions, and of new lesions in those already diseased, in these areas. To quote again from Bannen, the fact that most people sleep on their right side, which further restricts the movement of the right hemithorax, might possibly be used as another argument for this being the commonest site for new haematogenous lesions.

A Hypothetical Answer

A hypothetical answer to the questions would then read as follows: Pulmonary tuberculosis of the adult type is found in those persons who have been previously infected and is favoured by close contact with persons who are excreting tubercle bacilli. It is in these cases an exogenous superinfection. It is improbable that the initial lung lesions are caused directly by the inhaled tubercle bacilli. It is possible that these lesions, and new lesions in quiescent cases, may be caused by (a) bacilli which come to the lung in the blood stream from foci of living bacilli already in the lymphatic system, or (b) bacilli dormant in minimal and unrecognized lung lesions, whose activity in both cases has been increased by frequent massive ingestion into the system of tubercle bacilli from without whose destruction in the body liberates unknown quantities of the protein moiety of the bacillus to which the infected person is sensitive. That is to say, exogenous superinfection causes endogenous exacerbation.

This may appear a complicated hypothesis; but if a simple explanation fails, complexity is unavoidable. I put it forward as a possible hypothesis, not claiming that it is a satisfactory one, that it is true, or even that it excludes others; but as a basis for further study of this fundamentally important subject.

REFERENCES

- Bannen, J. E. (1942). *British Medical Journal*, 1, 309.
Jacobs, A. L. (1941). *Tubercle*, 22, 266.

OEDEMA OF THE GLOTTIS COMPLICATING MEASLES

A REVIEW OF EIGHT PERSONAL CASES

BY

J. H. MULVANY, F.R.C.S., M.R.C.P., M.R.C.O.G.

Surgeon, Emergency Medical Service

The recent account in the *Journal* (March 7, p. 325) of 4 cases of coincidental association of measles and laryngeal diphtheria prompts me to place on record the following brief summary of 8 cases of oedema of the glottis complicating measles. It is doubtful whether the possibility of this complication occurring save as a rare eventuality is generally recognized, as most textbooks of medicine and surgery do not include measles in the list of causative factors of oedema of the glottis, nor, in the general sections where reference to the complication may be made, is there anything to indicate that it is more common than such rare abnormalities as abscess of the brain, optic atrophy, or peroneal muscular dystrophy. Measles is as remarkable in the constancy of certain of its features as in the variation, in successive epidemics, of the dominant complication. Thus, bronchopneumonia, mastoid infection, and enterocolitis are common events which may take their turn in this fashion, and less frequently so may oedema of the glottis.

Clinical History of the Cases

In the widespread and severe outbreak of the winter of 1929-30, while in practice, my colleagues and I attended some 950 cases of measles. Of this number about two-thirds were moderately to severely ill, and included 186 cases of bronchopneumonia, 48 of them fatal; but the outstanding circumstance