Experience in the use of amnioplastin has been obtained in 8 intracranial cases, 12 peripheral nerve cases, and in one or two additional instances, such as a scar in the popliteal space. The type of peripheral nerve case in which the membrane has been used, the method of its employment, and the results so far obtained are here reported. These results are so encouraging as to suggest its continued and further use.

REFERENCES

- Chao, Y.-C., Humphreys, S., and Penfield, W. (1940). *British Medical Journal*, 1, 517.
Penfield, Wilder (1940). *Ibid.*, 2, 668.

A HISTORY OF THE RESCUE AND AFTER-CARE OF TWO CASES AFTER FOUR DAYS' BURIAL

BY

G. S. SWAN, M.B., Ch.B.

On a Thursday, at 1 a.m., a bomb struck a block of three-storied cellared houses. Some of these houses were completely destroyed. As it was known that a number of people were in them at the time of the disaster rescue work started almost at once, but with little hope of saving anyone alive. At 6 p.m. on Friday an elderly woman was rescued alive, but in such a serious condition that she died a few days later.

The First Rescue

At 7 p.m. on Sunday I was sent for, as the rescue squad had heard a voice under the debris. On arrival I made contact with the trapped girl, one of my own patients, Miss F. B., aged 21, whose voice could faintly be heard, obviously at some distance below the debris. She told me her name, that she knew her father and mother were dead, and that she was quite immobile except for a little movement of the left hand. It was astonishing to find her so completely intelligent, and throughout the six hours before she was finally removed at 1 a.m. on Monday—exactly ninety-six hours from the time she was first trapped—she retained her intelligence and was of great help in guiding the rescue squad in their difficult task.

At 9.30 p.m., debris having been removed entirely by hand and several wooden beams sawn through, a narrow hole four feet in depth allowed of contact with the girl's left hand. It was just possible to feel her pulse, which was well perceptible at about 90 beats per minute. A short time afterwards her head was uncovered, when it became obvious she was face downwards in position. By means of a length of rubber tubing and a feeding-cup we succeeded in getting her to rinse out her mouth, after which she was allowed to swallow a little hot well-sugared tea. Shortly afterwards, as more of her body became exposed, I gave her one-third of a grain of morphine, and about quarter-hourly sips of tea, by now out of the feeding-cup direct. By midnight she was completely exposed at the bottom of a deepish hole. The rescue squad also succeeded in burrowing sideways at her body level, a burrow sufficient to admit head and shoulders with debris supported by some flooring cross-beams. It was through this burrow that we succeeded in delivering her.

She was lying crouched, face down, her knees on her chest and heels under her buttocks, her right arm under her chest and left arm by the side of her face—a normal breech position. Across her back was the dead body of her father. It was evident we dared not remove him

before the girl, so we arranged the rescue squad with one man from above easing up the dead body as much as possible, another down the main hole supporting the chin, and a third grasping her hips and gently pulling her through the small narrow side burrow. As she was being removed she told us there was a woman alive next door. Once on the warm stretcher I immediately straightened her legs and sent her to hospital, where all was in readiness for her reception.

The Second Rescue

On investigating her report that a woman was alive next door, we found, to our surprise, that this was true. After shouting we heard a voice say, "I am A. D. [aged 34]; I am all right, and walking about in the cellar. I have a baby with me, who is not very well, and we have some bottles of milk." This turned out to be delirium, and it was some hours before contact was made. It was then discovered that she was lying on the cellar stairs, her head higher than her feet, and completely immobile.

The only way to deliver her was through a narrow aperture formed by the absence of the upper panel of a door, the rest of the door supporting a mass of debris. It was impossible to risk cutting away the side of the door, but it was possible to remove a triangular piece of the lower panel. Her pulse was weaker than the girl's, but countable at about 100. She also received morphine, and was fed in a similar way to the first patient. The dead body of a 3½-year-old boy was lying under her right arm. To deliver her through this narrow aperture one man was able to step inside the hole and just scramble down to place both hands under her shoulders; another man, lying on the debris held by his heels, was just able to reach her legs. I lay over the aperture and, as she was eased upwards, succeeded in grasping her shoulders and bringing her out of the hole over my own right shoulder to stretcher-bearers behind. This was at 6 a.m. on the Monday, 101 hours after being buried. She also was immediately sent to hospital.

Treatment of the Patients

When admitted both patients were placed in warm beds with suspended cradle lamps and received a transfusion of two pints of plasma. Blood pressure on admission was: Miss F. B., systolic 85, diastolic ?; Miss A. D., systolic 90, diastolic ? 70. After transfusion both blood pressures were 140/80.

General Examination of Case I.—Severely shocked, but perfectly conscious and mentally very alert. Very thirsty. Eyes bright and free from grit; lips and tongue abraded and some dirt ingrained; air passages clear—no evidence of gritty lungs. A few small pressure sores on back and large area over sacrum. Full range of movement in upper limbs; small superficial patches of skin gangrene over patellae; both lower limbs stiff, and foot-drop present on left. One gramme of sulphanilamide four-hourly for forty-eight hours, followed by one tablet (0.5 gramme) four-hourly for four days. Frequent hot drinks taken very well. All pressure sores dressed with gentian violet; nursed on an air-bed; right foot splinted; rest of skin perfectly clear. Extremities were warm.

General Examination of Case II.—Severe shock; cold; mental condition incoherent with periods of delirium; very thirsty. Eyes clear; tongue dry; air passages clear—no evidence of gritty lungs. Bruising of both legs, with large gangrenous patch on right thigh posteriorly, lateral aspect of right leg, and above left outer malleolus; a smaller patch of superficial skin gangrene on right side of chest, also on both wrists posteriorly. Patient was not able to move either of lower limbs; the right leg and foot were oedematous, also right hand and arm. Warm fluids were

taken freely. All patches were treated with gentian violet, and sulphanilamide was given as in previous case.

Both patients also received 2,000 units A.T.S. Within twenty-four hours mental condition became perfectly clear; right foot-drop developed, and foot was accordingly splinted.

Subsequent History

Both patients steadily improved. No vomiting; solid food was begun on the third day, following an enema. On the fourth day both began to bend the affected knee; oedema steadily diminished. Urine, tested daily, revealed nothing abnormal, and the output was never less than 75% of intake. As in cases reported in the *Journal* of March 22, the non-protein nitrogen was estimated on both, and was found to be well within the normal limits—25 mg. per 100 c.cm. in one and 27 mg. in the other.

By the end of the week the oedema had practically disappeared; there had been no vomiting, the bowels were acting normally, blood pressure in both was 148/85, and fluid output was equal to intake. After ten days all gangrenous patches were separating at the edges, leaving clean granulating surfaces underneath. No oedema was present. Slight movement of the foot was noticed.

At this stage Miss A. D. showed evidence of slight left ulnar nerve injury with some median involvement, only fingers affected; the hand was splinted in dorsiflexion. At the time of writing, thirteen days after admission, this is already showing signs of clearing up, and the general condition of both patients is excellent. Stringent precautions have been taken all along, no visitors having been allowed until the eleventh day and the cradle lamps still being in use. Both are eating and sleeping normally, and there has been no change in their mental condition. They have not talked much of their experiences, nor have they as yet been questioned.

Commentary

These two cases raise several points of both general and medical importance. I have suggested to the municipal authorities that it would be of great advantage if some form of sound-detector could be employed on the site of a disaster as early as possible. Both patients state that whereas they could hear everything going on above them—and Miss F. B. says that her father was alive for two days and Miss A. D. that the child lived for nearly three days—their voices were not heard until sufficient debris had been removed nearly four days afterwards. The reason for and advantages of a sound-detector are therefore obvious. I suggest also that rescue work should be carried out on the assumption that people are alive though buried, and that work should be done in relays without ceasing. All ambulances should normally carry hot-water bottles (I used three of my own on this occasion) and six feet of rubber tubing with feeding-cup.

The important medical points that occur to one, bearing in mind the positions in which the patients were found and the length of immobility, are: (1) The severe limitation of oxygen supply, and the lack of fluid intake, both acting together, failed to produce the obvious mental and physical reactions. (2) The extraordinarily good condition of the skin and the warmth of the extremities. (3) The absence of irritation conjunctivitis. (4) The clearness of the air passages and the absence of gritty lungs. (5) The comparatively mild nerve-pressure paralysis.

I believe a doctor working in conjunction with the rescue squad is the ideal arrangement when there is any chance of rescuing people alive, and would stress the importance of some form of sound-detector and my belief that the dangerous period after prolonged burying is that between the moment when exposure takes place and the

patient's arrival in hospital. Thus treatment on the lines already indicated should be instituted during the actual rescue. Lastly, I would support Prof. H. A. Harris in his criticism of over-treatment. Briefly put, the treatment of "crush cases" should provide a maximum of rest, both physical and mental; a maximum of warmth, both external and internal; and a minimum of interference.

I would like to thank my resident, Miss Walsh, and the nursing staff of the hospital, for their whole-hearted co-operation.

Clinical Memoranda

Oestradiol Benzoate in Hypoglandular Rheumatoid Arthritis

During the past season a number of female patients with rheumatoid arthritis had very limited time to complete a satisfactory course of spa treatment owing to home duties of an urgent nature. Six weeks would be an ideal course for the average case, but some patients could spare only two weeks, and less than three is of very little use.

In the early spring of 1940 I noticed that those who were taking small doses of oestradiol preparations or the synthetic substance stilboestrol for menstrual disorders showed remarkable improvement in coexisting hypoglandular rheumatoid arthritis. In all these cases I decided to supplement treatment with intramuscular injections of oestradiol benzoate in oil. The results have been beyond all expectation, and I now use this method as a routine. Many patients felt distinct improvement within twelve hours of the injections.

The cases varied from slight periarticular thickenings of the hands to marked multiple thickened joints. In one there was complete ankylosis of the knee-joints, but the general health greatly improved. Two patients had multiple thickened joints, pathological deformity of hands, knees, and feet, and very poor mobility. They were becoming totally disabled and were very depressed, and considered their cases to be hopeless. After several weeks' treatment all the mild cases cleared up completely. The two advanced ones made a remarkable recovery. One is almost free from any signs, and the other patient is fit for light occupation.

Oestradiol benzoate has been of great benefit in combination with spa treatment owing to the fact that patients suffering from depression often state that they have a feeling of general well-being during the first day of its use. Injections were given twice weekly intramuscularly in addition to the spa treatment. A high dosage of 50,000 units was used in severe cases, and 10,000 units and upwards for milder ones. Ampoules representing both strengths or a bulk quantity can be used, and a 1-inch-long No. 21 needle was found suitable. Injections were made into the deltoid and were quite painless. The syringe can be kept free from any moisture in a pocket spirit case containing 5% lysol added to the spirit. It is very important to warn these patients, irrespective of age, not to be alarmed if a menstrual cycle begins some days or weeks after the injection. The flow, if troublesome, can be regulated by giving calcium and ergot preparations, but this was not required, or if the cycle date is noted a corpus luteum preparation could be given daily for a week before the next cycle. This preparation is costly and its use in rheumatoid arthritis may not be of any particular advantage.

Oestradiol benzoate is tolerated by all, and has no harmful effects even in pregnancy. Male cases of rheumatoid arthritis also respond to oestradiol benzoate injections, but I have tried only a short course on a few cases in view of any disturbance to the prostate gland. Stilboestrol can be used as a substitute, but in a few cases nausea was complained of even with a small dosage. A recent improvement has been made to stilboestrol which obviates any nausea.

JOHN A. MURRAY, L.R.C.P. & S.Ed.
Strathpeffer Spa, Ross-shire.