CHEST SURGERY IN WAR

This is one of a short series of articles based on lectures given at the British Postgraduate Medical School, Hammersmith

ABDOMINO-THORACIC INJURIES *

BY

GORDON GORDON-TAYLOR, M.S., F.R.C.S.

Temporary Surgeon Rear-Admiral

TREATMENT OF PENETRATING ABDOMINO-THORACIC INJURIES

The surgery of the thoracic wall and cavity is being dealt with in this series of lectures by those most expert in the surgery of this region. My contribution is therefore only concerned with the surgical treatment of injuries of the diaphragm and of those abdominal organs whose anatomical proximity renders them specially vulnerable in cases of diaphragmatic wounding.

I. Wounds implicating the Liver

The dimensions and the velocity of the missile play an important part in determining the type of hepatic lesion, which may be almost protean in character; large fragments of liver may be torn off, and as much as one-third of the right lobe has been found loose in the peritoneal cavity of a patient who recovered from a penetrating abdomino-thoracic injury in the last war.

Before their total destruction the examination of museum specimens in the College of Surgeons collection illustrating gunshot wounds of the liver revealed many lesions for which surgery must be futile or incomplete: thus it would be impossible by operative measures alone to prevent infection throughout the length of a ragged track running from the right lateral surface of the organ to its extreme pole on the other side of the body. In many cases of liver injury due to enemy action there are few arresting clinical signs, and it is significant that out of all the penetrating injuries of the abdomen which recovered in the last war without operation the wounds were mostly situated in the liver area. *Surgical intervention* directed towards the liver is indicated under the following circumstances only: (1) *gross haemorrhage from the liver*, which mainly depends on whether large veins in the organ have been injured—haemorrhage otherwise ceases spontaneously, as does oozing of bile, unless main ducts are severed; (2) the association of some thoracic or another abdominal lesion demanding exploration; (3) the retention of a missile in the liver, especially a large missile in an accessible position in that organ. In the case of severe laceration of the liver packing may be necessary to control haemorrhage. In some such patients the diaphragm may be sutured, and the thorax closed in the approved manner with airtight drainage of the pleura by means of a Malecot's tube inserted through a separate intercostal incision. The gauze pack is brought out through another separate incision in the anterior abdominal wall. In other cases the diaphragmatic opening may be enlarged parallel with its fibres from mediastinal to lateral border; the diaphragmatic edges are now approximated to the intercostal muscles and the pleura is thus shut off from the drainage track.

Early jaundice in cases of hepatic injury may be present, and is usually slight and evanescent; the possibility of a collection of bile in peritoneum or pelvis should always be considered. *Late jaundice* has a much graver significance, and indicates serious infection.

Secondary hepatic haemorrhage in penetrating abdomino-thoracic injuries is almost always fatal, and is usually associated with signs of sepsis such as pain, elevation of temperature and pulse rate, abdominal pallor, restlessness, and loss of strength.

The formation of an *intrahepatic abscess* round a retained fragment may be dealt with successfully by surgical measures.

II. Gunshot Injury of the Spleen

A study of the specimens of *gunshot injury of the spleen* in the War Collection formerly housed in the Royal College of Surgeons used to suggest that wounds of that organ are frequently complicated by damage to adjacent viscera; yet in some collected clinical series it is stated that in more than half the cases of splenic injury revealed at operation there was no damage to other abdominal organs. This was certainly not my experience (Gordon-Taylor, 1939), which is in agreement with that of Major Jolly in his surgical work with the Spanish Republican Army (Jolly, 1940).

The propriety of primary splenectomy in cases of gunshot injury will still remain a subject of controversy. The removal of the organ should certainly not be resorted to light-heartedly, but in my opinion the procedure has not greatly increased the risk. The suture of the spleen may not be difficult when the organ is approached through the chest and diaphragm, but when explored by other routes cobbling of the organ is wasteful of time and may leave a sense of insecurity. Jolly (1940) and some of the Spanish surgeons write enthusiastically of splenectomy.

Numerous cases of successful transdiaphragmatic splenectomy in left-sided abdomino-thoracic wounds were recorded in the war of 1914-18, some of which were associated with the operative repair of injuries to the stomach (Richard Charles, Gask, J. E. H. Roberts, Lockwood, Bryan, Gordon-Taylor, etc., and very recently by Valentine Logue). Recovery has even taken place in cases of left-sided abdomino-thoracic wounds in which left-sided nephrectomy as well as splenectomy had to be performed for complete disruption of each organ (Gordon-Taylor).

III. Wounds of the Pancreas

These injuries were less frequent in the war of 1914-18, or at any rate were less often recognized, than injuries of any other of the abdominal viscera. The intimate anatomical relation of the gland to the large blood vessels doubtless explains the paucity of wounds coming under the observation of the casualty hospital surgeons; doubtless some were unrecognized and were designated retroperitoneal haematomata. I have knowledge of only three abdomino-thoracic injuries implicating the pancreas in which recovery took place (Saint, Alan Curry, Gordon Bryan).

IV. Injury to the Kidney

Each kidney is equally liable to damage. It was estimated that in 40% of gunshot wounds of the kidney investigated at an English surgical base in France during 1914-18 the thorax was also implicated. The propriety of immediate operation on the kidney in abdomino-thoracic wounds demands even more careful consideration than in purely abdominal injuries. Probably the best course is to excise the wound leading down to the damaged kidney; the organ can then be inspected and any foreign body removed.

* Concluded from p. 864.

In the majority of cases the renal injury plays a minor part in menacing life, and only for severe haemorrhage, a hopelessly pulped organ, or gross damage to the renal pedicle is primary nephrectomy indicated. In about 20% of cases of gunshot injury of the kidney in the last war life was threatened by secondary haemorrhage, consequent on sepsis, a retained missile, or an infarct of the kidney produced by damage to some branch of the renal artery.

V. Injury to the Hollow Organs

Of hollow organs the *stomach and splenic flexure of the colon* are those most liable to injury; in suitable cases transdiaphragmatic laparotomy affords excellent access to these organs. Concomitant involvement of the *small intestine* denotes a more vertical track of the missile, and demands a separate approach or a much-extended incision. Each measure, though necessary, increases risk to life. Multivisceral damage also augments mortality, and vertical wounds, which may even extend from supraclavicular fossa to femur, tend to be very fatal.

The possibility of injury to the *duodenum* must be borne in mind, and any haematoma in its vicinity should be suspect, lest a retroperitoneal rupture of this fixed part of the small gut be overlooked. These cases were very fatal in the last war; successful cases have recently been operated upon by Rex Williams, Henry Thompson, and Hedley Atkins.

Summary of Treatment

1. In many traversing (through-and-through) abdomino-thoracic wounds of the right side produced by a small fragment no immediate active surgical treatment is required, provided that: (a) no gross damage has been inflicted upon the thoracic or abdominal wall—fractured ribs, explosive effect, etc.; (b) the direction of the track of the missile does not appear to compromise the general peritoneal cavity or suggest the desirability of its exploration; (c) the signs of abdominal haemorrhage or of injury to a hollow viscus are clearly absent.

2. In cases of right-sided abdomino-thoracic wounds in which a small fragment is retained in an inaccessible position in the substance of the liver an expectant line of treatment is the correct procedure; accessible fragments, unless of small dimensions, should be sought and removed.

3. When there is an open, blowing thoracic wound or a "stove-in chest," the chest wound should of course assume chronological priority of treatment.

4. If the position of the wound of entry and exit in a left-sided abdomino-thoracic wound adumbrates a track implicating that fatal left subphrenic area of the abdomen, or if a radiograph demonstrates a fragment of metal retained in this region, the thorax should be dealt with first and access to the upper abdomen obtained through the diaphragm.

5. When the thoracic injury appears insignificant but there is evidence of severe widespread intraperitoneal damage, especially involvement of hollow viscera, the abdomen should be explored through an appropriately placed laparotomy incision. This instruction applies to wounds of thorax and abdomen produced by the same or by separate missiles.

6. When the thoracic injury seems slight and when the evidence of a radiograph or the direction of a missile track in a through-and-through wound suggests an extra-peritoneal course of a small fragment, such an abdomino-thoracic injury may often be left alone.

If in such cases there is evidence of injury to the kidney the parietal wound down to the kidney should be excised, the organ investigated, and any foreign body removed.

7. In cases in which the thoracic wound is situated low down on the lateral or antero-lateral aspect of the chest access to both supradiaphragmatic and infradiaphragmatic areas may be obtained by cutting through the costal arch after the manner of Bland-Sutton, Duval, Bérard, etc.

8. If an abdomino-thoracic injury has been approached from the abdominal aspect, it is important not to waste time trying to complete a difficult suture of the diaphragm in a critically ill patient, unless the aperture in the midriff is so large that immediate or early herniation of the abdominal contents is certain to occur.

Prognosis

In one large series from the war of 1914-18 it was found that operated abdominal cases (excluding abdomino-thoracic injuries) gave a recovery rate of 51%; while abdomino-thoracic wounds operated upon produced a recovery rate of 44.5%. But some series of abdomino-thoracic cases in the casualty clearing zone gave a more favourable recovery rate than this: in the autumn of 1918 it attained the figure of 66.6% in the hands of certain surgeons in the British Fourth Army; and some operators (Jack Anderson, Saint) who were perhaps fortunate in their percentage of right-sided cases saved as many as 79 and 80% of their patients (Gordon-Taylor, 1939).

Wounds that traverse the diaphragm from below upwards are said to be more grave than those in which the missile has passed from thorax to abdomen. In the former group an intestinal injury increases the seriousness of the prognosis.

The mortality figures attending the surgical treatment of injury in one campaign are scarcely comparable with those of another war at a different period, in which the type of enemy missile, the degree of explosive force, the facility of transport from the field of battle, and other circumstances differ; but the steady improvement in the results of abdomino-thoracic cases from year to year in the war of 1914-18 was a memorable triumph of surgery.

NON-PENETRATING INJURIES OF THE ABDOMINO-THORACIC ZONE

In contrast to the last great European conflict non-penetrating injuries of the abdomino-thoracic zone do not play the same small and infrequent part in war surgery to-day; the thoracic and abdominal viscera, solid and hollow, may be damaged under a variety of circumstances in "modern" warfare. Fragments or masses of timber, stone, or metal may be confusedly hurled with explosive force against chest and abdomen in the indiscriminate air bombardment of civilians; women and children have suffered "crush" contusions of thorax and belly from falling masonry and iron girders; visceral damage above and below the midriff may result from the blast of detonating bombs, mines, or other high-explosive machinery in close proximity to men and women of every age, condition, and calling, for in "total war" none can claim privilege or sanctuary. Nor does the diaphragm remain immune when chest and belly suffer violence.

Effects of Rupture of the Diaphragm

Rupture of the diaphragm may be produced by the passage of a heavy vehicle obliquely across the abdomino-thoracic zone, by the crushing force of buffer accidents, or from the demolition and collapse of buildings. An excellent account of the injury has been given by Gordon Bryan (1922), who points out that the lesion is not very rare in children and adolescents, whose chest wall is more elastic and can be compressed to a considerable degree without fracture. In some cases costo-chondral dislocations may occur, as shown by some displacement of the

chest wall and a clicking sound on respiration; at other times the diaphragm is torn by stretching or bursting, or may be lacerated by a fractured rib.

The convexity of the diaphragm is the part that usually suffers; on the rarest occasion the crus is damaged. The left side of the midriff is that most prone to rupture under such conditions of violence; the liver is more likely to bear the brunt of an injury due to compression of the right side, yet the right dome of the diaphragm has been torn, with a resultant haemothorax and even the escape of bile into the pleura. The diaphragmatic lesion may readily be overlooked in the presence of cataclysmic intraperitoneal bleeding. Rupture of the diaphragm produces shock, dyspnoea, and a disordered action of the heart. The pulse is rapid and irregular; the breathing is almost entirely thoracic, but diaphragmatic respiration may be restored after a day or two by the plugging of the rent by omentum, spleen, etc.

Dyspnoea and tachycardia occur with herniation of the stomach into the chest; and severe pain with rigidity of the left hypochondrium is complained of, and also pain in the neck or shoulder from irritation of the diaphragmatic peritoneum. Thoracic distress is produced after meals; radiological examination will readily establish the diagnosis.

Rupture of one or both crura of the diaphragm may occur as the result of a subparietal injury. Bryan (1922) records a rupture of the left crus associated with a compression fracture of the eleventh and twelfth thoracic vertebrae resulting from a fall of thirty feet on to the pavement in a sitting posture. The intraperitoneal haemorrhage, which suggested a ruptured abdominal viscus, determined a laparotomy.

Severe crushing injuries of the chest have been known to produce a *tearing lesion of the mediastinum*, even when there has been no fracture of ribs or sternum. This has been found associated with a large retroperitoneal haematoma from damaged vessels in the root of the mesentery (Säuerbruch and O'Shaughnessy, 1937). The displacement of the mediastinum may suggest the diagnosis of a tension pneumothorax; investigation of the intrapleural pressure and the use of radiography should prevent unnecessary and harmful surgery.

The diagnosis and treatment of *subparietal injuries of the spleen* are well established and require no comment. For similar *hepatic injuries* thorough surgical exposure is necessary. Subparietal or non-penetrating *injuries of the kidney* are generally isolated lesions, and most often are mere contusions of the organ without rupture of the renal capsule. In injuries of the abdomino-thoracic zone which menace life concomitant damage rather than the renal lesion demands most attention. The chief indications for early surgical intervention on the kidney are: (1) cataclysmic haemorrhage; (2) persistent pain and aggravation of the clinical symptoms and signs of renal injury during the first forty-eight hours, especially increasing tenderness in the iliac fossa and hypogastrium with spasm of the psoas muscle; such clinical phenomena denote a spreading extravasation of blood, urine, or both. The size of any perinephric swelling that may be detected is some criterion of the degree of kidney injury. An appeal may also be made to urographic methods of diagnosis.

It should be remembered that early exploration in a doubtful case will render this service to surgeon and patient—that a deliberate conservative resection of the damaged portion of the kidney can be considered at a period when urgency is not so extreme that only a hurried total nephrectomy can save life. In arriving at a decision as to the exact technique to be employed in a case, it behoves the surgeon to remember the slogan: "First save the patient, and secondly, if possible, save the kidney."

Sequelae of Rupture of the Diaphragm

Other effects of injury of the diaphragm include: (1) the lodgment of fragments of foreign bodies in the diaphragm itself; (2) diaphragmatic hernia; (3) scarring and adhesions.

The most dramatic of these sequelae is *diaphragmatic hernia*; this may develop immediately or soon after the injury; on the other hand symptoms are commonly delayed for several months or longer. Traumatic diaphragmatic hernia rarely produces symptoms on the right side, and at most contains only a small piece of omentum in that situation. Only very rarely does a rent occur between the abdomen and the pericardium. The left dome is that most often implicated, and the herniated viscera include the omentum, stomach, transverse colon and splenic flexure, spleen, and jejunum; it is stated that the pancreas, liver, and duodenum are only very rarely herniated.

Such cases are best operated upon by a thoracic approach: an incision is made in the eighth or ninth intercostal space; the resection of adjoining ribs may also be required. Positive-pressure anaesthesia is employed; the phrenic nerve is crushed as it lies upon the pericardium before entering the diaphragm. The extruded viscera are first separated from the edges of the diaphragmatic aperture and thoracic structures, and are then coaxed into the coelom.

The repair of the rent in the diaphragm may be difficult if the muscle has undergone fibrotic or atrophic changes. Strong silk is employed: in some cases fascial grafts seem to give greatest promise of permanent closure. In others the gap can only be closed by suture of the liver, spleen, or stomach to the margin of the opening. In cases of some duration the stomach may have undergone hypertrophy and dilatation, and a laparotomy may also be required to anchor it satisfactorily inside the abdomen. The performance of a temporary gastrostomy in some such cases has proved of inestimable value in my hands. The complication of *strangulation* in connexion with a *diaphragmatic hernia* is fraught with the gravest peril. The colon is the segment of the alimentary canal most frequently obstructed; sometimes the jejunum is the part compressed; very rarely is the stomach strangled.

A correct diagnosis is not often attained unless there is previous knowledge of the existence of a hernia; but where such knowledge is lacking an antecedent history of a severe injury in the abdomino-thoracic zone, a previous complaint of post-prandial discomfort and even dyspnoea, hiccup, precordial oppression, and pain in the shoulder, together with the evidence of competent radiology, should enable a pre-operative diagnosis to be made.

The symptoms of strangulation usually supervene suddenly; the baleful results of operation for this disastrous complication constitute a powerful argument for early operation upon traumatic diaphragmatic hernia, at any rate if the symptoms to which it gives rise show any evidence of worsening. The transthoracic approach should be employed.

Familiarity with the transdiaphragmatic approach to the vault of the abdominal cavity has been turned to useful purpose by thoracic surgeons in the operation of cardio-omentopexy (O'Shaughnessy, J. B. Hunter). The successful operation by Pierre Duval (1918) for the extraction of a bullet from the suprahepatic portion of the inferior vena cava, in which the sternum was split and the diaphragm divided backwards to the caval opening still remains to my mind the most brilliant operation in the history of surgery.

Even the general surgeon avails himself of an abdomino-thoracic approach in operations for the surgical removal of

malignant disease of the lower end of the oesophagus or the cardiac orifice of the stomach.

No reference to the employment of the sulphonamide derivatives has been made in this communication; this is now a routine measure, and its value is incontestable. The severe shock which appears so constant in wounds of the trunk in this war demands a bountiful supply of blood and plasma, which on occasion is preferably introduced rapidly into the veins of the recipient.

In this war abdomino-thoracic injuries due to fragments of high explosive are probably less grave than those due to crushing force or the effects of blast.

REFERENCES

- Bryan Gordon (1922). *Brit. J. Surg.*, 9, 119.
 Duval, Pierre (1918). *War Wounds of the Lung*. Bristol.
 Edwards, A. Tudor. Memorandum, E.M.S./Gen./326, p. 10.
 Gordon-Taylor, G. (1919). *Brit. J. Surg.*, 7, 142.
 — (1939). *Abdominal Injuries of Warfare*, p. 13. John Wright, Bristol.
 Jolly, Douglas (1940). *Field Surgery in Total War*, p. 211. Hamish Hamilton, London.
 Lockwood, A. L., and Nixon, J. A. (1918). *British Medical Journal*, 1, 105.
 Nixon, J. A. (1919). *Ibid.*, 1, 399.
 Saint, Charles, quoted by Gordon-Taylor (1939). *Loc. cit.*, p. 59.
 Sauerbruch, F., and O'Shaughnessy, L. (1957). *Thoracic Surgery*, p. 234. Edward Arnold, London.
 Wallace, Sir Cuthbert (1922). *Official History of the War: Medical Services*. Surgery, 1, 567.

PROTECTING MEDICAL SUPPLIES AGAINST CONTAMINATION BY GAS

The problem of protecting medical supplies against contamination by poison gas is analogous in many respects to that of protecting foodstuffs, and doctors and hospital authorities will therefore get much help from the handbook issued by the Ministry of Food, *Food and its Protection against Poison Gas*. Some medical supplies, however, present special problems of protection and decontamination, and the Ministry of Health's circular (No. 2385) to local authorities and hospitals contains the following advice and suggestions.

The normal conditions of storage of medical and surgical supplies in suitably warmed and ventilated buildings should provide for most articles all that is practicable in the way of protection against liquid contamination. Some additional protection can also be afforded by using existing stocks of waterproof sheeting, batiste, cellophane, or other impermeable material if they are not required for immediate use as a covering for unprotected stores or stores whose nature or wrapping does not of itself provide protection.

Surgical dressings, cotton-wool, bandages, etc., wrapped in paper, cardboard, or cartons.—These would not become contaminated by exposure to vapour providing the wrappings are undamaged. Traces of vapour can, if necessary, be removed from such materials by opening them out and airing for an hour or so in a clean, well-ventilated warm atmosphere. Sterilization might be undertaken where the means are available. If contaminated by liquid it is extremely doubtful whether any decontamination treatment would be satisfactory and destruction by burning may be unavoidable.

Drugs and medicines in metal containers.—If adequately sealed no contamination of the contents can occur. If contaminated by liquid the containers should be decontaminated by repeated swabbing with solvent such as paraffin, petrol, etc., and airing; the corks, etc., unless protected by metal or other seal, should be destroyed and replaced by new ones.

Rubber materials, surgeons' aprons, waterproof sheeting, hot-water bottles, etc.—Vapour contamination will, in general, not be serious and could be removed by airing. If contaminated by liquid, decontamination by immersion in boiling water for a period of one to three hours, depending on the thickness of the rubber, would be essential.

Gum elastic catheters, surgeons' gloves, fine rubber drain sheeting, etc.—The finest grades of rubber materials are unlikely to stand boiling, and if contaminated by liquid should be destroyed. Covering to prevent such contamination is therefore most desirable.

Wooden furniture, bedding, etc.—The decontamination treatment for such articles is detailed in *A.R.P. Handbook No. 4*. In the case of bedding in general and mattresses in particular treatment in a steam disinfectant is the only practical process.

Metal instruments and utensils.—These, depending on their nature, can be decontaminated by immersion for thirty minutes in boiling water or repeated swabbing with solvent.

MEDICINAL HERBS IN WARTIME

The Minister of Health recently appointed an expert committee "to review the present and future requirements of vegetable drugs in the light of Empire consumption and trade, and facilities for cultivation within the Empire; to consider the steps which should be taken to secure organization of cultivation and collection; and to make recommendations to the Minister." This "Vegetable Drugs" Committee has now presented an interim report and recommendations. The report differentiates between two groups of drugs—a "long-term" and a "short-term" group.

Long-term Group of Drugs

It is unlikely that any long-term policy will materially affect the supply of these drugs during the present emergency, as the latent period between initiation of schemes and production for the market is so long. One primary essential for a long-term policy is an assured and steady market and remunerative prices for the producers within the Empire, particularly in respect of drugs requiring many years of experimental cultivation and considerable financial outlay before regular supplies can be available. The establishment of such stable conditions would probably necessitate (a) financial assistance to growers in the Colonies and Dominions over a period of years, and (b) protective tariffs against supplies from pre-war producing countries. The present difficult economic and political conditions and the impossibility of knowing or forecasting the economic circumstances and political considerations that may operate at the end of the war make it impracticable and inadvisable at the moment to formulate any proposals involving such questions. The committee strongly recommends that the position should be further considered as soon as possible after the end of the war.

Short-term Group

Here the position is quite different. Supplies can be rapidly created and producers can recoup themselves during the war. After the war stocks may come on the market from countries now cut off from international trade, which may create serious difficulties for growers in the United Kingdom who may have considerable stocks on hand at the cessation of hostilities. Therefore, so far as production in the United Kingdom is concerned, it is suggested that H.M. Government might consider safeguarding the position of persons who have undertaken the cultivation of drugs, in the same way as is being done in other branches of agriculture. It is hazardous to grow these plants in climates or soils whose suitability has not been clearly tested by experience, and it is preferable to expand production in areas where the plants are already grown and collected. Much could be achieved during the war by intensifying the collection of wild plants and developing the use of related species or substitutes.

Production of Vegetable Drugs in the U.K.

The committee considers that attention should be concentrated on the following drugs:

Agar	Fragot	Psyllium
Dill	Male fern	Sphagnum moss
Belladonna (leaf and root)	Liquorice root	Stramonium
Irish moss	Hyoscyamus	Dandelion root
Colchicum	Peppermint	Valerian
Digitalis		

Arrangements have been made by the Ministry of Health for regular growers in the United Kingdom to extend the areas under cultivation of certain extremely important drugs, which with other steps taken to secure supplies, ensures that the wartime requirements of these are safeguarded.

The committee points out that while the development of organized collection of herbs growing wild is most valuable and should be encouraged, the growing of medicinal herbs is a highly specialized industry which is not without its dangers to the inexperienced and that therefore growing by amateurs is to be discouraged. Considerable experimental work is being done at research institutes and agricultural stations on strains, seed production, growing, and utilization of various drugs, the results of which may be important. Steps have already been taken to encourage methodical collection, proper drying (which is most important), and marketing of drugs growing wild in this country.

Cultivation, Collection, and Marketing

During 1940 the Ministry of Health was engaged in correspondence with amateur growers and collectors regarding cultivation, collection, and marketing of vegetable drugs, and held several conferences with trade representatives. The Department also sponsored, in co-operation with the Ministry of Agriculture and Fisheries and the Board of Education, a