proximal portion of the excretory apparatus. Even after 84 hours the kidneys cast an abnormally dense shadow, indicating that some of the dye was still being retained.

Volhard's test was carried out on Sept. 19, with the following result: Specific gravity of urine during 24 hours with forced fluids: minimum reading 1000. Specific gravity of urine during 24 hours without fluids: maximum reading 1028.

Discussion

Prominent in the consideration of the treatment of this case was the assumption that the renal tubules were blocked with sulphapyridine crystals. With the aim of promoting active hyperaemia of the glomeruli and increasing secretion it was decided to employ inductotherm (short-wave) treatment. This method of heating deep tissues, the most efficient in use, has been applied previously in cases of renal disease (Stark, 1939), but we have been unable to find any reference to its employment in anuria resulting from the use of the sulphonamide series of drugs. The results in this case indicate that it may be of considerable value in the treatment of cases in which the condition is due primarily to mechanical obstruction of the renal tubules. It is unlikely that this treatment would be equally effective when glomerular damage is present.

It would appear that in cases with oliguria or anuria arising from the use of the sulphonamides this method of treatment is worthy of trial before submitting the patient to ureteric catheterization and lavage of the renal pelvis, a procedure which may be inadvisable when an infection of the urinary tract is present. Catheterization will of course be required when the ureteric lumen is completely occluded.

The x-ray findings in this case indicate the presence of intrarenal changes, gradually diminishing during convalescence. It is therefore apparent that, even though ureteric catheterization may have to be carried out, short-wave therapy may be a useful adjunct in clearing the renal tubules to the greatest extent at the earliest opportunity.

We wish to express our thanks to Brigadier C. H. Stringer and Lieut.-Col. J. W. Craven, R.A.M.C., for permission to publish this article.

REFERENCES

- Antopol, W., and Robinson, H. (1939). *Proc. Soc. exp. Biol.*, N.Y., **40**, 428.
 Backhouse, T. C. (1939). *Lancet*, **2**, 736.
 Gross, P., Cooper, F. B., and Lewis, M. (1939). *Proc. Soc. exp. Biol.*, N.Y., **40**, 448.
 Lebel, H., Schroeder E., and Simesen, M. (1940). *Acta med. scand.*, **105**, 395.
 Leedham-Green, J. C. (1941). *British Medical Journal*, **1**, 586.
 Pepper, D. S., and Horack, H. M. (1940). *Amer. J. med. Sci.*, **199**, 674.
 Plummer, N., and McLellan, F. (1940). *J. Amer. med. Ass.*, **114**, 943.
 Scurry, M. M., and Wittenborg, M. H. (1940). *Ann. intern. Med.*, **13**, 2164.
 Stark, G. W. (1939). *Arch. phys. Therapy*, **20**, 94.
 Stryker, W. A. (1940). *J. Amer. med. Ass.*, **114**, 953.
 Tsao, Y. F., et al. (1939). *Ibid.*, **113**, 1316.

CEREBROSPINAL FLUID IN ACUTE ANTERIOR POLIOMYELITIS

BY

ARTHUR L. EYRE-BROOK, M.S., F.R.C.S.

Assistant Surgeon to the Bristol Children's Hospital

Paralysis in infancy does not constitute infantile paralysis (acute anterior poliomyelitis), and we need to be in a position to diagnose this, "the saddest of diseases," when we are confronted with paralysis in childhood. The nature of the paralysis and the tenderness of the muscles are characteristic features in this disease, in which the diagnostic picture with its pre-paralytic stage and "back signs" can be very typical. Many cases, however, are atypical; how are we to diagnose them?

The Literature on Diagnosis

H. Howton Merritt and Frank Fremont-Smith (1937) state that before the onset of paralysis the diagnosis of anterior poliomyelitis cannot be made with certainty on the basis of the examination of the cerebrospinal fluid because numerous diseases of children and young adults are accompanied by similar cerebrospinal fluid changes. Fortunately, about the time of the onset of the paralysis changes begin to occur in the cerebrospinal fluid which are very characteristic, but these can only be fully appreciated if two or three specimens are obtained during

the first three weeks of the illness. These changes are: (1) an early rise in cell count, falling to normal by about the end of the second week of the illness; (2) a protein content which is almost normal until some days after the onset of paralysis, and which rises to a maximum in the third week. These very characteristic changes were first demonstrated by Peabody, Draper, and Dochez in 1912, and have often been confirmed since then.

As to the nature of the cells which cause the early rise, Howton Merritt and Fremont-Smith state that 80% of their cases showed a lymphocytic predominance within five days of the onset of the paralysis, although the early counts might consist largely of polymorphonuclear cells. Similarly, Peabody, Draper, and Dochez state that "polymorphonuclear cells, except in small numbers and probably due for the most part to a slight admixture of blood, were, except in a very few instances, absent after the first week of the disease."

This glance at the literature gives us a clear idea of the possibilities and limitations of the findings in the cerebrospinal fluid as an aid to diagnosis.

Findings in a Series of Seven Cases

This short paper is designed merely to stress the importance of these known facts by reporting the findings in seven cases of infantile paralysis admitted to the Bristol Children's Hospital during a small epidemic which occurred in the autumn of 1938. It was found that the cerebrospinal fluid had been examined on various occasions between the fifth and twenty-first days of the illness. The paralyzes had in every case developed between the second and seventh days, and the fluid was in all cases examined during the paralytic stage.

In those cases in which the fluid was examined within the first twelve days the cell counts were high, reaching 50 in two cases, while in those examined after the second week of the

Chart showing the cerebrospinal fluid findings in the seven cases described. o = Cell count. • = Protein content. (Cerebrospinal fluid from A.P. examined on two occasions.)

illness the cells never exceeded 6. No fluid was examined before the fifth day of the illness, which explains why no very high cell counts were recorded.

The protein content was below normal in two fluids examined on the seventh and tenth days, but there was one specimen on the fifth day which gave a reading of 70 mg. per 100 c.c.m. Three estimations of 300 mg. and one of 200 mg. were, however, obtained during the third week, revealing the magnitude of this late protein rise. On the twenty-first day a specimen showed a cell count of 5 and a protein content of 70 mg., suggesting that at this stage the protein content was also on the wane.

One case was subjected to two lumbar punctures, and the cerebrospinal fluid findings were: on the seventh day—cells 50, protein 80 mg. per 100 c.c.m.; on the fifteenth day—cells 5, protein 200 mg. Such findings are diagnostic of acute anterior poliomyelitis.

It must be noted that a single specimen of fluid taken towards the end of the second week may give a reading which will show little that is abnormal, because the cell count will

be returning to the normal level while the protein rise may hardly have begun. This observation lays stress on the importance of obtaining two specimens of fluid from each case, preferably at the end of the first week and in the middle of the third week. Such routine examinations should enable a case of acute anterior poliomyelitis to be picked out with ease from among the paralyses of childhood and young adult life.

Summary

The findings in a small series of cases in an epidemic of acute anterior poliomyelitis are reported. These show the typical early rise in cell counts and the late rise in protein content. The cases are recorded in detail on the chart.

An appeal is made for the taking of more than one specimen of cerebrospinal fluid in a case suspected of acute anterior poliomyelitis, and for the interpretation of the findings in the light of the changing picture presented in this disease.

It is stressed that a single specimen at the end of the second week may give findings approximating to the normal, as the cell rise is almost at an end and the protein rise may hardly have begun.

I would like to acknowledge the help of Prof. Perry, Drs. Birrell, Beryl Corner, and Lloyd Price, under whom these cases were treated and who called me in as their orthopaedic colleague; and of Prof. Davie, who was at the time professor of pathology at the University of Bristol, and whose staff carried out all the cerebrospinal fluid examinations. My thanks are especially due to Prof. Perry, who has stimulated me to publish this series from a small epidemic in an attempt to widen the recognition of the entirely typical picture here presented.

BIBLIOGRAPHY

- Gordon, R. G., and Brown, M. Forrester (1933). *Paralysis in Children*, London. Hobhouse, N. (1932). *Nervous Disorders of Infancy and Childhood*, London. Merritt, H. H., and Fremont-Smith, F. (1937). *The Cerebrospinal Fluid*, Philadelphia. Peabody, F., Draper, G., and Dochez, A. R. (1912). Monogr. Rockefeller Inst. med. Res., No. 4, New York.

TETANY IN BLOOD DONORS

BY

W. FITZGERALD FRAZER, M.R.C.S., L.R.C.P.

Surgeon Lieutenant, R.N.V.R.

AND

F. S. FOWWEATHER, M.D., M.R.C.P., F.I.C.

Reader in Chemical Pathology, University of Leeds

The removal of blood from donors by the methods now in use has little effect on the great majority of them. A small proportion—about 5% according to Brown and McCormack (1942)—show certain vasomotor phenomena, commonly referred to as “faints.” But in addition to these manifestations we have found that a few donors have developed unmistakable attacks of tetany, with typical spasms of hands and feet and positive Chvostek's sign. The number of donors affected in this way was small, amounting to 1 in every 1,000 on the average, but the condition is nevertheless of some importance, and we are not aware that it has been previously reported.

It is, of course, highly desirable that any untoward reaction occurring in a blood donor should be treated as promptly and efficiently as possible, not only for the sake of the subject of the attack but also, in centres where a number of donors are bled in the same room, to prevent the attention of others being drawn to the occurrence; for the observance of unusual reactions is apt to have a most disturbing effect on other donors, and the frequency and nature of the reactions tend to be greatly exaggerated by them. We have therefore attempted to ascertain the nature of the tetany we have observed and to apply a rational treatment.

Determinations of serum calcium in the first 2 of our cases proved that we were not dealing with low-calcium tetany; and gastric tetany was excluded by the entire absence of any history of attacks of vomiting. It was therefore concluded that the condition was probably a hyperventilation tetany, and in subsequent cases the serum inorganic phosphorus was determined, as well as calcium, and the appropriate treatment for hyperventilation tetany was instituted. This consists in supplying to

the patient air enriched with CO₂. In only one case was a cylinder of CO₂ immediately available; in the others the simple method of placing a paper bag over the patient's nose and mouth, and so causing him to rebreathe his own expired air, was adopted. Details of the cases are as follows:

Case Histories

Case 1.—Woman aged 42. First donation. Blood taken, 60 c.cm. Tingling and numbness noticed first in hands and feet, followed by carpopedal spasm and laryngismus stridulus. Attack lasted 25 minutes. No history of similar attacks. Serum calcium in specimen taken during the attack, 10.8 mg. per 100 c.cm.

Case 2.—Woman aged 30, very obese. First donation. Blood taken, 540 c.cm. Tingling and numbness, followed by carpopedal spasm and positive Chvostek's sign. Attack lasted 16 minutes. History of similar attack some years ago, after injection for varicose veins. Serum calcium in specimen taken during the attack, 11.2 mg. per 100 c.cm.

Case 3.—Man aged 34. Third donation. Blood taken, 540 c.cm. Full carpopedal spasm, laryngismus stridulus, positive Chvostek's sign. Treated by paper-bag method, following the application of which the spasm began to pass off almost immediately, and symptoms had disappeared within 10 minutes. Serum calcium in specimen taken during the attack, 11.2 mg.; inorganic phosphorus, 3 mg. Corresponding figures for specimen taken at the end of the attack were: calcium, 12.2 mg.; inorganic phosphorus, 2.2 mg.

Case 4.—Man aged 26; suffers from migraine; very apprehensive and nervous. First donation. Blood taken, 540 c.cm. Carpopedal spasm, laryngismus stridulus, positive Chvostek's sign. Spasm passed off very quickly after administration of air containing about 12% CO₂ through Minnitt's apparatus. Serum calcium in specimen taken at the end of the attack, 11.2 mg.; inorganic phosphorus, 1.9 mg.

Case 5.—Woman aged 32. First donation. Blood taken, 540 c.cm. Carpopedal spasm and laryngismus stridulus very marked; positive Chvostek's sign. Treatment by paper-bag method relieved the spasm, but patient would not co-operate, and persisted in hyperventilating; carpopedal spasm again supervened, lasting for a further 5 minutes. Serum calcium at the end of the attack, 11.3 mg.; inorganic phosphorus, 1.3 mg.

Case 6.—Woman aged 51. First donation. Blood taken, 400 c.cm. Tingling and numbness of hands complained of, and bleeding was stopped. Spasm developed in hands, but in feet there was only tingling and numbness. Treated by paper-bag method; symptoms disappeared after 10 minutes. Serum calcium at the end of the attack, 10.5 mg.; inorganic phosphorus, 3.1 mg.

Case 7.—Woman aged 23. First donation. Blood taken, 540 c.cm. Marked carpopedal spasm and laryngismus stridulus, with positive Chvostek's sign. There was very noticeable overbreathing. Treated by paper-bag method; attack terminated after about 7 minutes. Serum calcium at the end of the attack, 10.8 mg.; inorganic phosphorus, 3 mg.

Discussion

The first point to be noted is that the onset of the attacks is not wholly dependent on the amount of blood withdrawn, for while in most cases the attack occurred after the full quantity of blood had been taken, in one it supervened after the removal of only 60 c.cm., and in another after 400 c.cm. The next point is the preponderance of females among the cases observed. This is in agreement with the general experience that spontaneous hyperventilation tetany occurs mainly in females.

So far as the blood chemistry is concerned, the findings agree in the main with those of Fowweather, Davidson, and Ellis (1940) in spontaneous hyperventilation tetany, and of Haldane, Wigglesworth, and Woodrow (1924) in experimental hyperventilation tetany. The latter authors found that on every occasion the blood inorganic phosphorus fell below 1.25 mg. per 100 c.cm.; moreover, this fall and the subsequent rise to normal values after the attack are rapid, and consequently minimal concentrations are present for only a very short time. Fowweather *et al.* found, in specimens of blood taken at the end of the tetanic attack, that the serum phosphorus was reduced (in 2 cases to only 1 mg. per 100 c.cm.), while the serum calcium showed a high-normal or even a definitely raised value. Other authors have mostly reported normal or slightly raised calcium figures, with low-normal phosphorus figures (2 to 3 mg.) in specimens taken during attacks. Fowweather *et al.* argued that the greater changes shown by them, compared with those of