BRITISH MEDICAL JOURNAL

LONDON SATURDAY JULY 5 1941

THE TREATMENT OF BURNS AND WOUNDS BY THE ENVELOPE METHOD

BY

JOHN BUNYAN, *Surgeon Lieut.-Commander (D), R.N.V.R.*

(WITH SPECIAL PLATE)

Introductory Remarks

The outbreak of war has naturally resulted in much research into the treatment of burns and wounds. Experience has led to the modification and even rejection of some previously accepted methods of treatment. The most notable incident has been the discovery of the disasters that may follow the application of tannic acid to the hands and face. Some of the therapeutic advance up to date has been of an empirical, trial-and-error type, and has left much room for improvement. The scientific theoretical aspect of the question is still, it is suggested, capable of further development.

My interest in inflammatory disorders and wounds of the mouth led me to form certain hypotheses and opinions on the problem of the local treatment of inflammatory conditions in general. My theoretical views were as follows:

First, sound healing of the tissues is assisted by gentle but thorough cleansing, with removal of dead tissue and avoidance of irritant applications, together with every encouragement to the physiological outpouring of serum and leucocytes into the injured area. This assistance is, of course, strengthened by any bactericidal treatment that can be given without damage to living cells and by the prevention of secondary infection.

Secondly, it is desirable that any treatment given should be painless, and, if possible, alleviate existing pain, and that it should reduce to the minimum the necessity for general anaesthesia. It would be a great advantage if the local treatment permitted the active and passive movements of the injured part at all times.

Thirdly, a local treatment of burns should have the following qualities: it should not aggravate primary shock; it should diminish and inhibit the loss of body fluid from the burnt area; it should reduce toxæmic shock from the absorption of protein breakdown products and bacterial toxins; and it should disinfect without irritating or damaging the healing tissues.

In the past less success than was hoped for has followed the treatment of burns and wounds because irritating, destructive antiseptics were used, which killed tissue as well as organisms. At the same time the type of frequently changed dressing, conventionally applied, gave rise to pain on removal, to destruction of newly formed tissue to which it was adherent on each removal, and to immobility and disuse of the part to which it was applied. Frequent changing of a dressing introduced the inescapable danger of secondary infection from such sources as droplets from the nose and mouth of other individuals or from ward dust.

To obtain improvements in the local dressing of burns and wounds it is therefore desirable to employ a dressing which may be kept applied without changing for a considerable period—preferably during the whole course of treatment. The dressing should be flexible, frictionless, bland to the tissues, and, if possible, transparent, in order that progress may be constantly observed. The dressing should combat bacterial contamination and be watertight, so that fluids applied to the injured area would be confined within bounds.

Apart from avoiding the necessity for changing, the dressing should permit free movement if this is desirable, and permit constant drainage and cleansing of the affected area.

The Therapeutic Agent

In attempting to devise an improved form of dressing to meet the above-mentioned requirements, the form of solution to be applied to burns and wounds was attended to first. It was desired to find one which was non-toxic, non-irritant, capable of dissolving dead organic matter, able to encourage transudation of serum and cells, and to exert some bactericidal power in concentrations that would be harmless to healing tissue, and which would be easily obtainable anywhere and at any time.

From personal clinical experience, and from literature on the subject, it was decided that sodium hypochlorite solutions most fulfilled these desiderata, and that the hypochlorite solution produced by electrolysis of sodium chloride was the best preparation, having the least irritant action. The following are a few important points in the manufacture of hypochlorite solutions.

First, sodium hypochlorite solution can be made by a chemical process. Chlorine gas can be passed through dilute caustic-soda solution at suitably low temperatures. The reaction is as follows:

In solutions prepared this way sodium hypochlorite and sodium chloride will usually be found in equal quantities, but there will always be an excess of caustic soda remaining in solution.

Sodium hypochlorite solution can also be made in an indirect chemical manner by the use of bleaching powder. The danger of causticity is present in this method also, as exemplified in the case of liq. sod. chlorinatae (B.P.C.). In Dakin's solution causticity is reduced by the addition of boric acid, but some of this is absorbed by the tissues, so that free soda is always present. Calcium hypochlorite solution made from bleaching powder is also combined with boric acid to form eusol.

In all solutions made from bleaching powder there is one serious disadvantage—namely, the variability in composition of the bleaching powder. Careful analysis is necessary before a final product of definite strength and alkalinity is obtained. It is impracticable to carry out such an analysis every time one of these solutions is made up in a dispensary or hospital.

Secondly, sodium hypochlorite can be manufactured electrolytically from the electrolysis of brine. The reaction can be expressed in the following equations:

There is evidence to suggest that the increased efficacy of electrolytic sodium hypochlorite as compared with the chemical form is due to the presence of ozone. Electrolytic sodium hypochlorite has an oxidizing action on organic matter. Glycerin, for example, can be converted by successive stages to oxalic acid. Proteins, pus, mucus, etc., are dissolved and converted to inoffensive products. In the case of proteins chloramines are formed, which in themselves perhaps have some degree of germicidal action.

Unlike any other antiseptic, electrolytic sodium hypochlorite is completely non-toxic. HOCl gives either of two actions—oxidizing or chlorinating. The end-product from oxidizing is NaCl, and from chlorinating H_2O . Electrolytic sodium hypochlorite is thus the nearest approach to a physiological and natural cleansing medium. On living tissue, in proper dosage, the effects are beneficial and stimulating to growth, in contrast with the destructive action on necrotic tissue. This automatic selectivity—the property of differentiating between dead organic matter to be dissolved* and live tissue to be repaired—is a most interesting one.

Electrolytic sodium hypochlorite does not dissolve necrotic matter quite so well as chemical hypochlorite because the latter contains alkali, but this difference is unimportant compared with the freedom from causticity of the former.

The use of hypochlorites has been criticized because of the process of inactivation: when brought into contact with dead organic matter the hypochlorite is broken down and therefore cannot continue to exert a bactericidal action. It is partly in this property that its value lies, and it is only necessary to provide a continual supply of solution for a given period to take the utmost advantage of it. In order to get the greatest value from the solution it was considered important to use a large quantity of a low concentration rather than a small quantity of a high concentration.

The Dressing

Having decided to employ electrolytic sodium hypochlorite, the next step was to devise some method of applying the solution to the injured area in a simple manner and in such way that it could be left in contact with the tissues for as prolonged a period as desired. The allowance of adequate time to exert its full effect is important in treating sepsis with sodium hypochlorite.

A dressing made from rubber sheeting with tyre valves as inlet and outlet was tried on an old injury dating from the last war. The clinical result was excellent, but the rubber apparatus was by no means satisfactory. An injured limb was then put up in plaster, with a window of celluloid inserted over the wound, and hypochlorite solution was pumped in through a Higginson syringe three times a day for twenty minutes. Again the results were excellent clinically, but the method had many mechanical drawbacks.

It was at this time that we had the extreme good fortune to interest Mr. Wm. Stannard of Leek in our idea. He agreed to make an envelope of oiled cotton which would allow a fluid to be pumped in and let out while attached to the limb. The first one made was very crude, but was good enough to show that the idea had many possibilities. Shortly afterwards better envelopes were made, and these were tried out on several cases of burns and wounds. The results were such that encouragement was given to develop the idea, and during the past eight months Mr. Stannard has devoted himself entirely to the perfecting of the coated silk, so that it is now impermeable to electrolytic sodium hypochlorite, transparent, and absolutely bland to the tissues. The designing of the envelopes and the most efficient method of sealing to the limb also presented many problems that he has successfully overcome. The standard of perfection arrived at is such that it is now considered possible to release the envelopes to the profession with confidence.

It must here be stressed that the method is very new, and that far more time than six months is required in order to perfect not only the mechanical but the treatment side of the system.

Every facility was offered by Mr. Vaughan Hudson at the Middlesex Hospital, and Dr. Bruce Pearson at King's College Hospital. Mr. J. W. Hannay, at the London Hospital, took an active part in trying out the possibilities. At the same time I was given facilities at the Radcliffe Infirmary, Oxford, for the experimental side of the work. In order that all aspects of the subject might be investigated and any question answered the following team of workers was formed: Mr. Vollum, D.Phil., M.A., bacteriologist; K. O. Black, M.D., M.R.C.P., physician; R. G. MacFarlane, M.D., B.S., M.R.C.S., haematologist; Dorothy Russell, M.D., histologist; Mr. Manifold, B.Sc., Ph.D., biochemist; Prof. R. R. Macintosh, D.M., F.R.C.S., anaesthetist. I would like to acknowledge their helpful co-operation.

Description of Apparatus

(a) *Envelopes*.—The envelopes are simple covers designed to envelop any limb. Means of inlet and outlet are provided so that a lesion on any part may be irrigated by the insertion of a glass nozzle through any one of the openings. A simple adhesive seal is fixed on the proximal end of the envelope at the factory, so that by removing the protective gauze a watertight seal is produced by lightly pressing the adhesive to the limb, folding over the excess flaps, and covering with single-sided adhesive bandage. At present the following envelopes are in production:

		Infant	Small Child	Adult
Full leg ..	{ Circumference of thigh .. Length, groin to heel..	Up to 10"	10-14"	14-22"
		.. 15"	15-21"	21-33"
Foot and leg ..	{ Circumference (top) .. Length	.. 9"	9-13"	12-14"
		.. 14"	14-18"	18-23"
Knee	{ Length Circumference (top) (bottom)	12"	12"	12"
		13-16"	16-20"	16-20"
		10-13"	13-16"	13-16"
Elbow	{ Length Circumference (top) (bottom)	16"	16"	16"
		8-11"	11-14"	11-14"
		5-8"	8-11"	8-11"
		Infant	Large Child	Adult
Full arm ..	{ Circumference Length	Up to 6"	6-11"	11-14"
		.. 13"	13-24"	24-29"
Forearm & hand	{ Circumference Length	.. 6"	6-8"	8-13"
		.. 12"	12-17"	17-19"

* The word "dissolve" as applied to the removal of dead organic matter is not used in its true chemical sense, but is employed to describe the undoubted washing away of stale exudate and sloughs.

Envelopes for the head and a full bath are being developed, and will be available when sufficiently tested.

Different shapes and sizes are being tried out, and as the method develops it is hoped to be able to provide a simple means of effectively isolating any part of the body.

(b) *Other Apparatus :*

1. Special trolley for use in hospital.
2. Containers for hypochlorite solution, of glass or vulcanite.
3. Thermometer.
4. Rubber tubing and clips (sterile).
5. Glass cannula and vulcanite roses (sterile).
6. Higginson's syringe (sterile).
7. Waste container (bucket or bowl).
8. Silk sheeting.

(c) *Method of Application of Envelopes.*—See Treatment of Burns.

Technique of Treatment : (i) Preparation of Sodium Hypochlorite

Electrolytic sodium hypochlorite can be made with a small electrolytic cell, as used in modern water purification, producing three pints of 0.6% NaOCl (unstabilized) from 10% NaCl solution in six hours. With a larger plant considerable quantities of NaOCl can be produced as required. However, if this plant is not available, the simplest way to obtain as much sodium hypochlorite as required, in a standardized stabilized solution, was found to be to use "milton" solution, which is 1% NaOCl, standardized and stabilized, and which has been and is being kindly supplied by Messrs. Milton Proprietary Ltd. Since this concentration is a very convenient one for using as a stock solution, it has been decided to refer always to a 1% solution of sodium hypochlorite, electrolytically produced, as E.S.H., so that when we talk of a 5% solution of E.S.H. we mean 1 part of 1% electrolytic sodium hypochlorite to 20 parts of water. This 1% solution of E.S.H. has been used in all my experiments, and appreciation of the assistance given by the manufacturers must be expressed.

Various concentrations of E.S.H. were employed for different purposes. It should be remembered that solutions of E.S.H. above 5% in water are hypertonic, those of 5% almost exactly isotonic, and those below 5% hypotonic. The following guide to the concentration of E.S.H. for various purposes is given.

First, when gross sepsis is present, or dead tissue such as sloughs, scabs, etc., has to be removed, then E.S.H. in strengths up to 50% in water may be used. A concentration of 20% is a good average one. Hypertonic solutions assist the diminution of oedema.

Secondly, when the primary object of thorough cleansing has been attained, and daily irrigation is instituted, to remove stale exudate and to control infection, this daily irrigation should be carried out with a 5% solution. If the skin is very sensitive, or if healing is progressing rapidly, then the concentration may be reduced to 2½%. When there is no infection, or the infection is so slight as to be almost negligible, and stimulation of healing is the main objective, the 1% solution is adequate.

Temperature control is important. The solution should be kept at body heat. It should be made a rule that it be kept at 100° F., except that the initial irrigation done under analgesia or anaesthesia may be carried out at 110° F. Patients are extraordinarily sensitive to slight temperature variations, and great care should be exercised to ensure that the solution is at 100° F. exactly.

(ii) Treatment of Burns

The general treatment of shock, etc., need not be detailed here. It may be mentioned that when a general anaesthetic is necessary for the débridement and dressing of burns it need only be a light one. Various forms of anaesthesia and analgesia have been employed, including intravenous morphine, omnopon, and barbiturates. I have been impressed by the advantages of a vaporizer for delivering warm ether vapour of any desired concentration,

devised by the Nuffield Department of Anaesthetics, Oxford: anaesthesia, and even analgesia, have been most satisfactory, and the patients have had no ill effects from their use.

There is much confusion regarding the best method of first-aid treatment for burns, and guidance has hitherto taken the rather negative form of instruction to do nothing. Since some time may elapse before admission to hospital it is felt that the application of packs of E.S.H. to the burn is a considerable advantage—the relief of pain and the practical disinfection, together with prevention of fluid loss, are of great importance. Packs may be applied anywhere quite easily, and cannot possibly do any harm.

First-degree burns may be bathed in 50% E.S.H. for ten minutes, after which the concentration is lowered to 10% for ten minutes. The area is dried and then covered with adhesive plaster. Relief of pain is found to be immediate, and after twenty-four hours there will be little sign left of the burn. Second- and third-degree burns may be classed as either primary or secondary. Primary burns are those that arrive for treatment without having received any attention other than first aid.

It must here be noted that it has been found most difficult to decide at first whether a burn is a bad second or of actual third degree. There can be no doubt that many second-degree burns progress to third degree because of sepsis or trauma, and it is felt that the use of this method will enable treatment to be carried out without the possibility of any harm until the degree of the burn is established. By then any form of treatment may be undertaken without having run the risk of the catastrophes associated with the treatment of deeper burns.

A patient with a second- or third-degree burn is pre-medicated and anaesthetized as may be necessary, and local treatment is begun. When the blisters are unbroken the area is washed over on the surface with 20% E.S.H. at 100° F. Blisters should then be aspirated with a fine needle and some 10% E.S.H. injected into them. This should be left in for five minutes, and then withdrawn by syringe and needle, and the skin pressed down flat with a dry dressing. When the skin is broken the dead skin should be cut away under a stream of E.S.H., bathed for ten minutes in a 10% solution, and covered with a sheet of the special coated silk sealed at the edges with adhesive.

To deal with large burnt areas a suitable silk envelope, previously sterilized, is fitted. E.S.H. in the strength described is run through the envelope, washing all the affected area. The envelope is drained well and then inflated with oxygen, the inlets being sealed with adhesive tape or rubber-covered clips. Next day the irrigation routine begins: 5% E.S.H. is employed at 100° F., and is run through the envelope over the affected area for twenty minutes. This procedure is carried out three times a day, draining thoroughly for half an hour after the irrigation and then inflating with oxygen. A sheet of coated cotton is placed under the affected part to prevent wetting of the bedclothes, and the tube from the outlet is run into a basin, all this being explained in the directions for applying the envelopes.

In the case of hands the envelope may be filled with E.S.H. for the last ten minutes of irrigation to encourage the patient to move the fingers and wrist. A jet of E.S.H. from a Higginson syringe through the inlet opening may be used to remove sloughs and exudates.

In the case of third-degree burns of the hand and forearm an arm-bath is filled with E.S.H. solution up to a concentration of 20%. The more cleansing that has to be done the stronger should be the concentration. A jet from a Higginson syringe should be employed to scour-irrigate the limb, using the solution in the arm-bath. Dead skin should be removed almost to the margin of the burn. It should not

be removed right up to the edges, to its junction with normal skin, as this may result in subsequent pain. No scrubbing should be employed. Hosing with the Higginson syringe should be continued until spontaneous haemorrhage takes place and the burnt area becomes covered with a firm shiny fibrin coagulum. If profuse bleeding occurs no anxiety need be felt, since it has been found that it will cease spontaneously when the envelope is applied. Definite bleeding-points should be twisted or ligatured. Axillary depilation and nail-cleansing should be carried out.

The burned area is now covered with coated silk sheeting and dried outside this with sterile towels, the envelope being slipped into position over the limb and outside the sheeting. The latter is withdrawn from inside the envelope, the normal skin beneath the seal of the envelope is dried with aether meth., and the seal is closed. A small quantity of E.S.H. (of concentration previously described) is then run over the whole limb, the envelope is drained, oxygen is run in, and the inlets are sealed. Routine thrice-daily irrigation is now maintained for as long as necessary, as previously outlined.

These details apply equally to burns of the legs, but there are certain practical points of importance in addition so far as the legs are concerned. It is convenient to lay the patient on a length of coated cotton sheeting spread out on a table (e.g., an operating table), the sides of the sheeting being raised up by any suitable device. The lower end of the sheeting is formed into a trough leading to a receiver at the foot of the table. The table is slightly tilted to allow the fluid to run into the receiver. The whole of the affected area is then sprayed with E.S.H. of suitable concentration. The whole leg may be enclosed in an envelope, but if this is done care should be taken to see that the foot is thoroughly cleansed. In burns involving the upper thigh pubic depilation is necessary, and should there be no room between the perineum and the burn for the envelope to be sealed, then its use should not be attempted. Burns of this nature need treatment in a "perineal bath," which has proved of great value in the treatment of burns and wounds involving this area. Otherwise, primary third-degree leg burns are treated exactly as those of the hands and arms.

Secondary third-degree burns are those which have been subjected to treatment beyond that of mere first aid and which have failed to heal and have become infected. The technique of their treatment is similar to that of the primary type, but cleansing must be very thorough. When suppuration is present under the latter should be dissected away to the healthy margins. Any obviously necrotic subcutaneous tissue should also be removed. The dissection should be carried out under a stream of E.S.H. up to a 20% concentration, and a thorough scour-irrigation should be given for twenty minutes after the removal of the tan. The envelope is then applied and routine thrice-daily irrigation instituted as previously described. Healing usually proceeds rapidly in third-degree burns subjected to this treatment, and at the end of ten days, and sometimes earlier, any areas needing it can safely be skin-grafted.

Where the face and head are burnt, the initial cleansing is done in the normal manner, except that 10% E.S.H. is used. This is sprayed on through a vulcanite rose, with the patient bending his head over a wash-basin or bowl. If a patient cannot leave his bed he lies back with a silk sheet attached around his neck and beneath his head, forming a chute through the head of the bed into a pail. A cylindrical envelope which has slits for his eyes, mouth, and nostrils is then applied. In the two cases treated in this manner the patients have been free from pain, oedema of the eyelids has been reduced very rapidly, and the eyes have shown no signs of any irritation. Where necessary the eyes are washed with 1% E.S.H. solution, after which drops of sterile paraffin are inserted. Immediate skin-

grafting may be carried out in many cases or may be deferred for a few days to permit some granulation.

Burns of the limbs may be complicated by the presence of compound fractures. When this is the case it is suggested that treatment might be as described, and plaster or other splinting be applied outside the envelope. In some cases splinting is best fitted (if it is small) inside the envelope.

It has been noticed that most of the burns suffered by small children are extensive and are not capable of being treated in one or other of the limb envelopes. Small vests have been made for covering burns of the upper part of the body, and a nightgown of coated silk with hood has been fitted so that the whole of the child is enclosed, leaving sufficient length from the feet to form a trough over the end of the cot. Children can move freely in this, and are quite comfortable and free from pain. This device is very much in the experimental stage, however, but promises well.

(iii) Grafting

The envelope system of treatment has been successfully employed in the early preparation of an area for grafting. When a denuded area has progressed under treatment sufficiently to be ready for skin-grafting an irrigation is carried out and the envelope immediately removed. The whole area to be grafted is gently wiped over with gauze soaked in 1% E.S.H. Grafting is then carried out in the usual way. The skin-graft may be kept in position by a sheet of perforated coated silk stitched to the surrounding skin; a new envelope may now be applied, and after twenty-four hours have elapsed irrigation once daily with 1% E.S.H. can be resumed. In many cases the skin can be taken from the same limb and both areas healed in the one envelope. The system is not in any way meant to displace skin-grafting. It is left entirely to the surgeon in charge of the case to decide whether he will leave the area to epithelize or whether he will carry out an early graft. In any case, the use of the envelope will ensure the prevention of secondary infection.

(iv) Wounds and Lacerations

I have confined myself to the treatment of burns, but after watching the results of the treatment of various injuries and septic conditions by other workers have felt that the system will open up a new field of work in connexion with the problem of dealing with septic wounds—in fact, with most septic conditions. Various surgeons are using this method in treating septic fingers and infected tendon sheaths, cellulitis, osteomyelitis, varicose and trophic ulcerations, various types of dermatitis—in fact, all surgical cases in which non-traumatic cleansing, prevention of secondary infection, and early restoration of function are important. While the results from the closed-plaster method are excellent, the very fact that expert surgery and proper hospital accommodation are needed for success in this type of treatment prevents its use in most cases on active service, and it is here that the usefulness of the envelope system will be found.

About two hundred cases have now been treated by this method. Detailed below are five illustrative cases.

Case I

A man aged 64 was admitted on January 14, 1941, with phagedaena of the whole of the flexor surface of the left arm. He had grazed a knuckle two days before, and the arm had become painful and swollen. On examination the whole of the flexor surface of the arm had an appearance of one large sanguineous blister, and a vivid erythema was rapidly spreading over the shoulders and on to the face. The patient looked ill. Saline compresses and sulphapyridine were ordered, but without effect.

JOHN BUNYAN: TREATMENT OF BURNS BY THE ENVELOPE METHOD

FIG. 1

FIG. 2

FIG. 3

FIG. 1.—Tar burn, second degree. Slight areas of third degree.

FIG. 2.—After initial washing with 10% E.S.H. Oedema has started to subside; exudation has stopped. No pain. No anaesthetic.

FIG. 3.—After ten days.

FIG. 4.—Arm in full-length envelope: rubber band for use only during irrigation. Employed only as bath (instead of flow irrigation) where movement of injured fingers is required.

FIG. 4

JOHN BUNYAN: TREATMENT OF BURNS BY THE ENVELOPE METHOD

FIG. 5.—Severe third-degree burns of hands. Initial cleansing with 20% E.S.H.

FIG. 6.—After another month doing full household duties on farm. Absolutely full function.

FIG. 7.—Severe third-degree burn of child aged 4. Cleaned with 10% E.S.H. under analgesia. Note fibrin film.

FIG. 8.—Eight weeks later: almost healed; skin very strong and supple. Discharged two days afterwards, completely healed. Full movement of limb.

I was asked to see the case in the Radcliffe Infirmary, Oxford, by Mr. Abernethy, to whom I am indebted for permission to publish the record. The patient's arm was washed down for half an hour with 50% solution of E.S.H. and the arm fitted with an envelope. From the time of this irrigation the spread of infection was controlled, and within five days the whole of the skin and subcutaneous tissues had come away in one large slough, leaving the muscles and large blood vessels completely exposed. Daily irrigations of 10% E.S.H. were carried out, and in three weeks the whole of the area was covered with healthy bright red granulation. The patient was very well, was completely comfortable, and had almost free movement of his arm. A Thiersch graft was done on March 11 and the arm irrigated with 1% E.S.H. after twenty-four hours. This graft took perfectly. Further grafts were carried out. By March 28 all the skin-grafts had taken and the whole area was healed. The man had full function of his arm, and the new skin was quite supple. He was discharged healed with free function on April 7, and when seen on April 29 the skin was found to be very good, with only slight scarring, but complete function.

Case II

A man aged 25, admitted January 13, 1941, suffering from deep second-degree burns of back and side of hand and wrist, due to petrol on sleeve catching alight. Total area, 105 sq. cm. He was not shocked or in obvious pain, but there was some oedema and stiffness. He had had no previous treatment.

Morphine 1/4 grain was given, and in the theatre the hand was immersed in 12 pints of 10% E.S.H. at 100° F. and washed down for twenty minutes. All loose skin was cut away to periphery of burn. Patient experienced no pain, merely describing sensation as "tingling." A full-arm envelope was then applied. Next day the patient was completely comfortable, and slept well without a sedative. All oedema had disappeared, and a shiny film covered the wound. Thrice-daily irrigations were carried out with 8% E.S.H., and there was extensive epithelization, especially at the wrist. On January 21 a slight irritation was observed at the wrist, and oil was added after the irrigations. On the 26th the hand was cleansed and a new envelope fitted owing to the clouding effect of the oil. All sign of irritation was removed on giving ascorbic acid, 50 mg. thrice daily. By February 9 epithelization was complete, and the hand had perfect function.

At no time had the patient experienced any pain. There was no dermatitis under the seal, and the slight irritation in one part of the arm disappeared on administering 50 mg. of ascorbic acid t.d.s.

Case III

A woman aged 42 discovered in 1938 that she had diabetes. She went into hospital in London and was stabilized on a low starch diet, without insulin. Her urine was free of sugar for two days before she was discharged, but it had never been tested since. Diet worked until evacuation 6 weeks previously, when she began to eat starchy foods occasionally. Her condition became worse, so under protest she came into hospital. During the day her right leg had been badly burned by a hot-water bottle.

She was admitted on March 11, 1941, in semi-coma and was given 50 units insulin subcutaneously and 50 grammes glucose, also 2 pints of water by mouth. The following day she appeared fit and sat up in bed; felt better; given 15 grammes glucose intravenously. Blister burns from hot-water bottle on back of thigh burst, and were dressed with tannafax. On the 21st the burn was in very bad state, and sloughs were forming.

Operation, March 22.—Area of burn from flexor surface of knee extending upwards seven inches; two inches across top and five inches across at knee. The whole skin sloughed away under irrigation with 10 pints 10% E.S.H. at 100° F., and the area began to bleed. There was suppuration into the fat. Edges of wound were ragged. By the time the envelope was fitted haemorrhage had ceased.

Two swabs were taken of burn before irrigation, with the following results. 1 (a), Gram film: numerous pus cells, few Gram-positive cocci. (b) Culture: haemolytic streptococci, *Staph. aureus*, diphtheroids. 2 (a), Gram film: few pus cells, few Gram-positive cocci. (b), Culture: haemolytic streptococci, *Staph. aureus*, diphtheroids. The burned area filled with granulations in seven days, and was completely epithelized in three weeks. The patient was discharged healed, with good skin, on April 30. When seen on May 7 the skin was becoming pinker, and was quite supple and fine. She states that at no time did she experience any pain from the irrigations or any discomfort, either day or night.

Case IV

A man aged 26, admitted January 29, 1941, with petrol burn of palmar surface of wrist and hand, and two small areas on each side of dorsal surface of hand—second degree. Morphine 1/4 grain was given intravenously, and the hand was washed down for twenty minutes with 20% E.S.H., finishing with 10% for eight minutes. All blistered skin was excised. Patient did not complain of any pain, merely remarking that it was "tingling." After five minutes of washing the capillaries dilated and soon burst, but the slight haemorrhage persisted for only five minutes. The whole area became covered with a shiny fibrinous film. A hand envelope was fitted, and was inflated with oxygen.

The next day two washings with 5% E.S.H. were carried out, and the patient was discharged for daily treatment as an out-patient. The hand was irrigated with 5% E.S.H. each day, and epithelization was rapid. On February 10 the envelope was removed, and all areas were found to be covered with a pink cuticle. The hand was quite painless to touch, and full movement was possible. The patient was discharged, to return to work in a few days. He reported on February 20, having been employed, as a bricklayer for a week. The epithelium had cornified owing to the hard work, and local and general conditions were excellent.

This case shows that the method may be used successfully in the out-patient department.

Case V

On January 27, 1941, the patient, a woman aged 25, was cleaning the grate, and turned round to tie up one of her children's shoe-laces, when the back of her dress caught fire. She unsuccessfully tried to tear off the dress, thus burning her hands, and then plunged herself into a bath which happened to be full of water. The vicar's wife applied dressings of bicarbonate of soda, and the doctor was called in. At 1 p.m. she was brought into the Radcliffe Infirmary.

Examination showed extensive burns of the second degree from scapulae to sacrum, involving the right buttock; and of both hands, of second and third degree, extending over wrist and lower right forearm. At 1.10 p.m. morphine 1/4 grain was given, and at 2.50 p.m., in the theatre, the burns were irrigated with 10% E.S.H., all blisters and dead skin being removed. The left hand was immersed in 10% E.S.H., and some of the blisters were opened. The patient was encased in a coated silk envelope and the left hand in a hand envelope. She was returned to the ward and placed in a hot-box. At 7 p.m. the right hand was immersed in 10% E.S.H., and at 8.15 p.m. an envelope was applied.

The patient passed a comparatively good night, and next morning cleansing of the hands was attempted in the ward, but the burns proved to be too extensive. At 4.55 p.m. on the 28th blisters were removed in the theatre under gas-and-oxygen. Burns of the palmar surface were partially third degree. Hands were soaked in 10% E.S.H., and the envelopes reapplied. At 8 p.m. 1/3 grain of omnopon was given.

January 29.—Tolerable night. All three envelopes were irrigated with 5% E.S.H. and dried with oxygen.

January 30.—In theatre envelopes were removed. Back was washed with 5% E.S.H., and a new envelope enclosing the trunk and arms was applied. Movement of hands quite good.

January 31.—Envelope irrigated twice with 5% E.S.H. Sulphapyridine given.

February 1.—Patient not so well to-day. Complained of "stinging" on back. Arms irrigated 5% E.S.H. t.i.d. (An experiment was being carried out to determine whether the healing in the envelope method is due to the mere fact of isolating the part from secondary infection and from any irritation, or whether the E.S.H. is associated with the envelope in healing.) Patient's hands are being irrigated in the normal manner thrice daily, but the back is left untouched.

February 5.—Hands and arms irrigated with 5% E.S.H. t.i.d. Movement free.

February 7.—Hands and arms irrigated with 5% E.S.H. t.i.d. Back is covered with thick yellow exudate, which gives off a fetid odour. Where the exudate has run away from the burnt area healing has proceeded rapidly, but where the exudate has collected in the small of the back the granulations are unhealthy and healing is delayed. In the theatre the back was washed with 10% E.S.H. and all exudate removed. The whole was covered with a coated silk sheet.

February 8.—Back much cleaner, but still slightly painful when irrigated with 5%.

February 11.—Back is epithelizing well. Right hand has been removed from the envelope, and has perfect function.

February 13.—Bath given in coated silk. Right hand now quite well.

February 17.—Patient walked to bathroom and lay in E.S.H. bath for twenty-five minutes. Her back is almost healed, and feels "wonderfully comfortable." She is able to lie on the area, and moves quite freely.

February 18.—Patient walked to bathroom for her E.S.H. bath. Both hands are completely healed and have perfect function. There is no pain at all. The back has one small area on the lumbar region which has not healed over.

February 20.—Washings in bathroom are continued. Patient is happy and contented. There is still a small area on the back not healed.

February 28.—Patient discharged with hands perfectly healed, but one small area on the back not yet epithelized. She was given some E.S.H., with instructions how to use it on this small patch.

April 8.—Patient seen as out-patient. She uses her hands for heavy household duties, including the washing of clothes. The new skin on her hands is not shiny or tense, as in the case of treatment under tan, but is supple and allows free movement of all parts of the limb. There are no signs of scarring. Her back is quite healed, but the areas which were slowest in epithelizing are slightly raised. She can bend down and touch her toes quite easily, and is very well.

Clinical Progress

The preliminary irrigation rapidly cleans the whole area of the burn or wound. In many cases there is considerable capillary dilatation and transudation with spontaneous haemorrhage, which stops very quickly with the production of a shining fibrin coagulum; the patient feels little or no pain during the irrigation, and is almost always comfortable afterwards. Soreness may persist for a few hours in some cases of secondary burns. Oedema is often visibly reduced during the washing, and the patient may remark on the loosening-up of an affected limb. Excessive exudation tends to cease at the end of the washing. No signs of aggravation of primary shock have been seen. Tissue and bacterial toxæmia are favourably controlled, provided that the burn or wound has been adequately cleansed and all dead material removed. The latter is softened by irrigation, its adequate removal being thereby much eased.

So far as the daily irrigation through the envelope is concerned the patients remain comfortable and can lie on the burnt area with ease. They usually look forward to the irrigation, which they describe as soothing. In most cases temperature and pulse quickly settle and the burn or wound heals rapidly. Movement is encouraged, since it is certain that skin grown on a functioning joint is more likely to withstand use than skin grown on an immobile limb. Patients with burnt hands are made to use them as soon as possible from the time of the application of the envelope. They are encouraged to use a knife and fork and to grasp objects.

The morale of the patients is high and confidence is gained from the start; one never sees the frightened staring appearance so often associated with burns. Some apprehensive patients may describe the irrigation as giving a "tingling" or "pricking" sensation. This is never really unpleasant, and has usually disappeared after the first two irrigations. Secondary burns are more sensitive than primary ones. When epithelization is slow, short-wave therapy may accelerate it.

It has been found that the new skin grown over damaged areas as in third-degree burns is of sound and well-vascularized quality, though perhaps there have not yet been enough cases over a sufficient period of time to form final conclusions on this point.

It has been said before that the primary aim of this treatment is the control of infection. It has never yet been possible to sterilize a wound or the inside of an envelope, but from the clinical signs, and from the fact that the burns heal, it seems that the infection has been adequately controlled. There are some who would even go so far as to say

that a sterile wound is probably not desirable. Be that as it may, normal healing has been obtained in ten days by this method, and yet bacteria could be grown in the envelope. It must be said, however, that the numbers of bacteria are very small, and in some cases growth is difficult.

Results of Treatment Analysed

The following are the results of treatment of specific parts of the body.

Hand Burns.—In six cases treated by me relief of pain was immediate. The patients were encouraged to use their hands as much as possible, with the result that in all cases full function, with complete healing, was obtained within three weeks. Second-degree burns were healed, with full function, in ten days. In those cases in which it has been possible to follow up the results it is noteworthy that the new skin is quite free from scarring and is standing up to hard work. There is no sign of contraction.

Burns involving Joints.—Because full function is encouraged and obtained in every case, the new skin is grown to a fully flexed joint and is therefore not stretched at full flexion. Consequently there has been no breakdown of skin on a joint. In one case in which the joints were opened by the burn the hand was healed and is a satisfactory functioning member.

Face Burns.—In the few cases treated oedema of face and eyelids has been reduced completely within twenty-four hours and healing has progressed without interruption. There has been no sign of scarring or of involvement of the eyes or eyelids.

Body Burns.—Various areas of the body have been successfully treated in a vest, a "perineal bath," or a "full casualty bath."

Burns of the Perineum, Buttocks, and Upper Thigh.—The use of the perineal bath has given complete comfort to the patient and made the dressing very much easier than normally in this region. There has been no ascending infection of the urinary tract, and with the exception of one extremely severe burn of a quarter of the body surface, no infection from the urine or faeces. In the one case infection from pyocyanus occurred, but this was rapidly controlled.

Combined Burns and Fractures.—No case of combined burn and fracture has been treated, but from the experience gained in the treatment of compound fractures by this method it is quite certain that the problem can be solved by applying the necessary splinting outside or inside the envelope, whichever is desired.

Electrical Burns.—Uniformly good results have been obtained in the cases treated, resulting in full function within three weeks.

Chemical Burns.—One case of severe chemical burns has been treated under my direction at a military hospital.

A soldier was accidentally burned by mustard gas on both buttocks, involving the perineum, scrotum, penis, and anus. After ten days I was asked to treat the case. The patient had a thorough washing down with 20% E.S.H., the dead skin was removed, and a perineal bath was fitted. He was irrigated three times daily, and at the end of three weeks was discharged healed, and able to touch his toes without bending his knees.

Dermatitis.—There have been isolated cases of slight irritation of the skin. By changing the envelope and washing it before refitting, by lowering the concentration of the solution, and in one case by the administration of ascorbic acid, the trouble has always cleared up. It is rare to find the slightest irritation beneath the seal.

Chemotherapy.—It would appear that the sulphonamide type of bacteriostatic agent works at its best when dead organic matter is not present. An initial cleansing of the area to be treated with E.S.H. solution enhances the value of these agents. It may be that in a very septic case a

combination of the two types of treatment will be most successful. At present sulphanilamide is being tried out in the envelopes; hypochlorite is used to wash away exudate, and the envelope to prevent secondary infection.

Active Service.—The treating of burns and wounds with an envelope and sodium hypochlorite irrigation was devised as a simple method suitable for use under active service conditions—for example, in field ambulance, dressing station, or in the sick bay of a man-of-war. Its small bulk, its ease of application, and the fact that hypochlorites may be obtained anywhere should make it a most useful addition to modern surgery in warfare.

Conclusions

We feel justified in making claims with some certainty on several points, chief of which are: pain is relieved; primary infection is controlled, secondary infection prevented, and function restored at the earliest possible moment; primary shock is not aggravated, and secondary shock is minimized.

It is a system of treatment which will produce satisfactory results when used in small hospitals without specialist training, in ships, and in fact wherever the simple needs of the system are met. The catastrophes seen heretofore cannot occur when this method is used.

Future developments are in the orthopaedic field, in industrial accidents—in which immediate rehabilitation is so important—and in all branches of war surgery. Numerous varieties of treatment are possible, and many methods which were sound in conception but failed because of secondary infection may now be tried with an envelope, with the assurance that this problem will not arise.

Summary

Investigations into an improved form of dressing for burns and wounds are described.

The reasons for considering electrolytic sodium hypochlorite to be the best antiseptic to use are outlined.

The method of irrigating burns and wounds with this antiseptic, and dressing with coated silk envelopes which do not need daily changing and which shield raw areas from exposure to air and secondary infection, is described.

Burns and wounds covered with coated silk envelopes and irrigated with the appropriate concentration of sodium hypochlorite solution heal rapidly and painlessly, with the maximum preservation of function. The disadvantage of the painful daily changing of dressings is eliminated.

The envelope dressing can be combined with plastering or splinting in cases of fracture, and ulcerations of all kinds may be treated with advantage.

Five of the 200 cases treated by this method are described in detail.

Results are given of the treatment of burns on various parts of the body.

The advantages of this method of dressing for use during active service are emphasized.

I have to acknowledge my gratitude to Surgeon Rear-Admiral Cecil P. G. Wakeley, Mr. Vaughan Hudson, F.R.C.S., Mr. R. V. Hannay, F.R.C.S., Dr. Bruce Pearson, Prof. Hugh Cairns, F.R.C.S., the staff and sisters and nurses of the Radcliffe Infirmary, Oxford, and to all those who have so kindly helped and encouraged me in these investigations. Especially, I have to put on record my appreciation of the way in which Mr. Wm. Stannard has co-operated with me in the development of the envelope system.

M.M. Montgomery and J. D. Kirshbaum (*Arch. intern. Med.*, 1941, 67, 609) illustrate the rarity of anastomotic jejunal ulcer occurring after gastro-jejunostomy alone or combined with gastric resection by the fact that they could find only eight cases among 13,000 consecutive necropsies performed at the Cook County Hospital from 1929 to 1940. All the patients were white men and their average age was 49. Perforation and haemorrhage were frequent causes of death. The interval between operation for the original peptic ulcer and death varied from six weeks to twenty-one years, the average being eleven and four-tenths years.

WOUNDS AND BURNS TREATED BY COATED SILK FABRIC

A REPORT UPON AND CONCLUSIONS DRAWN FROM THE TREATMENT OF 82 CASES

BY

RUPERT VAUGHAN HUDSON, F.R.C.S.

Surgeon to the Middlesex Hospital and to an E.M.S. Hospital

On November 6, 1940, Bunyan and Stannard reported a case in which a large infected burn of the thorax and abdomen in a child was successfully treated in an occlusive envelope by immersion in and irrigations with electrolytic sodium hypochlorite. The envelope was manufactured by Stannard, and the electrolytic sodium hypochlorite was the proprietary preparation "milton" fluid, stabilized to 1% of sodium hypochlorite. The method seemed worthy of trial and capable of extension to wounds. Investigation was started in November, 1940, upon two main lines:

1. The use of coated silk fabric as (a) a fashioned dressing in the form of an envelope, and (b) an unfashioned dressing in the form of sheeting.
2. The clinical uses and properties of (a) electrolytic sodium hypochlorite, and (b) the sulphonamide group in conjunction with this special fabric.

In all, 82 cases were chosen because they showed ulceration in some form or another, or ulceration and infection. In a certain number the method was used because all other treatments had failed to relieve pain or to heal the wound, and one was at a loss to know how to proceed. The patients were treated at the Middlesex Hospital and the E.M.S. hospital to which I am attached. Although improvement in technique is continuing to be effected, it is felt that it is not too early to evaluate the results for the benefit of other general surgeons. In addition, there has been obtained a standard that enables us to contrast the quality of other substances. The following is a classification of the 60 cases whose treatment has been completed, 22 further cases being still under treatment:

1. Burns, primary and secondary.
2. Ulcers and gangrene.
3. Severe lacerating wounds, including compound fractures.
4. Acute pyogenic lesions.

Strict attention was paid to the treatment of shock, the use of prophylactic sulphanilamide compounds, surgical toilet, and constitutional and rehabilitation therapy when indicated.

Method of Treatment

In many cases the application of the envelope or sheeting, and in all cases the treatment, have been conducted by the nurses and dressers, who have also taught some of the patients to treat themselves. The envelopes were used as an occlusive or in occlusive dressing. The occlusive dressing was fixed to the intact skin of the patient by means of the seal incorporated in the envelope and reinforced by a light turn of the strapping; the in occlusive dressing by the removal of the seal, and the maintenance of the position by a light turn of the bandage, enabling the envelope to be removed at will for treatment or sterilization. The occlusive method provides the best means yet devised for minimizing the conduction and conveyance of infection, provided that the wound and the inside of the