and some infiltration at the left mid-zone. Cyst forms of *E. histolytica* were found in the stool. There was a leucocytosis of 10,200 per c.mm., with a 7% eosinophilia.

He was given a course of emetine and carbarsone, and his temperature came down to normal after forty-eight hours. A radiograph taken ten days later showed the lesion on the left side to be cleared, and there was some clearing of the lesion on the right. The course of emetine and carbarsone was repeated, and another radiograph after one month revealed still further clearing of the lesion on the right. At this stage the white cells numbered 8,700, with a 5% eosinophilia. Stool examination was negative for *E. histolytica*.

This patient is feeling much better; his cough and sputum have greatly decreased, and he is anxious to leave hospital. At the time of writing he is still under treatment, but will be discharged as soon as the lesion at the right base has cleared.

Commentary

The diagnosis of pulmonary amoebiasis is important because of its very rapid response to a specific therapeutic agent—emetine. In the diagnosis various factors must be considered:

(a) *History*.—A history of blood and mucus in the stools, no matter how long before the lung symptoms are manifest, is an important point. However, many patients do not admit to any previous bowel disturbances.

(b) *Site of Lesion*.—In secondary pulmonary amoebiasis the lesion is inevitably at the right base.

(c) *Stool*.—Examination of the stool very often reveals the presence of cyst forms of *E. histolytica*.

(d) *Sputum*.—The macroscopic appearance of the "anchovy-sauce" sputum is characteristic. Microscopically the sputum reveals an absence of other bacilli, such as tubercle bacilli. In primary amoebiasis *E. histolytica* is said to occur in the sputum, but in our series of cases, in both the primary and the secondary forms, we have been unable to demonstrate the entamoeba.

Differential Diagnosis

Tuberculosis.—Involvement of the right base with the apices clear is against tuberculosis, except in primitive races, in whom basal tuberculosis is not uncommon. Repeated negative sputum examinations would rule out pulmonary tuberculosis.

Bronchiectasis.—Bronchography would demonstrate the presence of bronchiectatic dilatation at the site of the lesion.

Foreign Body in Bronchus.—Unless there is a clear history of aspiration of a foreign body or the foreign body is easily seen on the radiograph, bronchoscopy is indicated in order to make a diagnosis.

Neoplasms.—A neoplasm may cause bronchial obstruction with resultant atelectasis, which may simulate secondary amoebiasis in the radiograph. Bronchography may show a characteristic "rat-tailing" of the affected bronchus. Bronchoscopy would be of value in differentiating a malignant from an innocent tumour by obtaining a biopsy specimen.

Syphilis of Lung.—As is well known, the symptoms and signs of syphilis are protean in their manifestations. Particularly in the Bantu, syphilis may show itself in the lung as a large tumour or as an extensive infiltration. The presence of a positive Wassermann reaction and the rapid response to the arsenicals would confirm the diagnosis.

Unresolved Pneumonia.—This may show as an opacity on the radiograph. Serial radiographs would show a gradual disappearance of the lesion.

Avitaminosis.—Jersild (1939) reported a case of haemoptysis and lung infiltration caused by avitaminosis. In the

Bantu in South Africa, more particularly in the urbanized native, symptoms of avitaminosis are very frequent. We feel, therefore, that a condition of avitaminosis should be considered in a differential diagnosis.

Where the diagnosis of pulmonary amoebiasis cannot be definitely established we feel that in suspicious cases the patient should be given the benefit of a therapeutic test with emetine and carbarsone.

Summary and Conclusions

Pulmonary amoebiasis is discussed.

Seven cases of primary and secondary amoebiasis are described.

The importance of examination of the stool, sputum, and blood is emphasized.

Bronchography may be of value in demonstrating a communicating hiatus between the primary lesion in the liver and the secondary lesion at the right base.

All cases of pulmonary amoebiasis show a striking response to emetine and carbarsone.

* We wish to thank Mr. M. E. Gibson, M.S.R., for taking the radiographs.

REFERENCES

- Jersild, T. (1939). *Lancet*, 1, 632.
Manson-Bahr, P. H. (1940). *Manson's Tropical Diseases*, 11th ed., London.

THE SURGICAL COMPLICATIONS OF AMOEBIC DYSENTERY

BY

H. W. S. WRIGHT, M.S., F.R.C.S.

Surgeon, E.M.S.

The British Empire is at the moment fighting for its existence in a vast area where amoebic dysentery is endemic. So ubiquitous is this disease that about one-fifth of the whole population is involved. After the evacuation of Gallipoli in the last war 23.7% of 971 cases were infected. In China the percentage of carriers in over 13,000 cases was 20.3, and amoebic dysentery is the direct cause of between 5 and 10% of all hospital admissions. Since its effects may extend over a period of from three months to thirty years it is important, at this time, to emphasize that it has significance in every branch of clinical medicine, and will continue to have for many years to come.

It is necessary first to recall some of the pathological features of the primary lesion, which is an ulcerative inflammation of the colon. The *Entamoeba histolytica* gains access to the large bowel through being able not only to attack and ingest the mucosa but to secrete a ferment which causes coagulation necrosis and tissue lysis. Having reached the submucosa, it multiplies in the centre of a small nutritive cytolysed area, which in its turn, although surrounded by some oedema, is not shut off by an extensive tissue reaction. Spreading in this way, it causes thrombosis of veins and venules, invades the clot, gets into the portal circulation, and is from then onwards caught in the wide mesh of the liver substance. That this is usually a most effective filter is shown by two facts: (1) it was found impossible to produce anything more than cloudy swelling of the liver by injecting very large numbers of amoebae into the portal vein of kittens; (2) amoebae rarely get into the systemic circulation, although amoebic abscess of the brain and the epididymis is well known to occur very occasionally.

It is true that cases of infection of the urinary and even of the uveal tract have been described; but these have not been accepted as genuine by many critical workers. Nevertheless the fact that such records do exist should be a

stimulus to further clinical observation. In any case, further involvement usually takes place by direct spread to contiguous tissues, from either the liver or the bowel, and its clinical results may be conveniently described under one of two headings—hepatic or colonic.

Involvement of the Liver and its Effects

As previously stated, hepatitis is secondary to an ulceration of the colon. This is widespread in about half the cases, but it is important to add that a large proportion of hepatic complications occur in just those cases in which the caecum of an apparently healthy patient is the seat of one or more solitary ulcers. When the liver is invaded by amoebae it may become suddenly enlarged and inflamed to a surprising degree. This phenomenon, known as acute hepatitis, is often accompanied by an acute inflammatory reaction in the contiguous pleura or peritoneum, and consequently may produce effects which equal in drama and interest anything that occurs in clinical medicine. Moreover, nothing is so therapeutically satisfactory as an acute condition which is resolved by the administration of 2 to 3 grains of emetine hydrochloride.

In the ordinary course of events, when the case is elucidated the following features emerge: (1) There is a history of diarrhoea or amoebic dysentery, which may coexist or may have occurred a long time ago. (2) There are fever and pain: the former is variable and is, in contradistinction to that caused by the "ordinary" acute abdomen, apt to be spiky, high, and associated with sweating and chills; the latter is classically referred to the shoulder or, if the parietal peritoneum is involved, to the abdominal wall, causing rigidity and tenderness. (3) The liver is enlarged. (4) There is a characteristic leucocytosis of from 10,000 to 30,000, in which the polymorphonuclear cells seldom make up more than 75 to 80% of the total. If the leucocytosis is more than 80% the condition is probably not amoebic or there is an associated secondary infection.

Differential Diagnosis

These cases, often admitted as emergencies, are diagnosed by the clinician who is not "amoeba-conscious" as diaphragmatic pleurisy due to basal pneumonia, or appendicitis, or the perforation of a hollow viscus. Malaria with pneumonia, and acute cholecystitis, are not uncommon preliminary diagnoses, made before or on admission. It is fortunate in acute cholecystitis that the necessity for intervention is not urgent and that emetine does it no harm, for the differentiation may produce real difficulties. But the word "diarrhoea" somewhere in the history gives a clue to the diagnosis; the blood count provides strong circumstantial evidence; and the examination of a little blood-stained mucus from the stool makes it a certainty. Emetine provides a cure for the immediate symptoms. Below are two typical case histories:

Case 1.—A woman aged 50, when seen, was doubled up with acute abdominal pain. There was extreme tenderness and boardlike rigidity over the upper right side of the abdomen, which extended beyond the midline to the left, and on the right, down as far as the right iliac fossa. She had vomited at the onset about five hours previously; temperature was 103°, pulse 110. There were no physical signs of pleurisy or pneumonia, and no history of dysentery except that everyone in the East has a short attack of diarrhoea at some time or other. There was a vague history of indigestion, more suggestive of cholecystitis than anything else. No special diagnostic methods were available, so with some trepidation she was given an anaesthetic, and on palpation of a relaxed abdomen the liver was felt to be enlarged.

Case 2.—This patient, aged 43, was ordinarily a robust healthy woman. She was three months pregnant, and during an in-

fluenza epidemic complained of some malaise and a little pain in the back, with a temperature of 100°. Two years previously she had suffered a mild attack of acute amoebic dysentery, which was adequately treated and carefully controlled in the interval. Her stools after treatment were consistently negative, both for amoebae and for cysts. On examination her liver was painlessly enlarged down to just below the anterior superior spine. She had a leucocytosis of 23,000, of which 72% were polymorphs. This condition rapidly subsided with emetine therapy, and for the last fifteen years no further symptoms have appeared.

Between these two extremes come the cases which unfold themselves with textbook uniformity. In all of them, once the diagnosis is made, the treatment is simple. It consists in giving 1/2 grain of emetine deep subcutaneously or intramuscularly thrice daily for about three days, after which the dose is reduced to 1 grain daily. More than 15 grains should not be administered at any one time, and the patient should be in bed. After a few days' rest the standard routine treatment for amoebic dysentery, consisting of emetine bismuth iodide and quinoxyl, should be given.

If this acute stage is unnoticed the case may progress until massive colliquative necrosis of a large or small area in the liver has taken place and the patient gets a liver abscess. Its physical signs and symptoms are usually clear-cut. They are well known and adequately described. But in spite of this it has to be realized that almost every experienced physician in tropical practice has at some time or other seen at necropsy a large liver abscess which has been unrecognized during life.

Within the last few weeks a patient was admitted to the French Hospital in a state so serious as to preclude exact diagnosis or treatment. A few weeks previously a perinephritic abscess had been opened elsewhere. He died within twenty-four hours, and at necropsy his liver, which was of normal dimensions, was found to contain an abscess the size of a large orange in the right lobe posteriorly. Pleural and peritoneal reaction was minimal.

For the exact diagnosis and treatment of liver abscess paracentesis is necessary, and when proper precautions are taken this is a safe procedure. These precautions are: (1) the needle track through the parietes should be anaesthetized with novocain; (2) the needle should be of fairly wide bore and be marked at intervals of half an inch; (3) it should not be inserted more than 3½ in. for fear of entering the vena cava, and its direction should be upwards and inwards; (4) the shortest route to the abscess should be taken, and this is indicated by local bulging or radiological evidence of a change in the height of the diaphragm. Aspiration should be repeated until no more pus is obtained, and the pus should be cultured. Amoebae are seldom found in the first aspiration.

One important fact should always be borne in mind. A liver abscess is a cold abscess and should be treated as such. If culture shows it to be secondarily infected drainage is indicated, and in such cases it is imperative that the drainage track should be sealed in some way or other. The growth of a few small colonies of staphylococci, however, is not in itself an indication for radical surgical measures. Where emetine has no effect on the temperature chart it is a sure sign of secondary infection. If the patient has a pleural effusion and a liver abscess beneath the diaphragm the following line of treatment should be followed:

Both pleurisy and the liver abscess should be treated by aspiration. If the former becomes grossly infected from the lung, pleural drainage should be deferred, if possible, until the liver abscess has been aspirated several times and the patient is well under emetine treatment. If both are secondarily infected, then radical transpleural and/or transdiaphragmatic drainage should be done. In the few cases of this kind seen by me the pleural cavity has been well sealed off by adhesions.

Further extension of the disease beyond the liver is as a rule by direct spread. It is well known that the diaphragm and lung may be involved and that a pleural effusion may occur as with any inflammatory process below the diaphragm, but it is not so well known that from the liver it may spread to the skin, producing a large ulcerating surface. Such a case I was privileged to see in consultation. It was reported and illustrated by Heimbürger (1925) so precisely that the account may profitably be consulted by anyone interested in the histopathology of amoebiasis. There was an area about the size of the palm of one's hand covered with pale oedematous granulations; its appearance was as if a quantity of red brickdust had been ground into the ulcerated surface. It healed rapidly with emetine therapy.

A liver abscess sometimes extends to the abdominal cavity. When this happens the size of the liver may be difficult to determine, and an inflammatory swelling arises which is, as a rule, somewhere continuous with the liver margin. This swelling may extend quite low down in the abdomen, limited either on the right or on the left by the falciform ligament. When it occurs on the left side it is very puzzling, because the occasional abscess arising in the left side of the liver is frequently overlooked. Such an abscess may extend well to the left above the spleen. In ordinary clinical practice it would be regarded as the result of a localized perforation of a peptic ulcer. Such localized intraperitoneal collections of pus from a liver abscess often have a somewhat indirect connexion with the liver, and it may be wise to open them and wash them out. If this course is taken, intentionally or inadvertently, the wound should always be sewn up without drainage and the parent liver abscess aspirated after localization. No harm comes if the patient is immediately put under emetine, and this should, if possible, be invariably done before any intervention is contemplated.

An extremely acute abscess may sometimes perforate into the peritoneal cavity, the case being admitted as an acute abdominal catastrophe. The condition is nearly always such as to preclude any surgical intervention.

Direct Spread from Colonic Lesions

Involvement of the muscular and peritoneal coats of the bowel wall is not uncommon. Sir Leonard Rogers has found amoebae in the lymph adherent to the base of an unperforated ulcer. Actual perforation is rare, but a localized slowly spreading peritonitis is by no means uncommon. It is important to bear in mind its possible existence, because it complicates the diagnosis of other abdominal conditions. The finding of amoebae in the stools makes the diagnosis simple.

Localized abscesses round the appendix are well known but not often seen, and the same may be said of retroperitoneal and perinephric abscesses. If their amoebic origin is not recognized extensive invasion of the abdominal wall may take place.

Ulcers of the colon may be secondarily infected and may produce a localized inflammatory mass which resembles diverticulitis. The lumen of the bowel may be so narrowed by oedema or fibrosis as to produce obstruction. In these cases routine treatment of the dysentery is not always sufficient, and a resection has to be performed.

Finally, the rectum is a common situation for amoebic lesions, and in severe untreated cases these involve the tissues in the ischio-rectal fossa, producing fistulae and ischio-rectal abscess. These may occur in patients who are extremely ill, originating in quite small ulcers without much surrounding infiltration. In more chronic cases the local ulceration may be widespread and, owing to second-

ary infection, the surrounding tissues be infiltrated widely. This may produce a particularly intractable form of stricture requiring colostomy for temporary relief.

Conclusion

It should be remembered that local lesions in the bowel do not as a rule progress thus far, but they do frequently and for a long time act as foci of irritation, exercising a profound influence on the physiology of bowel movement. Such patients have localized spasm and tenderness for years, their bowel is irritable, and they pass frequent stools containing a little mucus. They become bowel-conscious, and haunt the "colon lavatories" in the West End of London and the consulting-rooms of sympathetic physicians. They are diagnosed as cases of anything from spastic constipation to mucous colitis. They require above all things rest, both of the mind and of the colon.

This account by no means exhausts the widespread manifestations of amoebiasis, but enough has been said to show that a knowledge of its manifestations is necessary to every clinician handling cases in or from the East, and to suggest that to the use of the stethoscope must be added that of the microscope and sigmoidoscope as part of the routine examination of patients. By the employment of these methods we arrive at a positive diagnosis of a disease which can be "cured" in a sense that cannot often be used in the case of diseases commonly met with in this country.

REFERENCE

Heimbürger, L. F. (1925). *Arch. Derm. Syph.*, Chicago, 11, 49.

CYSTICERCOSIS EPILEPSY

BY

CECIL W. EWING, L.R.C.P.&S.I., D.P.M.

Late Medical Superintendent, West Riding Mental Hospital, Storthes Hall

The case reported by Dr. W. Blyth (1941), to which he applied the above diagnosis, has many points of similarity to one which came under my notice some years ago, and in which a diffuse cysticercosis of the brain was found post mortem. Out of a large number of post-mortem examinations on mental hospital patients extending over a period of thirty-odd years this was the only case I met with. The condition, though rare, is of interest because—as pointed out by Sir W. P. MacArthur (1941)—of "the importance of considering it as a possible cause of epileptiform attacks commencing in a previously healthy adult after service or residence abroad." The histories of all the cases I found recorded come within this category, though one might extend the term "epileptiform attacks" to those presenting other ill-defined and vague symptoms of cerebral disease. The value of radiographic investigation in the diagnosis is emphasized by Dr. Blyth, but it is not clear from the particulars he gave that these investigations did reveal the presence of cysticerci in the brain tissue of his patient.

References in Literature

The case which came under my observation, and in which unfortunately the possibility of cestode infection as a cause did not occur to me, demonstrated from the necropsy that a condition of cysticercosis of the brain diffuse in distribution produces a clinical picture of epilepsy with dementia difficult to distinguish from the idiopathic form except by the more rapid development of the dementia and of a fatal termination. The cysts examined showed no evidence of calcification. I was so impressed by this massed invasion of the brain by these parasitic cysts that I consulted a number of standard textbooks on psychiatry and cerebral