50,000 units of A.T.S. was given. A dose of paraldehyde was necessary at midday and avertin at night.

On the fifth day after admission to hospital spasms were less severe and more infrequent. Food was being well taken until evening, when spasms became frequent and intense. The patient complained of pain at the right angle of the jaw. A dose of avertin at night brought two hours' sleep, so 4 drachms of paraldehyde was given at 3 a.m. The condition of the patient was worse and spasms were still causing distress, consequently 20,000 units of A.T.S. was injected round the site of the injury and intramuscularly, and 60,000 units intravenously. A further dose of avertin caused some relaxation. Only slight improvement was noticeable the following day, and a dose of avertin was necessary to prevent restlessness at night.

There was improvement in the general condition the next day and avertin was not necessary as a sedative; muscular spasms were slight and caused no pain. The patient now made an uninterrupted recovery and was discharged on October 24.

During treatment nursing attention was paid to pressure points and an airway was placed between the teeth to prevent the tongue being bitten. Between spasms the patient could take food naturally. Egg and milk, glucose orangeade, egg custard and cream, with nourishing beverages, were given *ad lib.* One drachm of potassium bromide was given daily throughout.

My thanks are due to Dr. J. W. Wheeler for permission to publish this case.

F. R. BREBNER-SMITH, M.B., Ch.B.

A Case of Double Tubal Gestation

The following case is rare enough to merit being placed on record:

CASE REPORT

The patient, a 4-para aged 30, was admitted to Maryfield Hospital on March 18, 1940. All her previous labours had been without complication. On admission she had vaginal bleeding which began on February 9. Her two previous periods had been normally spaced, but for the five or six months before then they had been coming on every three weeks. She complained of pelvic pain between the periods, the pain being accentuated with the onset of the flow. On March 20 she was examined under a general anaesthetic, and swellings were felt in the lateral fornices. The uterine cavity measured $3\frac{1}{2}$ inches. Curettage was done, and the scrapings were sent for pathological examination. The report received was as follows: "The rather atrophic endometrium shows no evidence of decidual change. Small polypoid growths are present on the surface, and in the interstitial tissue plasma cells are present in fair numbers, indicating a degree of endometritis."

On March 29 the abdomen was opened: some free blood was found in the peritoneal cavity; both Fallopian tubes were distended and markedly pathological, and both were removed. We received the following pathological report: "This is a case of double tubal pregnancy. The larger Fallopian tube contains a gestation sac within which is a foetus of about the seventh or eighth week. The smaller tube is filled with a mass of blood clot in which chorionic villi are present. The embryo was not found. The appearances of this tube are those of a tubal mole." The right tube was the larger one and contained the foetus.

Various textbook writers mention the occurrence of intra-uterine and extra-uterine pregnancy at the same time, of multiple pregnancy in one tube, and of coincident pregnancy in both tubes. Also there may be a consecutive tubal pregnancy in the same patient. Eden and Lockyer state in their *Textbook of Gynaecology* that simultaneous binovular tubal pregnancy is the rarest form of all. They say that up to 1927 at least 42 authentic cases of ovarian pregnancy had been collected, while they had found only 28 cases of proved simultaneous pregnancy in both Fallopian tubes. They also refer to a particularly interesting case seen by Treub of Amsterdam who found quintuplets in one tubal gestation sac.

A. E. CHISHOLM, F.R.C.S.Ed., F.R.C.O.G.,
Obstetrician and Gynaecologist.

ABIGAIL LESSLIE, M.B., Ch.B., D.P.H.,
Late House-surgeon, Obstetrical and
Gynaecological Department.

Maryfield Hospital, Dundee.

Reviews

BACTERIOLOGY

General Bacteriology. By D. B. Swingle, Professor of Bacteriology and Dean of the Division of Science, Montana State College. (Pp. 313. 16s. net.) London: Chapman and Hall. 1941.

Of recent years a succession of textbooks has been published in the United States which deal with bacteriology as an independent science, and with its medical as only one of several important applications. Such a work is *General Bacteriology* by Prof. D. B. Swingle, who deplores, as other authors have, the manner in which the subject is usually taught to medical students. The average course in medical bacteriology is in fact confined to the study of pathogenic bacteria only, to those characters by which they can most readily be recognized, and to such aspects of their behaviour as bear on the phenomena of infection. It would of course be far better to approach the subject in the first instance as a pure science, to present a general instead of a particularized view of what bacteria are and what they do, before dealing with the special characters of pathogens. It need scarcely be said that what stands in the way of such an approach is not so much a wrong attitude on the part of teachers as the lack of time in an already overcrowded curriculum. Another factor is the disinclination of students to revert to long hours of laboratory work having no direct application to medicine, when clinical work has captured their interest, and qualification and practice are their only goal. If the subject is to be taught in this way it should be done in the first year, as a part of instruction in biology; the teacher of medical bacteriology in the clinical years would then find a foundation on which to build. If such a reform ever comes about such a book as this would suit the first-year student. It deals successively with classification, morphology, and bacterial metabolism in a general way, and then gives some indication of the functions or other importance of bacteria in various environments—air, soil, water, sewage, foods, and industrial processes. The few final chapters on pathogenic bacteria and the phenomena of infection are too rudimentary for the student of medicine and in places actually misleading, as in the brief description of a tuberculous lesion, which is said to be an example of "hypertrophy."

The book is profusely illustrated, particularly with line drawings, some of which are over-simplified. Each chapter concludes with a series of "review questions," often of disarming simplicity and rather obvious or redundant; the prize for these qualities goes to "What is meant by 'bacterial flora'?" and "In what ways are human excreta more objectionable than soil?"

MEDICAL ASPECTS OF BOXING

The Medical Aspects of Boxing. By Ernst Jokl, M.D. (Pp. 251; illustrated. No price given.) Pretoria: J. L. Van Schaik, Ltd. 1941.

"Pushing the thumb into the lateral region of the neck is a well-known rough-house trick with which the average clinician is naturally not too well acquainted." So states Dr. Jokl, the author of this fascinating book, which contains a very great deal with which the average clinician is not acquainted. This single quotation may give some idea of why the textbook may be claimed to be unique of its kind. Dr. Jokl has already contributed much to our knowledge of sports injuries, and between 1930 and 1933, in collaboration with Dr. E. Guttman, carried out what was probably the first comprehensive investigation of the neuropsychological aspects peculiar to the injuries