CHEST SURGERY IN WAR

This is the last of a short series of articles based on lectures given at the British Postgraduate Medical School, Hammersmith

DRAINAGE OF THE PLEURA

BY

R. C. BROCK, M.S., F.R.C.S.

Although pleural drainage does not afford a subject so dramatic as certain other features of war surgery of the chest, such as the removal of foreign bodies, it is perhaps more important than any other. It profoundly influences both mortality and morbidity, and is probably the greatest single factor in determining the length of invalidism, which in a mismanaged or unfortunate case may be lifelong. Like the treatment of pleural infection in civil work it can be conveniently considered under three headings: (1) when to drain the chest; (2) the provision of adequate drainage; (3) when to stop drainage.

When to Drain the Chest

Apart from the presence of frank suppuration the question arises as to drainage in less precise circumstances. It is well to remember two guiding principles which are of value in deciding whether or not to drain the peritoneum and which may well be applied to the pleura: (a) Is a condition present which the pleura cannot deal with by itself? (b) Is it anticipated that a condition will develop which the pleura will not be able to deal with by itself? We have a shrewd idea of what the peritoneum can deal with by itself, and I am coming to the conclusion that the pleura can deal almost as effectively with a certain amount of contamination, provided (a) the lung is healthy and no fistula is present; (b) no infected foreign body is present; (c) there is no infected chest-wall wound.

A penetrating chest wound (other than a small one without haemothorax or pneumothorax of any size) should be excised and either sutured, if only a few hours old, or packed open. The question arises as to whether the pleura should be drained in these cases. If the wound is large and with much laceration and contusion, or if there is a retained foreign body or damage to the lung, then the pleura should be drained by means of an air-tight intercostal catheter connected to an underwater seal and left in place for forty-eight or seventy-two hours. In fact, the surest general rule to follow is to drain every time. An intercostal catheter in use for two to three days can do no harm, and although perhaps unnecessary in some cases, will be responsible for averting disaster in many others. I have seen undrained cases of this type heal satisfactorily even when a haemothorax was present and had not been evacuated; the deeper layers have healed over and the haemothorax has been successfully treated by aspiration, thus proving that the pleura has been able to master some contamination at any rate. On the other hand, I have seen an even larger number of cases in which infection has supervened, drainage has been needed, and a long period of disability has resulted. There is no doubt that it is most unsafe not to drain if the pleura has not been sucked clear of blood and serous exudate before the chest-wall wound is closed; any such residuum is a dangerous culture medium to leave in the pleura.

Although routine drainage for two to three days is recommended, it is permissible for a surgeon familiar with the chest not to drain in selected cases, provided he is able to watch the patient closely himself for at least seven to ten days.

DANGERS OF SUTURING TOO SOON

When the first war casualties began to arrive from France we found that many of the chest wounds had been excised (often very incompletely), sewn up tight without drainage even when the lung had been incised to remove a large foreign body, and then transported within a few days. Much of our work consisted in taking out stitches, packing open the wound, which was the centre of a severe, spreading, sometimes gangrenous, cellulitis, and inserting an intercostal catheter into the infected pleura. Severe or fatal anaerobic infection may occur, and at the very least the patient has to face weeks and sometimes months of slow convalescence from a total pleural suppuration, with the lung perhaps completely collapsed and often containing one or more foreign bodies. The catastrophe could have been easily avoided in almost every case if the chest-wall wound had not been sewn up tight, but had been left open, and if a temporary intercostal drain had been used.

There is no doubt that these disasters have occurred owing to misunderstanding of the teaching about the treatment of an open sucking wound. It is generally taught, and generally accepted, that these wounds must be closed at all costs. That is quite correct; they must be closed, but not necessarily sutured. It is not safe to excise and sew up tight a big wound of the chest wall with lacerated and contused muscles; in the thigh or arm such a wound would not be sutured but would be packed open initially until the risk of spreading infection had passed. It is not possible to ignore a basic surgical principle such as this. The muscles and fascial planes overlying the thorax, far from being immune, are very susceptible to infection, which may rapidly give rise to a severe phlegmon. Unless, therefore, the wound is relatively simple, is completely excised soon after being received, and can be watched personally for at least a week, it must not be sewn up. A flat pad should be laid over it and secured by properly applied firm strapping. Such a wound is closed mechanically, but surgically it is still open. The pleura should be drained, as stated above, for two to three days. If it is then found that the lung is adherent and the deeper layers of the wound are shut off from the pleura the drainage tube may be removed, although careful watch will have to be kept for reaccumulation of effusion, which will need drainage again if aspiration reveals that it has become infected. The parietal wound can, and in fact should, be closed by delayed suture after a few days if the risk of spreading infection seems to have passed.

The management of a haemothorax may present many difficult problems, but one of the greatest dangers to be on the alert for is the onset of infection. This may not occur until as late as ten to twenty days after receipt of the wound, and may lead to death from anaerobic infection in a patient who had appeared to be almost convalescent. Presumably infection occurs from sloughing of devitalized lung tissue in the track of the missile. Close watch must be kept for the onset of infection; the temperature may be no true guide, but a rise in the pulse rate is significant. The fluid in the chest must be sampled at least every other day, and should be examined at once for evidence of infection. It is not enough to send a specimen to the laboratory. The haemothorax itself is an excellent culture medium, and much information can be obtained by its direct study; when infected the blood may

smell, haemolysis occurs, the colour changes to a purplish hue, and if a direct smear is examined micro-organisms will be seen. Culture in the laboratory will only confirm these findings. It is most important that samples of fluid aspirated from the chest should be kept by the patient's bed and properly labelled so that they may be available for inspection at any time. It is a common but nevertheless stupid habit to throw valuable evidence of this sort away and not to preserve it for future reference.

Once infection of the haemothorax has occurred drainage is needed. The type of drainage is considered in the next section.

Provision of Adequate Drainage

In a certain small number of cases pleural infection can be satisfactorily overcome by repeated aspiration of the exudate, but though this minor procedure is occasionally successful it is wrong to put it forward as a safe or standard method of treatment, except to tide over a few days, or two to three weeks at the longest. The same applies to intercostal drainage, which is invaluable as a temporary measure—in fact, is often life-saving—but will allow rapid and safe healing of pleural suppuration in only a few cases. It is a great mistake to imagine that all will go well and that rib resection can be avoided just because the patient looks and feels better and the temperature has subsided. The fundamental principle of civilian surgery still holds good, that in almost every case of empyema the adequate drainage provided by rib resection is essential to allow perfect healing in the shortest time and with the minimum of disability. To continue with a small intercostal drain for more than a few days is to expose the patient to the grave risk of a chronic empyema. The step of rib resection is a small one, should take only a few minutes, can be performed under local or a short gas anaesthesia, and should be no burden even to a very ill patient.

At the same time it is important not to drain the pleura too early. In war wounds, as in the synpneumonic pleural infections of civil life, there is usually diffuse infection of the whole pleura with no localization. If drainage is performed too early a total empyema results, the lung often remaining collapsed down on the hilum; in others the lower lobe expands more rapidly than the upper lobe, the drainage point is soon shut off, and a large residual or imperfectly drained anterior pocket is left, usually extending to the apex. These cases have been only too common and have proved to be very difficult to treat; they tend to become chronic, take months to heal, and leave the patient with an impaired respiratory function on the affected side. In many it has been almost unavoidable when aspiration has been difficult and unsatisfactory owing to the debris of infected clot, and the patient has been so desperately ill that early intercostal drainage has been needed as a life-saving measure. But unless this is the case, tube drainage should be postponed and aspiration persisted in until the pleural infection has become localized to give a true abscess. The aspiration must be performed regularly and efficiently and not infrequently and perfunctorily; by keeping the pleura as dry as possible not only is toxæmia diminished but the ultimate empyema will be a small one and therefore quicker to heal when it can be adequately drained by rib resection.

For the usual posterior basal empyema the best drainage is given by resecting about 3 cm. of the ninth rib just outside its angle. If a lower rib is taken not only is there risk of wounding the diaphragm but in the natural process of healing the ascent of the diaphragm will shut off the drainage hole and may cause a chronic empyema. Simi-

larly drainage too far forward is to be avoided; a tube inserted in the mid-axilla or anterior axilla cannot drain the posterior pleural recess efficiently, and a chronic empyema is almost certain to follow. Unless the patient has a pleuro-bronchial fistula there is no reason to fear turning him on his good side to resect a rib in the proper place at the back; to try to work with him lying on his back is inefficient and footling. If a broncho-pleural fistula is present the operation can be performed with the patient sitting up and under local anaesthesia; it is probable, however, that if the fistula is of any size a preliminary intercostal catheter will have been inserted earlier.

It is best to connect the drainage tube to a simple empyema bottle; this enables the pus to drain away cleanly, saving much labour, disturbance, and discomfort from constant changing of pus-soaked dressings, and it encourages expansion of the lung by maintenance of a mild negative pressure.

When to Stop Drainage

The termination of pleural drainage is a most important phase, and may be difficult to assess exactly. When drainage has been used as a safeguard in the presence of a certain amount of pleural contamination it can be stopped in two to three days; but when it has been needed because of an established infection the problem is quite different. There is no doubt at all that mismanagement of the drainage tube or its too early removal is responsible for much unnecessary suffering and a great deal of permanent crippling. Unfortunately the correct management of this phase has been understood during only the last decade or so, and the knowledge is still far too scantily appreciated. This is at once seen if standard surgical textbooks are consulted; it will be found that no correct rule is given as to when the drainage tube should be removed, but only some quite incorrect rule-of-thumb advice such as "after three weeks" or "when the discharge stops" or "when the discharge becomes thin." There is only one safe time to remove the tube, and that is when it is no longer possible to demonstrate a cavity in the pleura. This is a rule that cannot be too widely appreciated or stressed too much. Failure to observe it is responsible for more chronic empyemas than any other single cause.

In the healing of a simple acute abscess, provided drainage is maintained adequately, the walls collapse together and adhere by granulation until the cavity is completely obliterated; the discharge then ceases, the skin grows over, and healing is permanent. The proper healing of an acute empyema is no different: provided adequate drainage is maintained the walls come together (that is, the lung expands; collapse of the parietes is of course avoided by proper exercises) and adhere by granulation—that is, there is progressive symphysis of the visceral and parietal pleurae until the whole pleural space is closed. Then and then only should the drainage tube be removed. The final track from pleura to skin will close simply and certainly, and permanent cure follows. Should the tube be taken out when a pleural pocket still remains, progressive and satisfactory healing may follow, but it is much more likely that a chronic empyema will occur. Even if the skin heals over temporarily a latent pus pocket remains to be a constant source of toxæmia, and sooner or later (perhaps weeks or months, perhaps many years after) will flare up and lead to "recurrence."

The best way to decide the problem of tube removal is to investigate the size and extent of the cavity by a pleuro-gram—that is, radiography after instillation of a radio-opaque fluid or oil such as neohydriol or lipiodol. This should be done when it is felt that the cavity has

diminished to a reasonably small size; from a study of the size, shape, and disposition of the cavity any necessary adjustment in the size and disposition of the drainage tube can be made. Thus it is usually necessary to shorten the tube in the later stages of healing, but it may be shown that not shortening but lengthening is needed if a high-up or anteriorly placed loculus is not to be shut off and become chronic.

Measures Promoting Lung Expansion

It is impossible to speak of pleural drainage without referring to the measures to be used to promote lung expansion and, incidentally, to prevent flattening or deformity of the chest or scoliosis. In stubborn or difficult cases lung expansion may be aided by constant negative suction (e.g., 2 to 8 cm. Hg negative pressure) applied by means of a Roberts bottle or, when the patient is mobile, by Brock's portable suction apparatus. This suction can never make up for inadequate drainage; it must never be used to suck pus out through a small hole when it should really run out through a large hole, nor should it be allowed to replace proper breathing exercises, to which it is merely complementary. These should be taught by a trained masseuse, and the patient should be impressed with the need for daily hard work in their observance; it is not enough for him to spend a gentle ten to twenty minutes with his instructor each morning, but he should try to devote to them a total of several hours a day by himself. The best type of exercises are those of the inspiratory type described by McMahon, which aim at restoration of proper and symmetrical costal breathing and controlled diaphragmatic movements. They are supplemented by appropriate remedial exercises to maintain proper posture. The use of the homely "blow-bottle" is not to be recommended, although it has a place in certain selected cases.

The treatment of an empyema should, moreover, be ambulatory as soon as the patient's general condition allows him to get up. Proper expansion of the lung and obliteration of the infected pleural dead space is thereby hastened, always provided adequate and correctly placed drainage is maintained.

As a contrast, how often do we see those unfortunates who have been ill for months, perhaps years, still in bed most of the time, with a horribly flattened, deformed, and seriously impaired chest and a continuously running small sinus that either has a minute tube or no tube at all, or, worse still, has a strip of ribbon gauze tucked in it. Let us try to save our wounded fighting men and civilians from this fate—so tragic because it is avoidable, but one which has befallen many in the past.

Under proper and thoughtful guidance the healing of an infected pleural space can be achieved in a short time and with no permanent disability at all; inattention to the simple principles of drainage laid down here may result in the production of a chronic empyema which will entail perhaps a lifetime of suffering and disablement to the patient, and great economic loss not only to himself and his family but also to the nation, which may be called upon to grant him a permanent pension. In addition the services of a useful citizen or soldier may be unnecessarily lost. Not the least of the tragedy is that in most cases it could have been completely and easily avoided by taking proper care at one phase of management of the healing stage; in most this phase is that of regulation of the drainage tube, and occurs through failure to appreciate the seriousness of its casual premature removal.

Remember that the treatment of pleural suppuration does not consist in doing an operation on the patient. The operation for providing adequate drainage is only an

incident in treatment, albeit an important one. Treatment is concerned with securing and maintaining full lung expansion and thereby complete healing of the infected pleura. So often the surgeon's interest (and indeed the interest of all the patient's attendants) becomes progressively less as the days pass into weeks after operation. The reverse should be the case: interest and attention should increase as the stage of healing advances, and in the final stages of obliteration the most scrupulous care is needed to manipulate the tube and arrange its removal at the correct time. Only when this is fully appreciated will the number of crippling and chronic empyemas caused by this war be diminished.

CLEARANCE OF URBAN AIR-RAID CASUALTIES AT THE PERIMETER

BY

A. MASSEY, C.B.E., M.D., D.P.H.

Medical Officer of Health, City of Coventry

G. G. ALDERSON, B.A., M.B., F.R.C.S.

Group Officer, Coventry Area

AND

C. F. BROCKINGTON, M.A., M.D., D.P.H.

Deputy Medical Officer of Health, County of Warwick

One of the major lessons taught by experience of severe enemy air attack is the danger resulting from a central location of general hospitals in vulnerable towns. Fortunately indeed are the towns possessing general hospitals at the periphery. But in the majority of cases the hospitals are on central sites, and thus it is that heavy damage to their personnel and fabric is now an unhappy commonplace. These things have obvious bearings upon casualty clearing schemes in the localities. The hospitals in a town are ordinarily the natural centres for the initial admission and classification of the more severe air-raid casualties pending outward transfer to affiliated base hospitals. The sequence of events then is: (1) collection of street casualties—inward flow of laden ambulances to central hospitals; (2) "sorting" and initial treatment of casualties at central hospitals—outward flow of other laden ambulances to base hospitals. The minor cases are, of course, treated at the first-aid posts.

When the Inner Ring is Heavily Bombed

This works well enough in raids with circumscribed incidents. But when the entire inner ring of a town is subjected to heavy and continuous bombardment there are obvious and serious difficulties owing to: (a) actual or potential damage to the central hospitals themselves; and (b) the danger entailed in bringing loaded ambulances through the zone of maximum bombing. These conditions are more liable to arise in the county borough of ordinary size, for in the larger cities the inner ring is of such extent that it cannot be completely "plastered" in quite the same way.

In the average county borough of known or supposed vulnerability, therefore, whose hospitals are located centrally, it is well to have in readiness a scheme for the clearance of casualties at the perimeter. This means that, on a given signal to the ambulance depots, all ambulances will be instructed to turn outwards following the collection of street casualties and deliver them to the nearest prescribed clearing station at the perimeter. These stations may be provided by upgrading selected first-aid posts. In upgrading posts it is probable that staffing and equipment are of more importance than elaborate structural works, for "sorting," resuscitation, and recording are the primary functions of the clearing station. Patients usually travel better before operation than after, and the mass of