Case 24.—This patient appears to have died of a complication of the unconscious state, possibly due to inhalation of infected material, rather than from the meningitis itself.

Case 31 acquired a particularly lethal and rare complication—suppurative pericarditis.

There were no deaths among those who were seen during the first week of illness and who remained free from serious secondary infections of the viscera. For these reasons more than for the simple statistical fact of a low mortality rate on a limited series of cases, I feel justified in believing that sulphonamide E.O.S. has a pronounced curative action in this disease. Very diverse figures have been published by other observers using different remedial agents. This diversity is readily understood by anyone who has worked among these patients. The mortality rate is affected greatly by so many factors other than that of the chemotherapeutic potency of the drug employed. It varies with the epidemic, locality, age distribution, the speed with which the diagnosis is made, the amount of medical attention which can be focused on each patient, and, above all, with the perfection of the nursing care available. Superb nursing skill is required by delirious, comatose, or semiconscious patients. I know that I have been particularly fortunate in my house-physicians and nurses.

The figures given below furnish a suitable basis for comparison.

Epidemic 1914-17 (pre-sulphonamide):
5,306 cases; mortality 65%

These national figures were analysed by H. D. Rolleston (1919), who found in them evidence of a dwindling mortality as serum treatment came into more general use.

Present epidemic, first quarter, 1940 (sulphonamides widely used):
4,388 civilian cases; mortality 23.7%

Individual groups of smaller numbers but more unified medical control have yielded better results:

120 cases observed personally by Banks (1940) (sulphapyridine or other sulphonamides used in all cases):
Mortality 10%

154 cases at Wrexham (*Lancet*, 1940) (various doctors) (sulphonamides widely used):
Mortality 12.3%

The mortality in the present series treated with sulphonamide E.O.S.—34 cases (9%)—is therefore closely in accord with the current figures.

Summary

A new sulphonamide of low toxicity and full chemotherapeutic activity in acute cerebrospinal meningitis is described. Less critical clinical data lead one to expect that it will be equally effective in combating infections due to *B. coli* and haemolytic streptococci. More extended clinical observations will probably reveal other fields of utility. In animals it controls acute infections due to *Str. pyogenes* and pneumococcus Type III, and chronic urinary infection by *B. typhosus* and *B. paratyphosus* A. Its low toxicity has been investigated by experiment on mice. The rarity with which in therapeutic dosage it provokes unpleasant side-effects in man has been demonstrated by a clinical survey of 130 cases. It seldom causes vomiting, and malaise attributable to the drug is inconspicuous. It is hydrolysed in the body with the liberation of sulphanylamide *in situ*, and is excreted in forms which are far more soluble than those following the administration of sulphapyridine or sulphathiazole. It possesses the convenient property of ready solubility in water. Neutral and almost tasteless pharmaceutical solutions can easily be prepared for administration by mouth, rectum, or vein.

REFERENCES

- Banks, H. S. (1940). *Proc. roy. Soc. Med.*, **33**, 555.
Coplans, M. (1940). *J. Path. Bact.*, **50**, 507.
Green, A. G., and Coplans, M. (1940). *Chemistry and Industry*, **59**, 793.
Lancet (1940), **2**, 216.
Leedham-Green, J. C. (1941). *British Medical Journal*, **1**, 586.
Marshall, E. K., Emerson, K., and Cutting, W. C. (1937). *J. Amer. med. Ass.*, **108**, 953.
Rolleston, H. D. (1919). *Lancet*, **1**, 541, 593, 645.

THE SIZE AND VISIBILITY OF THE FILTERABLE OR VIRUS BODIES

BY

ALFRED C. COLES, M.D., D.Sc.
M.R.C.P.Lond., F.R.S.Ed.

Consulting Physician to the Royal Victoria Hospital and Royal National Sanatorium, Bournemouth

The filterable, so-called ultra-microscopic or elementary, bodies, generally spoken of as the viruses, are not easy to examine on account of their very small size and the difficulty of staining them; but fortunately they are nearly always present in enormous numbers, often in almost pure cultures. The measurement of such small bodies is very difficult by eye-piece or screw micrometer, and usually one has to resort to photography.

The Photographic Method of Measurement

Muir and Ritchie (1937) state:

"In the microscopical study of virus particles Barnard has utilized photography by ultra-violet light, which is of shorter wave-length than that of the visible rays, and thus has obtained greater microscopical resolution. By this method he has been able to demonstrate photographically virus particles of 0.075μ in diameter."

Let us compare this with the results obtainable by the use of the ordinary microscope and objectives in common use when these are employed correctly. The late Mr. Edward Milles Nelson (1909), who devoted the whole of his long life to the microscope and was undoubtedly the greatest authority on the subject, says: "Briefly, the method is to bring the object just to the point of invisibility by reducing the N.A. (numerical aperture), and then the correction for antipoint will be the measure of the object." I might add that one can determine the minimum visible of any objective by the following calculation: for white light of 45,300 waves per inch divide 1.041 by the N.A.; for green light of 50,000 waves per inch divide 0.942 by the N.A. of the objective. A full table of the values of the N.A. is given in Mr. Nelson's paper.

I have used this method of measuring by extinction for many years, employing for this purpose a Leitz $1/10^a$ fluorite objective fitted with an iris diaphragm graduated in N.A.s, immediately behind the back lens, and find it remarkably accurate and easy after one has had some practice.

From Nelson's table it will be seen that an objective of N.A. 1.4 will reveal a particle of 0.0744μ with white light and 0.0673μ with a green screen. Nelson later, in 1935, informed me that even greater visibility could be obtained by using a motor head-light bulb (3 amps 12 volts with a transformer, as suggested by Ivey-Cook), with the filament in focus, and a deep blue screen. Then the size of an object seen would be 0.841 divided by the N.A.; so that an objective of N.A. 1.3 would show a particle 0.065μ , and one of N.A. 1.4 a particle 0.06μ , in diameter. From these figures, given in much greater detail in a previous paper (Coles, 1929), ultra-violet microscopy does not seem very urgently needed.

Mr. Eliot Merlin (1929), the distinguished microscopist, in concluding his masterly article, says:

"In the writer's opinion it is of paramount importance that the true optical capacity of the ordinary worker's microscope, furnished with ordinary glass and fluorite achromatic or apochromatic object-glasses, should be clearly understood and not underestimated, this being the tool exclusively used by thousands of laboratory and technical workers all the world over. It cannot be too emphatically maintained that it is an experimentally absolute fact that an object-glass of 0.4 N.A. (i.e., a common

1/2-inch) will, in white light of wave-length of 45,300 per inch, just render visible a particle of 0.262μ in diameter; and similarly the 1/12 oil immersion of 1.30 N.A. in common laboratory use should, if properly used to its full defining capacity, reveal particles of not less than 0.08μ diameter in white light, of 0.0724μ diameter with a screen wave-length of 50,000 per inch, and of 0.057μ with the photographic wave-length of 63,500 per inch."

Stained Films and Filtration Method Compared

In recent times organic membranes such as collodion have been devised by Elford (1931) to test filterability. These are called "gradocol" membrane, and by their means Elford and his co-workers have computed the size of the particles. A list of filterable viruses, with their size, is given by Mackie and McCartney (1938). The unit used to express these minute viruses is $m\mu$ —i.e., one-thousandth part of a micron (μ) or one-millionth part of a millimetre.

When we compare the size of the filterable viruses as determined by microscopical methods on stained films we are at once struck with the extraordinary difference between these and those obtained by filtration methods. The following measurements of the elementary bodies of definite virus diseases will show how much the results differ.

The size of the virus of *variola* and of *vaccinia* as determined by filtration is stated to be 0.15 to 0.175μ . They were not recognized as the causal organisms till Paschen described them in 1906, but as a matter of fact Dr. John Buist of Edinburgh had fully described and photographed them nearly twenty years previously. Buist (1887), after staining them with aniline gentian violet, gave their size as from 0.15 to 0.5μ ; and von Prowazek (1912) gives exactly the same figures. Mr. Nelson and Mr. Eliot Merlin, knowing nothing of these figures, estimated the size of some of them as up to 0.45μ in preparations stained with Giemsa which I sent them in 1922. *Vaccinia* was the first of the filterable bodies that I dealt with in my work on the viruses in 1913.

The size of the virus of *foot-and-mouth disease* is given as 0.01 to 0.02μ , or 10 to $20 m\mu$, and it is regarded as one of the smallest of filterable bodies and quite invisible.

In 1929 Sir Charles Martin, then director of the Lister Institute, kindly sent me films from the vesicles of cases of foot-and-mouth disease, and later sealed ampoules of the filtrate; and in 1933 Dr. (since Sir) John Ledingham, his successor, sent some of the filtrate—both through the kind service of Dr. J. T. Edwards. Films made from the filtrate were fixed in alcohol and stained for twenty-four to forty-eight hours in well-diluted Giemsa. Films were also made with indian ink as in Burri's method, and control films of the ink alone were made at the same time. Lastly, the filtrate was spread in thin film on perfectly clean slides and merely allowed to dry in the air. When these were quite dry they were put under a cover-glass and the margins sealed with cement, taking great care that none of the cement should run in—i.e., they were mounted, unfixed, and unstained in air. It is interesting to note that these preparations and the indian-ink films mounted in 1929 and 1933 are to-day in as good condition as when they were made.

The Giemsa-stained films, when examined with the 1/10-inch oil-immersion objective, show clearly innumerable very fine virus bodies, differing from those seen in other virus diseases, such as *vaccinia*, herpes, molluscum, etc., on which I have been working for the last twenty-eight years, only by their enormous number and their small size. These stained preparations, when examined by dry dark-ground with only the 8-mm. or 1/3-inch apo-objective and an 18 compensating ocular, show the minute bodies exceedingly well. The indian-ink preparations under the oil-

immersion objective reveal the elementary bodies very distinctly, in marked contrast to the control film of ink. The merely air-dried unstained films mounted in air, when examined by dry dark-ground illumination with the same objective, distinctly show the minute bodies, and exclude the possibility of their being colloidal protein molecules or precipitates from the stain.

Mr. Eliot Merlin, who has during the past years examined nearly all my preparations of virus bodies of various diseases, writing in 1929, confirms my findings in every way. He describes the preparations of foot-and-mouth disease as consisting of densely packed strongly stained particles, measuring from 0.15 to 0.3μ in the stained films, and precisely similar bodies were seen in the indian-ink slides. He says:

"Your findings *re* size and appearance are in agreement with the facts as they appear to me, and the presence of the bodies in the unstained, merely air-dried, films is pretty conclusive that what we had seen in the stained films was not due to stain debris. The foot-and-mouth virus seems to me to greatly resemble the vaccine and many other virus bodies which we have examined previously. What surprises me is that the bodies with which the films swarm should be considered difficult to define with the ordinary microscope. I think the appearance of the bodies under the microscope is in itself fairly convincing to anyone with an open mind on the subject. If they be not virus bodies one wonders how they can be proved not to be."

Causal Virus of Various Diseases

The causal virus of *yellow fever* is said to be one of the smallest of the elementary bodies, and according to filtration experiments is estimated to measure 0.018 to 0.027μ , or 18 to $27 m\mu$. Blood films from monkeys infected with yellow fever were kindly sent me by Prof. Edward Hindle in 1929 and Prof. W. Schüffner in 1931, and when these were stained with Giemsa I found minute bodies in the red blood corpuscles and also free in the plasma. They varied considerably in size; many measured from 0.1 to 0.35μ , but some were only just visible specks. The larger bodies were resolvable in most cases into three or four minute spots not larger than 0.1μ . The number seen varied according to the day of the disease; they were most numerous on the second or third day of the fever, and in several of the films many free and intracorpuseular forms could be seen in every field.

In blood films of patients naturally affected and in volunteers inoculated with about 0.33 c.cm. of dried serum of infected blood from cases of *dengue fever*, somewhat similar bodies were found. These were numerous on the first three days of fever, considerably less numerous on the fourth day, and on the fifth and sixth days they were exceedingly few, being found only with difficulty. Blood films from three cases of *sandfly* or *pappataci fever* taken during the first twenty-four hours showed bodies almost exactly like those of dengue. Mr. Merlin kindly examined and confirmed my findings in these three virus diseases, which have been described in detail with photomicrographs elsewhere (Coles, 1937).

Lastly, let us examine another of the extremely small virus bodies, that of *poliomyelitis*, which measures according to filtration experiments 0.008 to 0.017μ , and according to other textbooks from 0.010 to 0.020μ . I have received air-dried films from proved infective filtrate of poliomyelitic monkey-cord suspension from three reliable sources—namely, Dr. J. A. Kolmer, Dr. A. B. Sabine, and Dr. E. W. Schultz, all of the U.S.A.—and after staining these in Giemsa very definite elementary bodies were easily seen. They varied in size from 0.15 to an average of 0.2 to 0.3μ . These have been described and photographed (Coles, 1939).

Discussion

How can the marked difference in size of the elementary bodies as determined by filtration experiments and microscopical methods be explained? It should be remembered that the size of the virus particles as measured by means of collodion filters represents the size of the *smallest* of the living bodies and not that of the average of the stained elementary bodies. Have fixation and staining any effect on the size of the particles? The answer to this is strikingly in the affirmative, and to a much greater extent than most of us realize. In 1929 I stained a film containing *Treponema pallidum* in Giemsa for fifteen to twenty minutes, and after washing and drying examined it with the 8-mm. or 1/3-inch apo-objective by dry dark-ground illumination, without any cover-glass or mounting medium. A spirochaete was very easily found by this means, partly because the eosin red of the Giemsa shows up brilliantly by dark-ground illumination, but chiefly because the difference in its refractive index and that of air is so great. The position of the spirochaete was logged, and ringed with the marker. The oil-immersion objective was then turned on, and the spirochaete was practically invisible; only after prolonged careful search was it seen, and it appeared as the very thinnest and finest spirochaete possible. The slide was placed back in the stain for twenty-four hours, and then it was markedly thicker and very easily seen with the oil objective. A film or a section stained by any of the flagella stains after mordanting, or in silver-impregnated preparations, shows spirochaetes and leptospira as very thick coarse structures, easily seen with quite low-powered objectives.

As the virus bodies are probably very minute organisms, why should the difficulty of examining them be further increased by restricting the examination to them in the living condition? Do we examine sputum for tubercle bacilli, or the diphtheritic membrane or its cultures for the bacilli, in the living unstained preparations? The size of all bacteria is determined in stained preparations, and in the case of the tubercle bacillus in films mordanted by carbolic acid and heat. The size of the tubercle bacillus is generally given as varying from 2.5 to 3.5 μ in length and 0.3 μ in thickness: has anyone detected this bacillus in its unstained living condition, and can anyone even guess at its size in its living state?

The only reason I can imagine why the elementary bodies are studied in their living unstained state is that the very few exponents of ultra-microscopy would have us believe that this is the only correct method: even if this method were adopted the viruses would never be seen but could only be photographed.

Conclusions

I am of the considered opinion (a) that all the filterable virus bodies, including the smallest and the so-called invisible virus of foot-and-mouth disease and that of infantile paralysis, are easily within the range of the ordinary microscope when used correctly; (b) that ultra-microscopical methods are not essential, and their use has so far added little to our knowledge.

REFERENCES

- Buist, J. B. (1887). *Vaccine and Variola: A Study of their Life History*, Churchill London.
 Coles, A. C. (1929). *British Medical Journal*, 2, 91.
 — (1937). *J. trop. Med. Hyg.*, 40, 209.
 — (1939). *Edinb. med. J.*, 46, 627.
 Elford, W. J. (1931). *J. Path. Bact.*, 34, 505.
 Mackie, T. J., and McCartney, J. E. (1938). *Handbook pract. Bact.*, 5th ed., p. 523.
 Merlin, A. A. C. Eliot (1929). *Watson's Microscope Record*, No. 17, p. 14.
 Muir and Ritchie (1937). *Manual of Bacteriology*, 10th ed., p. 717, Edinburgh.
 Nelson, E. M. (1909). *J. roy. micro. Soc.*, Ser. 2, 29, 549.
 von Prowazek, S. J. M. (1912). *Handbuch der pathogenen Protozoen*, p. 143, Leipzig.

CLINICAL MANIFESTATIONS OF TETRYL AND TRINITROTOLUENE

BY

J. HILTON, M.B., Ch.B.

AND

C. N. SWANSTON, M.R.C.S., L.R.C.P.

Medical Officers to a Royal Ordnance Factory

Since many filling factories have been built recently in different parts of the country it might be of interest to practitioners to know something about the effects on the human body of continued exposure to tetryl and trinitrotoluene.

Tetryl, or "C.E."

This compound is a sharp crystalline powder, yellow in colour. Its clinical manifestations are as follows:

1. The powder stains the skin of the hands, face, neck, and also the hair, and this colour deepens to orange on exposure to sunlight. It should be emphasized that the staining varies considerably in different workers and bears no relation whatever to any of the following manifestations.
2. New workers sometimes complain of a sharp tingling sensation in the nose. This causes sneezing and more rarely epistaxis, and is due to the irritating effect of the crystals on the nasal mucous membrane.
3. The commonest complaint from tetryl workers is dermatitis. This starts as an erythema, followed by a papular eruption accompanied by some exfoliation. It arises in the region of the sweat glands of the face—i.e., the naso-labial fold and submandibular regions of the neck. It may spread all over the face, causing severe irritation, and in some cases gross oedema of the eyelids, which may prevent the patient opening the eyes for two or three days. In severe cases secondary sensitization may occur on the hands, arms, and trunk. Should tetryl come into contact with the conjunctivae a mild conjunctivitis results. Constitutional symptoms are present only in the more severe cases, usually as headaches and general malaise. They are to some extent understandable, as patients suffering from extensive rash often lose sleep. The treatment of C.E. dermatitis is first and foremost removal from contact with the powder. It is not enough merely to change the worker's occupation; he or she must be removed right away from the vicinity of the powder, otherwise the condition will be aggravated by the small amount of dust present in the atmosphere in all shops where tetryl processes are carried on. It is also important that the worker be given a complete change of working clothes. The local treatment, generally successful, consists in frequent applications of calamine lotion or, if there is much desquamation and drying of the skin, lotio calaminae oleosum. A word of warning is, however, necessary when using the latter preparation: tetryl workers should be told on no account to use oily preparations on their skin as a preventive measure, as by so doing ideal conditions are produced for the development of a second attack of C.E. dermatitis. Mild cases clear up in a few days, but severe rashes often persist for two or three weeks. Suspension from work is seldom necessary. When the dermatitis has quite cleared up the worker may return to contact work, but if a second or third attack occurs it is an indication that the patient is very susceptible and should be removed permanently from tetryl work. In our experience those workers who develop oedema of the face and periorbital tissues are bad risks, and we remove them permanently from contact at the first outbreak of the rash.
4. Although it is not generally admitted that tetryl gives rise to any constitutional symptoms, we have found that numbers of new workers complain of epigastric pain, with nausea or vomiting. These symptoms, like those of T.N.T. toxic gastritis, bear no relation to food and are seldom severe enough to keep the patients away from work. Such workers are removed from contact until the symptoms abate. They are then allowed to return, and usually no further trouble arises.

Trinitrotoluene, or T.N.T.

T.N.T. is a yellowish crystalline powder more stable than C.E. and is used either by itself or mixed with other sub-