s-methyl isothiurea sulphate and observed no special difference as a consequence of taking the substance.

ITS POSSIBLE VALUE IN CASES OF SHOCK

Some observations have been made by me in man and in experimental animals to ascertain if this substance might be of value in the treatment of surgical shock. It has been noted in about twenty obviously moribund patients, mostly medical cases with terminal circulatory collapse, that the systolic blood pressure may rise promptly from, say, 60 to 120 mm. after the intravenous administration of 0.2 to 0.4 gramme. The diastolic pressure is raised proportionally. The blood pressure remains elevated for fifteen minutes to one hour. The heart rate, however, may fail to slow or does not diminish to the degree observed in health. Some of these results were encouraging in that several moribund patients made a temporary recovery. Two cases of severe surgical shock were treated by the intravenous injection of s-methyl isothiurea sulphate. Both patients survived, and improvement coincided with the administration of the substance. It is regretted that most of the clinical observations were not made on cases of surgical shock, but opportunities for a satisfactory study of this subject are lacking in New Zealand.

More definite evidence has been obtained that the substance is of value in treating acute shock produced experimentally in healthy animals. Thus crushing of limbs and extensive abdominal injuries were employed in anaesthetized cats and dogs to reduce the level of the blood pressure to 30 or even 20 mm. Hg. Ordinarily when the blood pressure has been reduced to such levels cessation of the heart beat usually follows within a minute or two. A majority of animals with blood pressures of 20 or 30 mm. Hg were revived for periods of one or more hours when treated with s-methyl isothiurea sulphate intravenously. It seems, therefore, that the substance delays death if given to animals in extreme traumatic shock. There is thus a chance that the substance would be useful to tide over air-raid casualties which cannot be removed to a place with facilities for transfusion.

The substance also may be worthy of clinical trial in those cases of traumatic shock in which a vasoconstrictor substance could be of value, but even so it should be regarded as an adjuvant and not as an alternative to methods of treatment which restore the blood volume. Patients whose blood pressures remain at a low level despite restoration of the blood volume might be benefited by s-methyl isothiurea sulphate.

In my patients dosage has been controlled by observing the pulse and blood pressure. It appears that in divided doses as much as 1 to 1.5 grammes may be administered in the course of three hours, and probably more would be tolerated. Solutions have been sterilized by tyndallization and kept in ampoules, 0.2 gramme in 2 c.cm. So far only one contraindication to the use of this substance has been found—namely, that it should not be employed in congestive heart failure.

F. H. SMIRK, M.D., F.R.C.P.,

Professor of Medicine, University of New Zealand, Dunedin.

Malachite in the Control of Bilharzia

The possibility of controlling bilharzia in Southern Rhodesia has been under investigation for the past two years. This work was begun at the instigation of the medical director of Southern Rhodesia, Dr. A. P. Martin, with the aid of a grant from the trustees of the State lottery. I proceeded to the colony in the first place in company with Sir Malcolm Watson, director of the Ross Institute of Tropical Hygiene. Certain results of this work seem to be important enough to merit immediate announcement.

An essential phase of the investigation has been a search for effective and practical means of destroying the intermediary hosts of human bilharzia—namely, the snails *Physopsis globosa* and *Biomphalaria pfeifferi*. Previous attempts to destroy these dangerous snails have centred largely around the use of copper sulphate. The principal objections to the use of this substance are its relative toxicity to man and other mammals, its great solubility (approximately ten parts of copper per hundred), and its high cost. Copper carbonate is at least as lethal to snails as is copper sulphate, it is much less soluble than the sulphate (approximately one part of copper per million), and in saturated

solution it has been proved by experiment in Southern Rhodesia to be non-toxic to white rats when given in repeated doses. The relatively high cost of copper carbonate as well as of the sulphate has led to a search for some less expensive form that would be suitable for control work on a large scale.

Malachite (mineralized basic copper carbonate) when ground fine enough to pass through a 200-mesh sieve has been found to be suitable for this purpose. In this condition malachite from two sources in Southern Rhodesia has been proved to be effective in destroying snails both in the laboratory and in actual bilharzial pools in the field. The present cost of malachite in many parts of Africa is less than £1 a ton. Finely powdered malachite yields an effective concentration (dissolved and colloidal) of 0.5 part of copper per million when placed in natural water (from the Macabuzi River, near Salisbury, S.R.). It has been found that the solubility of this substance is considerably increased by mixing it with fresh brewer's waste (i.e., fermented barley) or with the powdered pods of a common Rhodesian tree, *Swartzia madagascariensis*.

These methods of destroying snails will play an important part in a campaign for the control of bilharzia in Southern Rhodesia which will begin shortly. A full account of the control methods to be used is in preparation.

I am grateful to the Medical Director of Southern Rhodesia for permission to publish this note.

Salisbury, Southern Rhodesia.

ALAN MOZLEY, D.Sc Ed.

Melanotic Carcinoma: Sudden Spread after Hysterectomy

The following case may be interesting enough to be worth recording.

CASE REPORT

The patient, aged 41, had hysterectomy performed for fibroids of the uterus on April 25, 1941. Her only complaint was menorrhagia. At operation a routine inspection of the abdominal contents revealed no other abnormality. Pre- and post-operative examination of the chest showed no disease, and the patient left hospital after a normal convalescence. The pathological report on the uterus indicated simple myoma only.

The patient was readmitted on August 1 complaining of abdominal swelling and vomiting for two weeks; for one week she had been breathless. Symptoms had come on quite suddenly twelve weeks after operation. On admission she was cyanosed and breathless at rest; there were scattered subcutaneous black nodules, including three in the abdominal scar; the abdomen was distended with fluid, but a large nodular liver was readily palpated. Search for a primary melanoma revealed only a small scar over the left clavicle. The patient then gave a history that five years ago she had had a "black naevus" there. It was treated with carbon dioxide snow, but increased in size—whereupon it was excised under local anaesthesia. There had been no recurrence since.

She died on August 7. At necropsy melanotic carcinomatous deposits were found in almost every organ of the body—heart, thyroid, lungs, meninges, and brain. The liver weighed 110 oz., and there were profuse deposits in the mesentery, kidneys, spleen, suprarenals, pancreas, and stomach. The abdominal scar was studded with melanotic deposits. The urine gave a positive test for melanogen.

Twelve weeks previously there had been no macroscopical abdominal deposits. If the sudden spread of melanotic deposits was purely coincidental the rapid dissemination after five years' complete quiescence is noteworthy. It is, however, interesting to speculate whether the operation on a distant site initiated the spread; whether, in fact, the malignant cells had lain dormant for five years only to be stirred into activity by the mechanical or physiological disturbance of the operation. Although so far as I know no mechanism for this is described, it seems to be a more possible explanation.

My thanks are due to Mr. Edgar Freshman, under whom the case was admitted, for permission to report it, and to Dr. Norman Ashton, who carried out the necropsy, for the use of his report.

Kent and Canterbury Hospital.

LESLIE J. TEMPLE, F.R.C.S.