on July 29, two days after an indwelling catheter was resumed, and on July 30 his pulse was a steady 72 and his temperature normal. The catheter was removed on July 31. Even before its removal fluid injected into the bladder by the catheter returned freely by the urethra outside the catheter. On July 31 a rigor occurred with a temperature of 105° and an acute right orchitis. With

Drawing of right half-prostate of Case 2; natural size, weight 1½ oz.

Dr. G. S. Haynes's concurrence sulphapyridine was given; the temperature fell in forty-eight hours, and subsequent convalescence was uneventful. Urine was passed naturally and easily after August 1, and the patient describes the stream as "better than normal."

The specimen after removal was an almost globular mass 4½ cm. in diameter, and weighing 1½ oz. On section

the mass was a typical adenoma without cysts. On the left aspect was visible, adherent to but separable from it, a smooth elastic band representing the right half of the prostatic urethra. Above, this broadened out to terminate in a torn edge at the urethral outlet of the bladder. Below, it narrowed to a point short of the verumontanum, where the urethra had been half torn across (see Fig.).

BIBLIOGRAPHY

- Cabot, H. (1936). In *Modern Urology*, original contributions by American authors, edited by H. Cabot, vol. I, Kimpton.
Freyer, P. J. (1902). *Stricture of Urethra and Enlargement of the Prostate*, 2nd ed., Baillière, London.
Richardson, W. G. (1904). *On the Development and Anatomy of the Prostate Gland*, Churchill, London.
Riches, E. W. (1936). *British Medical Journal*, 1, 578.
Thomson-Walker, Sir J. W. (1936). *Genito-Urinary Surgery*, 2nd ed., Cassell, London.
Wardill, W. E. M. (1941). *Lancet*, 2, 127.

METHODS FOR THE LOCAL APPLICATION OF SULPHANILAMIDE

BY

FRANK HAWKING, D.M.

(From the National Institute of Medical Research, Hampstead)

Earlier papers have described the antibacterial effects of local applications of the sulphonamides to experimental wounds infected with the organisms of gas gangrene, and the concentrations in the blood and in the tissues around the wound resulting from such local applications (Hawking, 1941a, 1941b, 1941c). During this experimental work various practical methods have been devised, and these are here described in the hope that they may be helpful to those employing local applications of sulphonamide compounds to human lesions. For the sake of conciseness most references to other workers have been omitted. The compounds used have been either sulphanilamide or sulphathiazole.

Powder

In shallow open wounds this is the simplest form in which to apply the substance. It can be distributed by means of a spoon, a pepper-pot, or an insufflator. Whichever method is used, enough powder should be applied to cover the whole surface liberally, and yet not enough to form large concretions, which take a very long time to dissolve. Gently massaging the powder into the tissues with a spatula will help to get it effectively distributed and dissolved. For use in the operating theatre on fresh wounds it is convenient to have a supply of the powder in 5-gramme lots, put up either in paper packets or in screw-capped bottles. In order to obviate the very slight

risk of fortuitous contamination of the powder, during its manufacture or afterwards, by tetanus or other spores of pathogenic bacteria, it can be sterilized in such packets or bottles by dry heat at 150° C. for one hour. (If sterilized in the autoclave with the cap unscrewed the powder is apt to become crystalline and is no longer suitable for use in an insufflator.) For routine ward work the use of unsterilized powder would seem to be justifiable.

Gauze

In narrow deep sinuses which are packed at regular intervals with ribbon gauze the powder can be inserted into the depths on the packing. This can be done most simply by pouring the powder on to the gauze as it is pushed in, but a better way is to incorporate the powder on to the gauze beforehand. For this purpose small glass pots with a metal screw cap, such as are used for ointments, are taken. A wide slot is cut in the lid. The pot and the gauze are sterilized separately. The gauze is then passed through a thick paste of sulphanilamide and water under sterile precautions and is placed in the pot, the end of the ribbon protruding through the slot. The top of the pot is covered with a piece of sterile oiled silk. For use in the wards the ribbon is pulled out as required. If the gauze is autoclaved *after* smearing with the paste most of the sulphanilamide forms powdery crystals, which cease to adhere to the ribbon.

Tablets

It is difficult to transfer powder to the bottom of a deep sinus. The best method in these cases is the use of small tablets made of the compressed compound—e.g., tablets containing about 0.2 gramme of material and measuring about 8 mm. across. Such tablets were first suggested by Dr. A. S. Parkes; they were kindly prepared by the British Drug Houses Ltd. and May and Baker Ltd. The tablets can easily be inserted into the desired position by sinus forceps, and they are then crushed to hasten disintegration; eventually they are completely absorbed and leave no residue which might act as a foreign body. With sulphanilamide, which is fairly soluble, tablets of the pure compound will probably be dissolved in a reasonable time, although it is advantageous to crush them in order to accelerate the process. With sulphathiazole, tablets of the pure compound (uncrushed) were found to persist in a sinus for over two days without much visible change. It is better to use tablets consisting of 0.1 gramme of sulphathiazole and 0.1 gramme of glucose, the glucose being added in order to promote the disintegration of the tablet; when these are crushed after insertion most of the material is dissolved in twenty-four hours. Tablets could also be used in a wound when it was desired to secure a steady supply of sulphonamide over a prolonged period; by choosing tablets of an appropriate size the duration of action could be varied as required. The surface of the tablets can be sterilized by dry heat at 85° C. for one hour; heating to 115° C. causes the tablets which contain glucose to turn brown. This degree of sterilization will not kill spores, but the position is the same as that described for the powder.

Pastes

The results described in this section form the preliminary part of a wider programme, which will be described in more detail when it has been completed. Briefly, the ideal sulphanilamide paste for application to wounds, burns, etc., should have the following properties:

1. It should be satisfactory pharmaceutically—i.e., stable, non-odorous, non-rancid, etc.
2. Its physical consistency should be suitable for the required purpose; burnt surfaces might require a thinner preparation than that used for plugging wounds.

3. It should comprise only materials which are available under present war conditions. Thus, since cod-liver oil, which forms the base of two of the pastes described below, is now in short supply, a search for a suitable substitute forms part of the wider programme now in progress.

4. Sulphanilamide should be liberated from the paste at the optimum rate. Presumably the concentration of sulphanilamide in the fluid in contact with the paste should be as near saturation as possible, provided that: (a) The concentration is not so high as to injure growing tissues or to paralyse leucocytes. Jacoby, Medawar, and Willmer (1941) found that fibroblasts and macrophages are killed by a saturated solution of sulphanilamide (concentration 1:60), are temporarily depressed by concentrations of 1:150, and are not affected at all by concentrations of less than 1:1,000. According to these findings, it might be advantageous if the paste exerted a slight restraint on the rate of solution of the sulphanilamide. (b) The amount absorbed is not so great as to produce a toxic concentration in the blood; probably this will not occur except when the areas to be dressed are large; it can be decided only by blood estimations of treated patients. (In a private communication Colebrook reports that one patient who was treated by sulphanilamide paste for a very extensive burn became cyanotic, and the concentration of compound in the blood was 8 mg. per 100 ml.) The whole question is under investigation.

It may here be noted that mixtures of sulphanilamide and liquid paraffin appear to be in use in some hospitals. According to experiments in this laboratory sulphanilamide is insoluble in paraffin and cannot diffuse across it. In mixtures of paraffin and sulphanilamide, however, the particles of sulphanilamide usually sink to the bottom of the paraffin layer until they come into contact with the underlying watery fluids, in which they dissolve. Hence, if sulphanilamide suspended in paraffin is injected into rabbits subcutaneously it is absorbed almost as quickly as if it were suspended in saline. This occurs because the mixture is unstable and in the body it separates into its component parts; if this separation were prevented—for example, by using petroleum jelly—little or no absorption would presumably take place.

5. The paste should be non-irritant to the tissues and innocuous to leucocytes. It should not delay healing; rather, if possible, it should stimulate it.

The complete solution of all these problems will take a considerable time, and meanwhile casualties occur which require treatment. Therefore this short description is given of the best pastes which have been found to date so that they may be available in hospitals, and clinical experience may be gained which may help to orientate the further researches in hand. For the pharmaceutical side of the present work I am indebted to the British Drug Houses Ltd., who kindly co-operated with their experience in these matters and devised pastes of different properties for my trial. The physiological action of these pastes was tested by injecting them between the muscles of rabbits and into the peritoneum of rats; the irritation which they caused was not appreciably greater than would have been produced by any foreign body in these positions. Three pastes are provisionally recommended to suit three different types of lesion.

1. *Sulphanilamide, 30% ; oleic acid, 1.4% ; triethanolamine, 0.35% ; cetyl alcohol and beeswax, each 1.4% ; water to 100%.*—This is a thin paste to spread over inflamed skin surfaces—e.g., impetigo—in order to apply sulphanilamide with as little other material as possible. It can be used on granulation tissue instead of powdered sulphanilamide, but if inserted into deep wounds it leaves a residue. The sulphanilamide is absorbed from this paste almost as rapidly as when the powdered compound is applied as such. This paste was kindly tested by Dr. A. Glucksmann of the Strangeways Laboratories, who applied it every other day to standard wounds in rats. He found that it caused very

slight irritation, as shown by stimulation of collagen regeneration and slight retardation of epithelial regeneration; but this irritation was so slight as to be unimportant compared with bacterial infection.

2. *Sulphanilamide, 30% ; cod-liver oil, 49% ; oleic acid, 3.5% ; triethanolamine, 1.05% ; cetyl alcohol, beeswax, and other ingredients to diminish rancidity, 3.5% ; water to 100%.*—This is a paste of medium consistency. The absorption of sulphanilamide from it proceeds slightly more slowly than that which occurs when the compound is applied as a simple powder. Besides supplying a large depot of sulphanilamide, which will be taken up by any serous exudate to form an almost saturated solution, a paste of this kind provides an oily protective layer and will also prevent sticking of dressings. This paste was designed especially for the treatment of burns. Colebrook (personal communication) has observed the rapid disappearance of haemolytic streptococci following its use in such injuries; he also found that it is unnecessary to change the dressings more often than every two or three days. Robson and Wallace (1941) have reported very satisfactory results in burns treated with a similar paste made up with albucid (*p*-aminobenzenesulphonacetamide), which is considerably more soluble than sulphanilamide and which accordingly may exert a greater effect upon the less sensitive organisms such as staphylococci and *Ps. pyocyanea*.

3. *Sulphanilamide, 30% ; cod-liver oil, 60% ; beeswax, 10%.*—This is a stiff paste, to be used on gauze for packing wounds, in the same way as vaselined gauze is used. Its consistency can be modified by varying the proportions of cod-liver oil and beeswax. It provides a non-irritant, non-adhesive mechanical support to the sides of the wound, and at the same time gives off a supply of sulphanilamide, liberation of the compound being slow. It tends to become rancid, however, unless preventive measures are taken.

Summary

Methods are described for the local application of sulphanilamide: (a) as a powder to open wounds; (b) as tablets or on gauze to deep sinuses; (c) as various kinds of paste to treat inflamed surfaces, burns, and wounds which must be packed.

Details are given of three pastes which may be recommended for use in three different types of lesion.

Acknowledgments are due to Colonel L. Colebrook for inspiration and advice; to the surgical staff of the New End Hospital, Hampstead, for facilities; to Miss M. Kuck for technical assistance; and to the British Drug Houses Ltd. for co-operation in devising and preparing the pastes.

REFERENCES

- Hawking, F. (1941a). *British Medical Journal*, 1, 263.
 — (1941b). *Ibid.*, 1, 511.
 — (1941c). *Lancet*, 1, 786.
 Jacoby, F., Medawar, P. B., and Willmer, E. N. (1941). *British Medical Journal*, 2, 149.
 Robson, J. M., and Wallace, A. B. (1941). *Ibid.*, 1, 469.

M. B. Gordon, N. H. Solomon, and S. F. Pearlman (*J. Pediat.*, 1941, 19, 76) record their observations on 680 scarlet fever patients aged from 10 months to 41 years, most of them being between 8 and 21 years of age. They classify these cases into the three groups of mild, moderate, and severe. The mild cases were divided into two groups: (a) controls and (b) those treated with sulphanilamide. The moderate cases were divided into four groups: (a) controls, (b) those treated with sulphanilamide, (c) those treated with scarlet fever antitoxin, and (d) those treated with sulphanilamide plus scarlet fever antitoxin. The authors' conclusions are as follows: (1) Sulphanilamide reduces the incidence and severity of complications. (2) Scarlet fever antitoxin reduces the length of the febrile period to a further extent than sulphanilamide, but it is not so effective as sulphanilamide in reducing the incidence of complications; the combination of scarlet fever antitoxin and sulphanilamide offers better results than sulphanilamide or scarlet fever antitoxin alone. (3) The toxic effects of sulphanilamide are not grave enough to contraindicate its use in scarlet fever.