SURGICAL ECONOMICS IN WAR*

BY

Brigadier C. MAX PAGE, D.S.O., F.R.C.S.

In the organization of the Army the important part played by the medical services in maintaining the health of the troops is generally recognized. The possibility of keeping large armies in the field without undue wastage is accepted as being one of the striking achievements of preventive medicine of this century. This is effected by preventive inoculation, sanitation, and personal hygiene, and these subjects remain a primary and essential responsibility of the Service. When other methods are looked for whereby the medical branch can conserve manpower, it is apparent that, broadly considered, the efficient working of every department of the R.A.M.C. conduces to this end, and each branch has its special problems. I propose to consider in detail the responsibility from the surgical aspect. I will not refer to the handling of battle casualties, but to the less obvious and dramatic saving of man-power which can be effected by the management of the individual soldier suffering from diseases or disabilities that turn up apart from fighting. The views I express are personal, and are coloured by my observations of the results of various surgical work carried out in the Army and Ministry of Health Service during the past few years. The ideas put forward are not original, and the actions, advice, and attitude suggested are already operative in many centres. They are set forth here with the prime aim of giving the medical officer new to the Service a basis of action suited to our present needs.

The General Objective

A dogmatic definition of the procedures that will achieve an aim, the successful pursuit of which must in the main depend on an attitude of mind, is by no means easy. In medicine agreement between a group of doctors on the treatment and disposal of an individual case is seldom a simple problem. In peacetime the propriety and expediency of various methods of treatment are allowed to range over a wide field. In war the urgent need for the prevention or limitation of disabilities and for speed and efficiency in getting men back to work after illness requires no emphasis, nor does the importance of retaining a man in his unit, so far as this is possible, once he has become a trained and useful member of it. If every medical officer realizes his personal responsibility in the matter and considers his actions and their repercussions in this light something will be achieved. I will discuss the problem from the point of view of the unit M.O., the M.O. in charge of a reception station, and the operating surgeon.

The Unit Medical Officer's Work

The unit M.O. has an important part to play in the scheme, although the wide dispersal of troops complicates his work in war conditions. His position in relation to the men of his unit may be compared to that of a first-class general practitioner to his patients or, perhaps more appositely, with that of a whole-time school doctor. By means of periodic inspections he should be familiar with the physical condition and have some knowledge of the psychological attitude of all the men in his charge. He should be fully acquainted with the details of their routine work and training, and should take an interest in their recreations and study their intimate living conditions. By taking an active part in exercises and training, the M.O. will learn for himself the strain to which the troops are exposed, and will so acquire knowledge that may assist in preventing breakdowns; he will also in this way more easily win the co-operation of the officers and the confidence of the men.

Turning to matters of detail, it will be the responsibility of the medical officer to see that the unit is not carrying men unfit for the duties it is expected to perform. It is uneconomic to retain men in a job for which they are unsuited from physical or mental causes. Cases of acute disease or serious injury should clearly be evacuated promptly as they arise to the proper hospital, the location of which should be known, and with which personal contact should be made whenever possible.

* A lecture delivered at an R.A.M.C. Depot to a group of newly engaged medical officers.

If, in routine inspections, the M.O. should detect such defects as varicose veins, bubonocoele, or deformed toes, he should not be in too great haste to send the man for a specialist's opinion. This caution should certainly be adopted when a man does not himself complain of a defect, and is able to perform his full duties in spite of it.

It is in relation to the minor disabilities of the serving soldier that the M.O. will have the greatest opportunity for conserving the strength of his unit. Foot troubles represent an important proportion of this group. Prevention in these matters is clearly better than cure. He should make full use of the chiropodist and see for himself that boots and anklets fit properly. He should order such modifications to be carried out by the unit boot repairer as are possible or desirable. It should be noted that if a man has a foot defect which has proved no handicap in civil life the provision of light boots may sometimes solve the difficulty. The opinion and advice of a near-by orthopaedic surgeon will often help him in this respect. The M.O. should see that cases of foot-strain or tenosynovitis due to over-intensive training are rested before the condition becomes chronic. It must be recognized that the company officer is responsible for the care of his men's feet, but the M.O. should not on that account fail to take his part of the responsibility. A certain number of minor injuries may well be treated within the unit, but if it appears that recovery would be more rapid under better-equipped conditions the man should be transferred to hospital. The same remarks apply to cases of pyogenic infection. Cases of palmar whitlow in particular are best treated in hospital.

When a man is sent to hospital, whether for treatment or for opinion, the importance of sending notes bearing on the case may be emphasized. The M.O. should remember that the patient's history sheet does not always reach the consultant, nor does all the information which may be helpful. Reliable observations on the man's past medical history and on his value as an individual in the unit may be of considerable importance to the consultant if a proper decision on his treatment and disposal is to be made.

The Reception Station M.O.

Under quiet conditions the reception station can form an important buffer against hospitalization. Men may be retained for 10 days in these stations if they are not in need of specially skilled nursing and are likely to be able to return to their units within that period. At this stage an opportunity for some minor surgery arises. Certain nominated stations are now equipped to carry out injections for varicose veins. The selection of cases for this procedure is important, and observations on this point will be found in the *A.M.D. Bulletin*. Recent knee injuries should only be retained when the evidence suggests that there is no gross displacement of the meniscus or serious tear of a ligament. The accurate diagnosis of these injuries is of great importance, and further reference to the subject will be found below.

Reception stations will receive a good many cases of fibrositis. For localized cases injections of 1% novocain will often prove helpful and can do no harm. Hot-air baths can be arranged with a little resource either by extemporizing electric cradles or by the use of a paraffin stove. Cases of acute tonsillitis and laryngitis should not be automatically sent for examination by an E.N.T. specialist when the acute symptoms have resolved; unless there is definite evidence of chronic disease the man should be returned to duty.

The Surgeon's Responsibility

The surgeon can conserve man-power either by the advice he tenders or by his methods of operative management. In the presence of acute disease and serious injury, where active treatment is imposed, one can only suggest the importance of early rehabilitation once the acute phase is past.

It is in relation to what may be termed "cold surgery" that surgical economy or extravagance can be practised. One does not ask the surgeon abruptly to alter his peacetime views on technique. One asks him, however, to be persistently critical of his methods, and to assess so far as possible whether by his operation he is rendering the soldier more fit for his job with the least expenditure of time, or whether the soldier could

by some adjustment carry on an active life without it. The argument will be clearer if some practical examples are given.

Hernia.—A large number of hernias are being operated on in this country, and the results are not as good as might be expected. The proportion of recurrences is high, and the evidence available suggests that in many cases this is due to defective technique. An operation on a soldier or industrial worker for hernia, if effectively performed, should in general render him an A1 man. These operations should, therefore, be scrupulously carried out by an approved method and by an experienced surgeon. In the after-management of these cases graded physical exercises are valuable in restoring a man's efficiency in the minimum time. Cases of bubonocoele due to direct hernia, certainly in men showing a pronounced supra-inguinal bulge, should not be operated on. The condition does not carry any risk of bowel strangulation and can be controlled with a truss. Though the wearing of a truss relegates a man to a "B" category it does not prevent him from taking a useful part in some branches of the Army.

Appendicectomy.—When this operation is decided on in the "cold" period the evidence leading to the decision should be carefully sifted. I have seen a good many men submitted to this operation on account of right-sided pain who have been found subsequently to have a stone in the kidney or a hydronephrosis. Removal of the appendix for what is termed "appendix dyspepsia" should not be undertaken too readily.

Varicose Veins.—I would suggest that a mild degree of this condition does not involve any material disability for Army life any more than it does for a civil occupation, and that, therefore, treatment might often be deferred until after the war.

Operations on the Feet.—It has been generally recognized that operations on the bones of the soldier's foot seldom improve his capacity or category. The practice of cutting the foot to fit the boot was long ago found to be unsatisfactory. The use of less drastic measures and the fitting and modification of footwear, so far as is possible in the Service, deserve more attention than they are given. If a man has "got on" in civil life with a minor or even a major foot deformity, it will be found better to avoid operation except in specially selected cases.

Derangements of the Knee-joint.—Troubles with this joint are the cause of a good deal of short- and long-term disability in the Army. This source of wastage will continue so long as the troops play football. The minor cases may be quickly restored by a short period of rest and pressure followed by progressive, active use of the limb. In the more severe, treatment without accurate diagnosis is little better than a gamble. It is essential to differentiate clearly between damage to ligaments and a cartilage tear and displacement. The taking of a careful history of the circumstances and mechanism by which the injury was caused is of great value here. Tears of a ligament are produced when the leg is in the extended position by forced lateral deviation or hyperextension. Displacements of the menisci are caused when the knee is bent and a rotatory strain is applied. Operation is only justified if history and physical signs point to a tear or displacement of a cartilage. The site of the lesion should be determined with precision. It may be noted that the localization of pain and tenderness on the medial side is closely related to the position of the lesion, whereas with an injury of the lateral meniscus pain and tenderness are less well localized and are generally referred to the posterior aspect or centre of the joint. Routine x-ray examination should never be omitted or else cases of osteochondritis dissecans and other less common conditions will be missed. Operation duly carried out on a torn or displaced meniscus gives a high percentage of good results. If it is performed without a precise diagnosis being made and no gross lesion of the cartilage is found the outcome is usually unsatisfactory. In my experience the removal of both menisci (or parts of them) often leads to a joint which is not fully stable and seldom allows a man to be graded A1.

Fractures.—In these cases, which form so large a part of the surgeon's wartime practice, one can only recommend a critical attitude of mind. Does the method employed impede the speed of repair? Does it allow return of functional activity at as early a period as possible? Is everything done that can be done to avoid prolonged stiffening of joints in relation to the

fracture? Most surgeons agree that overtraction or "distraction" of the fragments at a sensitive phase of the repair process is responsible for delay in union. How far alternative methods of treatment lead to better results must be determined by the individual. The reduction and fixation of a fracture are mechanical processes, and the problem is therefore subject to many solutions as regards the exact method employed.

Ear, Nose, and Throat Operations.—I am here on delicate ground and I will limit myself to a few remarks. Operations for the removal of tonsils and for deflected nasal septum very often result in a man being "Y" listed. There have been wholesale attacks on soldiers' tonsils in certain areas, whereas in other localities with similar military populations such operations are few and far between.

In conclusion, I would say that the views which I have sketched in a wide field may well be subject for discussion, and I must admit they often lack precision, but I would emphasize that my main aim has been to induce an attitude of mind which will assist our Service to do its full share in saving man-power and man-hours at the present time.

I should like to acknowledge my indebtedness to Major-General S. Kyle for advice in assembling these notes.

ORGANIZATION FOR CANCER TREATMENT

MEMORANDUM BY RADIUM COMMISSION*

The National Radium Commission has been established 12 years and during that time has gathered much information about radiotherapy and the organization of radiotherapeutic centres. The most important part of the work of such centres is the treatment of cancer, and of late years the Commission has extended its observation, wherever possible, to cover the whole field of cancer. As a result, the Commission has been impressed by the fact that the death rate continues to rise although the centres with which the Commission deals treat a constantly increasing number of cases, with a slow but persistent improvement in the number of patients who survive several years after treatment.

Some Remediable Factors

Among the remediable factors which contribute to this paradox are:

(1) *Late diagnosis.*—This is due to (a) unsatisfactory conditions under which many patients are seen for the first time—e.g., ill-equipped consulting-rooms and inadequate facilities for examination and note-taking; (b) inexperience of cancer on the part of some medical men who see only a few cases a year and recognize them only when they are far advanced; (c) ignorance and fear of the disease among the laity, which prevent the victim from seeking advice in the early stage.

(2) *Scarcity of beds.*—This is a factor affecting the treatment of all hospital patients, but it has a more evil effect on the cancer patient than almost any other. In most hospitals cases fall into three classes: (a) acute emergencies, surgical or medical, which are admitted instantly; (b) patients admitted by a member of the honorary staff and generally admitted to his own beds; (c) patients who need admission for investigation or for treatment that can be postponed. These are placed on a "waiting list" and often have to wait a very long time. Under this heading are included a large number of cancer patients, and while on the "waiting list" their disease steadily progresses.

(3) *Failure to select the most appropriate treatment for each case.*—This is due to the absence in most hospitals of any regular and organized system for consultation, before treatment is started, between the various specialists whose skill may be needed during that treatment.

(4) *Inefficient treatment in inadequately equipped centres.*—This is due to the natural ambition of every hospital to carry out all kinds of treatment, and want of co-operation between medium-sized and small hospitals to build up joint centres in those specialties in which the needs are more elaborate than their means allow.

A Cancer Organization on a Regional Basis

The National Radium Commission puts forward the following suggestions for the organization of the treatment of cancer on a regional basis.

General.—The precise form of the administrative arrangements on which a cancer organization should be built up must depend in great measure on the general arrangements for the post-war hospital service which the Government has promised.

* Owing to shortage of space the memorandum has been abridged.