side the posturing is repeated. It is necessary, however, to perform the whole manoeuvre with as little delay as possible, as the oil becomes rather more viscid as it cools. Alternatively, the oil may be rendered thinner by the addition of olive oil (4 to 6 c.cm. of olive oil to 20 c.cm. of neohydriol or lipiodol), it being introduced without any preliminary heating. This, however, does not always give satisfactory results, as the thinner oil runs in the smaller bronchi and is there retained. By this posturing the patient is spared reintroduction of the needle into the neck, and excellent pictures can be obtained (Figs. 1 and 2)—A.-P. and lateral of the side first filled, A.-P. and oblique after filling of the second side.

Subsequently, whatever method is used, the patient is told to expectorate as much as possible and not to swallow any sputum. If necessary he should posture to get rid of iodized oil from any cavities where it may have lodged. Oil seen on screening in an iodine-sensitive patient's stomach should be removed by stomach wash-out or administration of emetics. A patient who has received cocaine should be warned not to eat or drink for three hours subsequently.

Discussion

The crico-thyroid route has been used merely because I get the best results with it, and because cocainization can be cut down to a minimum in all patients and omitted in many. Sensitivity below the vocal cords to injections seems to vary greatly, and many patients can control their cough reflex without cocaine anaesthetization (Jewesbury, 1939). Be that as it may, transnasal, translingual, or other routes may be used if preferred. The importance is stressed not of the site of introduction of the oil but of the areas reached after introduction. A lateral view of the affected side is essential, and, if posturing be faulty, oil on the opposite side will render this view confused, one bronchial tree being superimposed on the other in the film. To fill the middle lobe properly is also essential,

FIG. 1.—Radiograph showing outlines of the left side of chest.

and lateral views are again necessary. The hard-pressed radiologist should be spared the pain of reporting yet again: "Those few basal bronchi outlined are normal."

Summary

The importance of adequate posturing on bronchography is stressed, and a standard technique is described which allows delineation of all parts of the bronchial tree. Without such

technique, imperfect filling of the bronchial tree will be the rule instead of the exception.

My thanks are due to Drs. W. E. Lloyd and W. H. Coldwell for their kindness and help, and to the Sister and staff of the x-ray department, Westminster Hospital.

FIG. 2.—Right side outlined by posturing the patient. The oil from the left side has been made to run over and outline the right.

REFERENCES

- Erwin, G. S. (1936). *Lancet*, 1, 1236.
Jewesbury, E. C. O. (1939). *British Medical Journal*, 1, 772.

THE EFFECT OF BLEEDING ON THE BLOOD-SUGAR LEVEL IN BLOOD DONORS*

BY

B. J. LAWRENCE, M.B., Ch.B.

AND

G. PLAUT, M.D.

The object of the present investigation was to determine whether the level of the blood sugar is related to the incidence of "fainting" in blood donors. The clinical characteristics of such faints have been previously described by Brown and McCormack (1942). Certain factors responsible have been analysed by a statistical survey (Greenbury, 1942). In the latter it was suggested that lack of food might be a predisposing cause. If a constant low blood-sugar level occurred in donors who fainted, as opposed to those who did not, it would favour such a hypothesis.

Method

In a series of 50 unselected donors blood-sugar estimations were made upon samples of capillary blood obtained from the ear immediately before and immediately after the withdrawal of between 400 and 450 c.cm. of blood by the Medical Research Council technique (Brown and McCormack, 1942). In a second series of 46 unselected donors blood-sugar estimations were made upon venous blood taken at the end of a routine bleeding from the antecubital vein. In these cases a pressure of between

* A report to the Medical Research Council from the N.W. London Blood Supply Depot.

60 and 80 mm. Hg had been maintained by a sphygmomanometer cuff on the arm for 3 to 5 minutes before the sample was obtained. During this period, in order to facilitate the blood flow, the donor had continuously clasped and unclasped a rolled bandage held in the palm of the hand. All samples were collected in small bottles containing dried oxalate (equivalent to 1 c.cm. of a mixture of 1.2% ammonium oxalate and 0.8% potassium oxalate). Owing to the nature of the transfusion service it was not possible to standardize conditions of temperature, previous exercise, or the time of the last meal. The majority of donors had had a meal within 3 hours of venesection.

The method used for the estimation of glucose was Harding's modification of the Schaffer-Hartmann method as described by King, Haslewood, and Delory (1937).

Results

In the series of 50 unselected donors examined before and after bleeding only 2 fainted. Their blood-sugar levels were 85.8 mg. and 98.6 mg. per 100 c.cm. before and 85.8 and 100.92 after bleeding. The findings in the other 48 donors are summarized in Table I. No change in blood-sugar level was found

TABLE I.—*Capillary Blood-sugar Values*

Time	No. of Donors	Blood-sugar Values in mg. per 100 c.cm.	
		Range	Mean
Before bleeding ..	48	60.32 to 100.40	94.91 $\pm$ 6.12
After bleeding ..	48	67.30 to 147.32	100.67 $\pm$ 6.56

in 13 instances; in 27 there was a slight increase after bleeding, the average figure being 16.34 mg., while in 10 there was a decrease of the same order. In the second series of 46 unselected donors in whom the level of the blood sugar in venous blood obtained at the end of bleeding was determined, 16 fainted either while being bled, so that the full amount could not be withdrawn, or immediately afterwards. The results are summarized in Table II. Again the figures recorded reveal no

TABLE II.—*Venous Blood-sugar Values*

Donors	No. of Donors	Blood-sugar Values in mg. per 100 c.cm.	
		Range	Mean
Group 1 (normals)	30	28.01 to 106.42	66.14 $\pm$ 7.14
Group 2 (faints) ..	16	50.11 to 100.33	77.27 $\pm$ 5.90

significant difference in the blood-sugar level of donors who faint and those who do not.

There was no apparent relation between the age of the donor and the blood-sugar level. One extremely low figure of 28.01 mg. occurred in an old man of 73 who had had a good breakfast about 2½ hours before bleeding and who felt perfectly well. No correlation could be established between the time that had elapsed since the last meal, the blood-sugar level, and the incidence of fainting.

Discussion

The figures for the blood-sugar level recorded in the present investigation suggest that a low blood sugar is not a factor associated with fainting in blood donors, and that withdrawal of 500 c.cm. of blood does not result in any definite change in the blood-sugar level. The level in the first series, made upon capillary blood, was higher than that in the second series, made upon venous blood—a finding in agreement with that of other workers (Jonas, 1933; Himsworth, 1933; Shoji *et al.*, 1938). Shoji and his co-workers found that the contraction of the forearm muscles decreased the sugar content of venous blood considerably in non-fasting subjects. Such contraction of muscles occurs in all donors, as they are asked to clasp and unclasp their hands during the bleeding procedure in order to facilitate the blood flow. It is probably responsible for the rather wide difference observed.

Summary

The blood sugar was estimated in 96 donors, 18 of whom fainted. No significant difference was found in the level of the

blood sugar after bleeding. This level was without effect upon the incidence of fainting.

REFERENCES

- Brown, H., and McCormack, P. (1942). *British Medical Journal*, 1, 1.
 Greenbury, C. L. (1942). *Ibid.*, 1, 253.
 Himsworth, H. P. (1933). *Clin. Sci.*, 1, 1.
 Jonas, L. (1933). *J. clin. Invest.*, 12, 139.
 King, E. J., Haslewood, G. A. D., and Delory, G. E. (1937). *Lancet*, 1, 886.
 Shoji, R., Shima, K., and Sasaki, T. (1938). *Jap. J. med. Sci. III Biophys.*, 5, 41, 42

CONDITION OF RETAILED RAW MILK IN AN URBAN AREA

BY

C. A. GREEN, M.D., Ph.D., D.P.H.

Surgeon Commander, R.N.V.R.

The incidence of tuberculosis of the bovine type in the human subject is regrettably high in this country. Bendixen *et al.* (1937) recorded aggregate figures which showed that the percentage frequency of this type in cervical gland infections at all ages was 46.2 and 52.3 in England and Scotland respectively. The corresponding figures in bone and joint infections were 22 and 29.8; in lung infections, 0.9 and 5.5; in lupus, 48.6 and 53.8; in meningeal infection, 25.7 and 29.8. Lesions due to the bovine type in the human subject result, in the majority of instances, from the ingestion of milk from infected cattle. It is difficult to obtain exact figures as to the incidence of infection in dairy cattle, but approximately 40% react to the tuberculin test and about the same proportion of cattle are found infected on slaughter in abattoirs. That raw milk should suffer as a result is a natural consequence. Thus Brownlie (1933) found that 10% of churn samples of retail milk in urban areas in Scotland contained tubercle bacilli.

Contamination of milk supplies by an organism such as the tubercle bacillus, whose virulence is capable of easy demonstration both in the experimental animal and in the unfortunate human patient, rightly has greater claim to attention than the general cleanliness of the milk, by which term is meant freedom from bacterial contamination, irrespective of the type of organism present. Nevertheless, questions of both economic and public health importance depend upon this latter quality.

The following data were recorded as the result of an investigation into the raw milk supplies of an urban community in which it was estimated that 50% of retailed milk was of this type.

Methods

A sample of milk measuring 300 c.cm. was taken into a sterile container from the raw milk exposed for sale in the retailer's premises on the morning of delivery and brought immediately to the laboratory for examination. Three tests for cleanliness were then applied. Two of these were the plate-count and presumptive *B. coli* tests, recommended in the Milk Special Designations Order, 1936, Scotland. The third test was the methylene-blue reductase test, introduced under the Milk Order, 1936, England and Wales, as a substitute for the plate-count test. In addition, many of the milks were examined for the presence of tubercle bacilli. The methods used are, in brief, as follows:

1. *Plate-count Test*.—Decimal dilutions of milk in 90 c.cm. of diluent were prepared. From the 1/100 and 1/1000 dilutions three separate 1-c.cm. volumes were plated in Wilson's (1935) yeastrel agar medium. After incubation at 37° C. for 48 hours colonies were counted by means of a hand lens and tally counter.

2. *Presumptive B. coli Test*.—Three tubes of MacConkey's fluid medium were each inoculated with 1 c.cm. from the 1/1000 dilution of milk. The presence of acid and gas after 48 hours' incubation at 37° C. was noted.

The standard upon which the quality of samples was judged by the above tests was that demanded of standard milk by the