induced in an animal by the injection of saponin. It seems permissible to assume that these chemical changes, and perhaps also others as yet unknown, affect the glandular system without producing any appreciable anatomical alteration. They naturally vary according to the nature and degree of the disorder.

Through all the portals of entry employed the rays have access to the sympathetic nervous system. Radiotherapy therefore can modify the function of the sympathetic, and also of its perivascular plexuses, in various ways according to the dosage and the degree of excitability of the nerves. Gouin and Bienvenue produced by irradiation of the sympathetic the same reactions in the blood and the same clinical and therapeutic phenomena as Leriche produced by surgery. Moreover, besides the endocrine glands, which secrete their products into the blood, P. Masson has demonstrated the existence of endocrine tissues which secrete directly into the nerves. These two actions, humoral and nervous, proceed together. For instance, adrenaline acts by exciting the sympathetic nerves. Comparative anatomy and embryology show the close association of chromaffin with the sympathetic, and experimental physiology shows the close interaction of certain hormonal influences with corresponding neural factors—"sympatheticoglandular synergy." It is therefore easily conceivable that radiotherapy applied to the sympathetic and glandular systems may excite modifications which produce the beneficial effects that have been noted. As yet no one knows why or how it restores a disturbed balance by acting on that phase of the reciprocal vegetative function which happens to be impaired. This and many other difficult questions remain to be answered, and the answers will be as interesting as they are valuable.

TREATMENT OF PHOSPHORUS BURNS

BY

A. THELWALL JONES, M.D.

A notable increase in the incidence of phosphorus burns must be expected under war conditions, and the article by E. W. Godding and H. E. F. Notton in the *Journal* of April 4 (p. 433) is therefore timely and useful. There are several points in this article which deserve further comment. It should be stated that the "official treatment" which they critically examine has been based on the practice gradually established through long clinical experience within the chemical industry and is the result of careful observation over many cases. These authors are under a misapprehension on one minor issue, as they assume that the copper sulphate treatment is designed chemically and completely to destroy the phosphorus particles left in the affected area after the preliminary washing. This is not the primary object, which is rather to destroy minute particles and to make obvious, by blackening, the larger pieces so that they can be removed mechanically. There is no present clinical evidence that the concentrations, both of alkali and of copper sulphate, suggested "officially" are inadequate, and indeed the extant case histories indicate that they are well chosen. Godding and Notton have done well, however, to stress that the alkaline treatment must be thorough, and the importance of this feature is borne out by my practical experience. A novel problem associated with present conditions is the satisfactory treatment of phosphorus burns under field or Service conditions. Here the practice found good enough in industry would often be difficult to apply with the requisite speed, and a satisfactory field dressing would be a valuable contribution to the casualty services. The powders suggested by Godding and Notton may prove of value in this connexion, and should certainly be given a thorough trial on human cases; it may be possible to get the requisite experience industrially. One further caution is necessary: Godding and Notton experimented with material from S.I.P. (self-

igniting phosphorus) grenades, and this is not strictly comparable to the phosphorus as encountered in industry, though the effects may reasonably be assumed to be similar. The following notes on the treatment of phosphorus burns may be of interest:

Characteristics of the Burns

Burns caused by phosphorus have long been recognized as particularly severe and troublesome lesions. They have these characteristics: (a) the initial pain is severe; (b) a greyish area is produced almost immediately at the site of the lesion, and blistering is rare; (c) sepsis is particularly apt to ensue; (d) healing is prolonged, and the resulting scar is often thin and atrophic. During the past ten years I have had the opportunity of studying burns arising from a variety of chemical irritants and corrosives. Generally, the severity of burns is proportionate to the chemical nature of the causative agent, and may be placed in the following order: (1) hydrofluoric acid, (2) phosphorus, (3) sodium sulphide, (4) caustic soda, (5) sulphuric and other acids.

Method of Treatment

The essential factor in the treatment of all chemical burns is the immediate flooding of the area affected with water or sodium bicarbonate solution. It is necessary to organize an adequate first-aid service for this purpose. Buckets of water, running water, eye-wash bottles, and, in special cases, baths must be instantly available. Experience has indicated that immediate flooding with water and *prolonged soaking in warm sodium bicarbonate solution* is the essential feature in treatment.

Altogether 65 cases of burns caused by phosphorus have been observed, the records of which are set out in the accompanying Tables. (In every case the time taken to heal is recorded as a period which elapsed before the employee was able to return to normal work without any dressings.) Several methods

Cases treated with Sodium Bicarbonate, Copper Sulphate, Acriflavine Emulsion (1/1,000), and Boroseptic Ointment

Age	Date of Burn	No. of Burns	Area Affected	Degree	Total Size (sq. in.)	Days Absent from Work	Days taken to Heal	Comments
22	24/1/39	4	Fingers	3	2½	Nil	30	
34	14/3/39	1	R. forearm	1	8	"	17	
34	14/5/39	1	Eyelids, face	2, 3	9	Hosp. 28	17	
29	29/6/39	1	Heel	3	6	30	22	
46	8/6/39	1	Ankle	3	4	16	10	
27	9/7/39	4	Wrist, fingers	2	2	Nil	30	
36	14/7/39	4	Neck, chest	2, 3	5½	"	49	Skin eruption with acriflavine
36	14/10/39	1	Forearm	3	1½	Hosp. 81	79	Sensitive thin scar. Skin eruption with acriflavine
42	26/10/39	1	Finger	3	1½	Nil	19	
41	12/11/39	1	"	3	½	"	17	
42	16/11/39	1	Palm	3	½	"	4	
30	17/11/39	2	Fingers	3	½	"	18	
25	13/12/39	1	Arm	3	6	"	25	
34	18/12/39	4	Hand, fingers	2, 3	1½	"	17	
—	10/1/40	2	Fingers	2	½	"	12	
46	6/2/40	4	"	2	6	"	4	
32	8/2/40	1	Hand	2	1	"	4	
37	24/2/40	1	Forehead	2	½	"	9	
28	7/3/40	1	Finger	3	½	"	11	
36	14/3/40	1	Hand	2	½	"	7	
38	15/3/40	1	Forearm	3	1½	"	23	
46	28/3/40	1	Finger	3	½	"	12	
36	2/4/40	1	"	3	1	"	18	
35	10/4/40	3	Fingers	3	1½	16	24	
34	25/4/40	1	Forehead	3	4½	Nil	60	
68	13/5/40	2	Forearm	3	1½	"	24	
47	15/5/40	1	Wrist	3	1	"	16	
42	23/5/40	2	Fingers	3	1	"	8	
29	2/6/40	1	Hand	3	4	"	35	
29	2/7/40	1	Finger	3	½	"	20	
36	4/8/40	3	Arm, forearm	3	32½	"	32	Skin eruption with acriflavine
45	21/11/40	1	Palm	2, 3	½	"	18	
38	25/11/40	1	Thumb	3	1½	"	27	
24	25/11/40	2	Wrist, hand	3	1½	"	39	Given light work
25	28/11/40	3	Wrist, fingers	2, 3	3	"	33	" "
37	29/11/40	1	Thumb	2	½	"	8	
—	4/12/40	3	Fingers	3	2	"	24	

Cases treated with Sodium Bicarbonate, Copper Sulphate, Picric Acid, and Boric Ointment

Age	Date of Burn	No. of Burns	Area Affected	Degree	Total Size (sq. in.)	Days Absent from Work	Days taken to Heal	Comments
35	10/11/33	1	Foot	2	1	Nil	8	Sepsis
36	13/11/33	1	Eyelid	3	1	"	17	
37	14/11/33	1	Thumb	2	1	"	7	
47	15/11/33	1	Forearm	3	1	"	12	
35	20/11/33	2	Fingers	3	1	"	13	
42	25/11/33	1	Finger	3	1	"	12	
34	28/11/33	3	Hand, fingers	2	1	"	5	
33	30/11/33	2	Fingers	2	1	"	7	
37	1/12/33	1	Hand	3	2	"	21	
35	4/12/33	1	Wrist	3	1	"	10	
—	25/8/35	2	Fingers	3	1	"	18	Sepsis
—	28/8/35	3	"	3	1	"	15	
34	2/9/37	1	Hand	3	1	"	42	

*Cases treated with Sodium Bicarbonate, Copper Sulphate, and Burnogel**

Age	Date of Burn	No. of Burns	Area Affected	Degree	Total Size (sq. in.)	Days Absent from Work	Days taken to Heal	Comments
28	20/2/37	2	Hand, thumb	3	3	Nil	19	Sepsis—ichthyol substituted
41	1/3/37	1	Hand	3	5	27	31	
45	28/5/37	1	Forearm	3	1	Nil	19	Pain ichthyol substituted
43	3/9/37	4	Arm	3	2½	"	29	
38	12/11/37	1	Knee	3	30	12	28	Ichthyol used when healing
36	17/11/37	1	Calf, leg	3	5	Nil	43	"
29	12/2/38	1	Thumb	3	½	"	15	"
36	5/5/38	1	L ft heel	3	2½	32	35	Thin scar. Sepsis—ichthyol substituted
40	17/5/38	6	Fingers, thumb, cheek	3	6	Hosp.	43	Thin scar. Off work 60 days
35	17/10/38	1	Finger	3	1	Nil	18	Pain after 2 days—ichthyol substituted
34	15/11/38	1	Hand	3	12	51	59	
38	10/6/39	1	"	2	1½	Nil	10	Ichthyol required
39	20/8/39	1	Ankle	3	6	36	48	

* Copper guaiacol sulphonate, 0.2%; acriflavine, 0.05%; benzamine borate, 0.1%; perfume and jelly base, ad 100%.

Cases treated with Sodium Bicarbonate, Copper Sulphate, and Tannic Acid

Age	Date of Burn	No. of Burns	Area Affected	Degree	Total Size (sq. in.)	Days Absent from Work	Days taken to Heal	Comments
40	18/1/37	2	Two fingers	3	1½	Nil	32	Painful
21	3/2/37	1	Hand	2	½	24	5	"

of treatment have been tried, and as a result of the experience gained the following procedure is advised:

- (1) Immediately flood the affected area with water.
 - (2) Thoroughly flood and wash with warm sodium bicarbonate solution (roughly two tablespoonfuls to a pint of water). Pick off with forceps any obvious particles of phosphorus disclosed by examination in the dark.
 - (3) Swab the affected area with 1% solution of copper sulphate and make a further exploration for phosphorus particles so revealed. Soak the affected area (immerse if possible) in warm sodium bicarbonate solution for a prolonged period. The duration of this soaking depends upon the size and depth of the burn—i.e., small burns 1/2 hour, large burns 1 to 2 hours.
 - (4) Dress daily with acriflavine emulsion 1/1000. Burns near the eyes are best treated by a mask of lint soaked in sodium bicarbonate solution renewed four-hourly.
 - (5) Immediately healing begins (7 to 10 days) boroseptic ointment (P. D. & Co.) provides a very satisfactory dressing, and its use can be continued until the scar produced is sound.
- Tannic acid and other coagulants do not give satisfactory results, as is also the case in the treatment of all chemical burns. Acriflavine emulsion has the disadvantage that in some

individuals local and generalized skin eruptions may be produced. A daily inspection will detect such lesions at an early stage, when boroseptic ointment should be substituted.

As a result of such treatment results are obtained which compare favourably with a similar series of burns caused by caustic soda and sulphuric acid. Healing is slower and scars are thinner in burns caused by phosphorus.

RELATION OF PYOGENIC SKIN INFECTIONS TO THE SKIN-CARRIER RATE

BY

N. H. MARTIN, B.M., M.R.C.P., F.I.C.

Major, R.A.M.C.

Bacteriological investigations have demonstrated that a fraction of the presumed healthy population carry potentially pathogenic staphylococci on their wrists. The actual proportion of carriers varies from 5% found by Nimmo-Smith (1941), examining R.A.M.C. personnel, to 19.5% found by Gillespie, Devenish, and Cowan (1939), examining students. Devenish and Miles (1939) appeared to show that a previous history of pyogenic skin infection was not given by a significantly higher percentage of skin carriers than normal.

The Present Investigation

The object of the investigation was to discover whether or not the carrier rate varies with the part of the body sampled, and if there was a significantly higher carrier rate in people suffering from pyogenic skin infections as judged by multiple swabbing. The series investigated consists of group A, 140 patients suffering from pyogenic infections of the face or neck, and group B, 100 individuals having no demonstrable pyogenic lesions. An effort was made at first to ascertain whether or not individuals of group B had at any time suffered from pyogenic skin infections. This was abandoned because it was clear that they were not equally reliable as witnesses. So far as the inquiry went there seemed to be no evidence of a significantly higher carrier rate among those giving a previous history of skin infection.

Technique.—Three swabs were taken from each person. The first swab was from the upper lip in group B and from the upper lip where possible in group A, but in every instance at least one inch from the nearest demonstrable lesion. The second swab was taken from the right wrist in right-handed people and from the left wrist in left-handed people. The third swab was taken from the chest in the midline opposite the second rib. At each site an area of 2 sq. cm. was sampled with a broth-impregnated swab which was stroked over that area 30 times in 30 seconds. The swabs were immediately plated out on blood agar and on gentian-violet agar at 37° C. for 24 hours. The plates were then examined, and selected colonies of staphylococci were subcultured and tested for their ability to form coagulase. At the same time their ability to form pigment was noted. The coagulase test was chosen as a basis for assessing pathogenicity; and it was observed that 5 out of 151 strains isolated from group A failed to produce any pigment though they were coagulase-positive. McFarlan (1938) showed that a number of non-pathogenic strains of the staphylococcus would ferment mannitol, and this test was not carried out. The plates containing gentian violet were used for the isolation of streptococci.

Discussion

The results are shown in tabular form. There is in the series under review a significant increase in the "complete carriers" of potentially pathogenic staphylococci in group A. Though this may be, and probably is, a transient phenomenon, the carrier rate appears to be extensive. A comparison of the carrier rate for the face, wrist, and chest in each group shows that in group B it varies, being highest, as might be expected, on the face. In group A the variation though present is not so marked, the difference between face-carrier rate and chest-carrier rate being 13.2%, compared with 24% in group B.