be the most important factor in determining the amount of protein entering the circulation after haemorrhage and also in deciding the rate at which this addition is made. If the stores are depleted completely, then the plasma protein concentration after haemorrhage will be less than that expected, as the protein requirements of the body must be met from the circulating plasma protein. The gap between expected and observed values will continue to widen unless new protein is added direct to the blood by blood or plasma transfusions or is synthesized from the breakdown products of protein taken by mouth.

If the adequate stimulus to movement of protein into the blood stream is a low plasma protein content of the blood, then some correlation might be expected between the degree of protein deficit (difference between plasma protein values before and 30 minutes after haemorrhage) and the amount of protein entering the blood stream at, say, the end of 1 hour after haemorrhage, indicated by the difference between the expected and observed plasma protein values. Such a correlation does not exist. Consequently, it would seem that either hypoproteinaemia is not the adequate stimulus or, alternatively, such stimulus is effective only when protein reserves exist outside the blood stream. The balance of evidence so far would suggest that the latter view is more likely to be correct. It is not suggested, however, that the protein in the plasma stores exists precisely in the form of the albumin-globulin complex present as plasma protein in the blood. The speed with which protein does appear in the blood stream and the amounts contributed after haemorrhage lend strong support to Whipple's view that the precursor of the plasma proteins in the plasma protein stores requires very little change before it can be discharged across the cell membrane into the blood stream.

The state of the protein reserves of the body may be determined by the response to plasma transfusions. After allowing 15 minutes for thorough mixing of the infused plasma, a blood sample is taken, followed by further samples at 45 and 75 minutes after the end of transfusion. The movement of the haemoglobin values is determined, and from these are calculated the expected plasma protein value for the 45-minute and 75-minute samples. If the observed values are greater than those expected it can be concluded that the movement of protein from blood to body cells is slow, and presumably protein stores are adequately filled. If the expected and observed values are equal or approximately equal, then protein-free fluid only is moving out of the blood stream. If the expected value is greater than the observed value, then protein movement from the circulation is rapid, and unless there are signs of massive protein movement into the tissue fluid—e.g., oedema—it must be presumed that the plasma protein stores are low. Such results give a useful indication of the state of the protein stores.

Conclusion

It is clear from these experiments that after haemorrhage the restoration of blood volume is independent of the restoration of plasma protein. Restoration of both concurrently would imply considerable protein stores. The speed of blood volume restoration in the cat apparently always outstrips the rate of plasma protein movement into the blood stream. In man the data of Wallace and Sharpey-Schafer make it clear that even after large haemorrhages (1,150 c.cm.) the plasma protein concentration can remain unchanged, although blood dilution is proceeding rapidly. One subject (19) apparently restored his blood volume after a haemorrhage of 750 c.cm. within 4 hours, and at the end of that time had increased his plasma protein concentration from 6.6 to 6.9 g. per 100 c.cm. This subject had a blood volume of 4 litres before haemorrhage and one of 4.2 litres after haemorrhage.

After haemorrhage and subsequent dilution of the blood, a blood transfusion is followed by a rapid increase in haemoglobin concentration. This haemococoncentration is not pathological but is due entirely to fluid movement out of the blood stream. The restoration of the blood volume to its pre-haemorrhage level takes place independently of a movement of erythrocytes into the blood stores or of plasma proteins out of the circulation. A movement of erythrocytes out of active circulation after blood transfusion would give an expected plasma protein value lower than the observed value. The fact that no animal showed such excess of observed plasma protein over

that expected even more strongly suggests that there had been rapid movement of plasma protein out of the circulation.

Summary

There is evidence of movement of plasma proteins into the blood stream within 2 hours after haemorrhage in cats anaesthetized with nembutal.

Blood transfusions at varying times after haemorrhage are followed by movement of plasma proteins out of the circulation within 1 hour of completion of transfusion.

Plasma transfusions after haemorrhage are followed by plasma protein movement out of the circulation within 1 hour of completion of transfusion.

The rate and amount of plasma protein movement into the circulation are independent of the amount or rate of bleeding, the amount or rate of dilution by fluid from the tissue spaces, and blood-pressure changes.

The amount of plasma protein contributed to the blood, and therefore the plasma protein concentration at any time after haemorrhage (provided no food containing protein has been administered), is probably dependent on the state of the plasma protein stores outside the circulation.

A method of testing the adequacy or otherwise of plasma protein stores is suggested.

We wish to thank Drs. Wallace and Sharpey-Schafer for permission to use the data in Table VII, and for allowing us to inspect some of their unpublished results.

REFERENCES

- Barbour, H. G., and Hamilton, W. F. (1926). *J. biol. Chem.*, **69**, 625.
Ebert, R. V., and Stead, E. A. (1941). *Amer. J. med. Sci.*, **201**, 655.
Madden, S. C., and Whipple, G. H. (1940). *Physiol. Rev.*, **20**, 194.
Price, P. B., Hanlon, C. R., Longmire, W. P., and Metcalf, W. (1941). *Johns Hopk. Hosp. Bull.*, **69**, 327.
Robertson, J. D. (1935). *J. Physiol.*, **84**, 393.
Robinson, H. W., and Hogden, C. G. (1940). *J. biol. Chem.*, **135**, 707, 727.
Stead, E. A. (1940). *J. clin. Invest.*, **19**, 776.
Wallace, J., and Sharpey-Schafer, E. P. (1941). *Lancet*, **2**, 393.
Weech, A. A., Reeves, E. B., and Goettsch, E. (1936). *J. biol. Chem.*, **113**, 167.

CIRCULATORY OVERLOADING FOLLOWING RAPID INTRAVENOUS INJECTIONS

BY

E. P. SHARPEY-SCHAFFER, M.R.C.P.

AND

JOHN WALLACE,* B.Sc., M.B.

WITH TECHNICAL ASSISTANCE OF

A. LATHAM AND A. C. PINCOCK

(From the Department of Medicine, British Postgraduate Medical School)

Authorities differ widely in advising the amount of serum or blood to be injected and the rate at which it should be given. Many stress the dangers of overloading the circulation. In 1894 Bayliss and Starling showed that rapid injection of crystalloid solutions or blood caused a rise of venous pressure, observations which have been amply confirmed by others in the experimental animal (Bainbridge, 1915; Wiggers, 1921). Meek and Eyster (1922) showed in dogs that the immediate effect of an intravenous injection was a rise of venous pressure associated with an increase in the diastolic size of the heart; these effects were only transitory, and there was evidence that the peripheral capillaries and venules were acting as reservoirs. There have been fewer observations in man. Eyster and Middleton (1924) investigated 4 normal subjects. In the case showing the most pronounced change 600 c.cm. of blood was given at the rate of 43 c.cm. a minute; this produced a rise in venous pressure of 2 cm. H₂O but no increase in the size of the heart. Caughey (1935) found no rise of venous pressure in normal sub-

* Working for the Medical Research Council.

jects receiving 1,500 c.cm. of saline at the rate of 50 c.cm. a minute, while a rise of up to 6 cm. H₂O was produced in some patients with heart disease. No attempt was made to measure the degree of retention of injected saline in the circulation. Altschule and Gilligan (1938) gave from 500 to 1,500 c.cm. of crystalloid solutions to 35 subjects at rates of 6 to 71 c.cm. a minute. The venous pressure was raised more than 4 cm. H₂O in only 6 subjects, pulse and blood-pressure changes were small, the cardiac output was increased, but there was no change in vital capacity. This paper reports the results of rapid intravenous injection of saline, serum, and, in a few cases, blood on the venous pressure and certain other cardiovascular measurements.

Methods

The observations were made on convalescent subjects without cardiovascular disease. The conditions of study have been given elsewhere (Sharpey-Schafer and Wallace, 1942). The haemoglobin was estimated on 0.02 ml. freely flowing ear samples by a photo-electric method (Hill and Pincock, 1941). Venous pressures were estimated in the antecubital vein by a continuous citrate method (Wood, 1940), and changes were checked by direct observation of the neck veins (Lewis, 1937). Up to 45 minutes may have to be allowed for the venous pressure to reach a steady level. Vital capacity was measured on a Benedict-Roth spirometer. Radiographs of the heart were taken in the sitting posture, with the tube at 6 ft. The exposure was 1 second to obtain the diastolic heart size. Differences in inclination or respiratory excursion are indicated in the text. Usually two control and several post-injection films were taken. The right pectoral-F or the right pectoral-R leads (Wood and Selzer, 1939) were employed in the electrocardiographic studies. The following solutions were used: (1) 0.9% saline; (2) normal strength reconstituted lyophil serum; (3) 4-times-concentrated reconstituted lyophil serum; (4) citrated blood.

Results

The results are shown in Table I. When the rate of injection was between 50 and 150 c.cm. a minute the venous pressure began to rise after about 1,000 c.cm. of saline or

serum had been given, and reached a maximum toward the end of injection (Fig. 1). If the rate of injection was very rapid, as little as 500 c.cm. raised the venous pressure. The subject shown in Fig. 2 received the first 500 c.cm. of serum

FIG. 1.—Case 1. Venous pressure and Hb% change after saline. Zero represents control level of venous pressure or Hb%.

FIG. 2.—Venous pressure and Hb% change after serum in Case 11a.

TABLE I

Case	Age and Sex	Amount Injected (c.cm.)	Rate of Injection (c.cm./min)	Initial Hb %	Immed. Hb % Fall	V.P. Rise (cm. H ₂ O)	Pulse		Vital Capacity		E.C.G.	Symptoms		
							Before	After	Before (Litres)	After (Litres)				
0.9% Saline														
1	43 M	2,000	91	96	20.0	10.5	90	60	2.3	2.3	0	Headache Slight constriction in chest. Headache		
2	39 M	1,800	112	114	21.0	10.0	75	75			0			
3	16 M	2,000	91	89.5	19.5	8.0					0			
4	17 M	2,000	90	98.5	19.5	6.5					105		105	T change
5	20 M	1,500	75	90	11.2	6.0					84		96	0
6	17 M	2,000	168	116	12.0	6.0	75	75	75	75	0			
7	56 F	2,000	104			4.0	92	88						
8a	54 M	2,000	100			0	75	75						
9a	21 M	300	20			95	5.7	0						
4%														
Serum														
10	44 M	1,620	80	64.5	15.5	11.0	75	105	3.8	3.4	Slight T change Slight T change 0	Slight constriction in chest. Headache Headache		
11a	17 M	1,600	114	102.3	23.5	10.5			3.0	2.3			0	
12a	52 M	2,000	83	104	20.7	9.0			90	105			2.9	2.4
8b	54 M	700	100	110	11.5	9.0	75	75	3.5	2.9	0	Constriction in chest. Headache Headache Severe constriction in chest		
13a	59 M	2,100	154	95	18.0	8.5	60	60			0			
14	28 M	1,200	95	68.5	18.3	8.0	75	90			0			
15	19 M	1,200	54	104.8	19.3	8.0	80	84	3.36	3.0	0			
16	46 F	1,400	108	62.8	5.3	4.0					75			75
17	61 M	1,500*	60	71		8.0	75	75			3.36	3.0		0
18	40 M	1,080*	71			4.5	75	105	4.3	4.0	T change			
Concentrated Serum														
19a	59 M	300	30	84.2	10.6	10.0						Headache " "		
9b	21 M	300	23	95	19.9	6.0								
11b	17 M	300	21	93.2	11.6	4.0								
20	44 M	300	25	66.6	9.6	2.0								
21	29 M	300	25	78	0.5	0								

* Blood.

at a rate of 250 c.cm. a minute. All subjects given saline or serum showed a rapid fall of venous pressure once the injection had been stopped. Saline often left the circulation quickly, as indicated by a rising haemoglobin percentage (Fig. 1). The venous pressure still fell in a matter of minutes, however, if the haemoglobin remained at the lower post-injection level

fall of blood pressure and syncope in the upright posture after sodium nitrite is preceded by a fall in venous pressure. Injection of serum immediately after venesection restored venous pressure, blood pressure, or postural response to normal. When giving intravenous injections at rates of 100 c.cm. or more a minute considerable pressure must be maintained in

TABLE II.—*Injection after Venesection*

Case	Age and Sex	Initial Hb%	Amount of Venesection (c.cm.)	Rate of Venesection (c.cm./min.)	Hb% Change directly after Venesection	Amount Injected (c.cm.)	Rate of Injection (c.cm./min.)	Immed. Hb% Fall	V.P. Change (cm. H ₂ O)	Symptoms
0.9% Saline										
22	31 M	108	900	90	-3.0	1,650	138	15.5	0	
23	37 M	101.5	740	105	-8.5	1,800	140	18.5	+2.5	
24	57 M	108	1,000	71	+1.0	1,500	88	9.0	0	
Serum										
12b	52 M	104.7	1,100	100	-8.0	2,000	77	33.5	0	Fall of B.P. Syncope
13b	59 M	84	1,080	81	-4.0	2,100	84	20.9	+4.0	
11c	17 M	104.4	1,100	36	-5.8	1,600	80	20.6	0	
8c	54 M	106.3	740	146	-1.0	700	140	10.8	0	
25	53 F	95	900	112	-4.0	1,300	60	24.6	0	Fall of B.P. Syncope
19b	59 M	83	1,080	71	0	300*	43	17.3	0	

* Concentrated serum.

(Fig. 2). One subject given blood showed a rise of venous pressure that persisted for 50 minutes. When the blood volume was first reduced by a large venesection, injection of serum or saline in similar quantities and at similar rates produced little or no change in venous pressure (Table II and Fig. 3). The subject of Fig. 3 showed a small fall of venous pressure

the bottle of fluid; measurements of venous pressure in the opposite antecubital vein might therefore merely reflect this transmitted pressure. That this is not so, and that a rise of venous pressure is due to an acutely increased blood volume, is shown by the following considerations: (1) When the blood volume has not been previously reduced a rise of venous pressure does not occur until about 1,000 c.cm. has been given, at rates of about 100 c.cm. a minute. (2) There is a statistically significant correlation between the rise of venous pressure and the fall of haemoglobin (Fig. 4). (3) There is little or no rise of venous pressure when saline or serum is injected immediately after blood-volume reduction by venesection. (4) If, however,

FIG. 3.—Case 12a and 12b: interval of 10 days between observations.

after venesection, associated with a fall of blood pressure and syncope. In this and other subjects with syncope after large venesections (Wallace and Sharpey-Schafer, 1941) a small fall of venous pressure (1 to 2 cm.) preceded the fall of blood pressure, and there was a further fall of venous pressure with collapse of the circulation. If there is no fall of blood pressure after venesection, there is, however, an abnormal fall on sitting upright. Weiss, Wilkins, and Haynes (1937) showed that the

FIG. 4.—Relation between rise of venous pressure and fall in haemoglobin: coefficient of correlation (r)=0.84.

enough time elapses after venesection for the haemoglobin to fall considerably, transfusion of similar quantities then causes a rise of venous pressure.

Vital Capacity.—The vital capacity, measured within 2 minutes of the end of injection, showed an average fall of 470 c.cm. in 6 of 7 subjects (Table I). McMichael and McGibbon (1939) have shown that there is a fall in the lung volume and vital capacity of normal subjects on assuming the horizontal posture; they ascribed the fall to the presence of more blood in the lungs. Venesection, on the other hand, increases the vital capacity (Glaser and McMichael, 1940). That the fall of vital capacity on overloading the circulation is due to increased blood in the lungs is borne out by the x-ray appearances. None of these subjects showed clinical evidence of pulmonary oedema, and the respiratory rate and tidal air showed no change.

Radiograph of Heart.—Films were taken before and immediately after injection. The results are shown in Table III. The area of the cardiac shadow was measured by a planimeter, and the right, left, and oblique cardiac and transverse chest diameters were also measured. In 5 of the 8 subjects examined the average increase in the heart area was 15.3%. Table I shows

TABLE III.—Radiological Measurements

Case	Heart Area (sq. cm.)		Right Diam. (cm.)		Left Diam. (cm.)		Oblique Diam. (cm.)		Transverse Chest (cm.)		
	B.	A.	B.	A.	B.	A.	B.	A.	B.	A.	
14	107	123	3.5	4.1	8.4	9.4	14.0	16.2	27.0		Diaphragm 1 cm. higher in films after Slight inclination of body forward in films after Diaphragm 1.2 cm. higher in films after
15	92.5	109	3.4	4.5	7.4	8.0	14.0	14.8	27.2		
3	92	105	5.0	5.1	6.9	7.4	14.2	15.1	29.8		
5	83	101	4.1	5.0	6.4	6.9	13.1	14.0	26.6		
6	105	119	3.3	4.1	9.1	9.3	15.5	16.3	23.1		
7	97	101	3.7	4.3	8.3	8.7	15.3	16.2	26.2		
2	93	96.5	4.7	4.5	9.6	9.7	14.2	14.4	27.6		
8b	150	158	4.0	4.7	10.6	10.6	15.9	16.4	29.0		

B. = Before injection. A. = After injection.

that there was a considerable rise of venous pressure and fall of haemoglobin in each of these 5 subjects. Although the latter changes were present in Case 2, there was but little cardiac enlargement. So far as could be judged from postero-anterior films the enlargement appeared to involve all chambers. In addition, the following changes were present after injection: the superior vena cava shadow became prominent in those subjects in whom there was a considerable rise of venous pressure, the pulmonary arteries and vascular markings in the lungs increased in size and density, and the whole lung fields were relatively less translucent (Figs. 5 and 6).

FIG. 5.—Radiograph of Case 14 before transfusion.

Electrocardiogram.—The right pectoral-R arm or right pectoral-L leg electrocardiogram is the chest lead that gives one of the earliest indications of acute right heart stress, as from packed pulmonary embolism (Wood, 1941). Changes in the T-wave are in the direction of inversion when chest leads

are recorded in the manner agreed by the Cardiac Society of Great Britain and Ireland and the American Heart Association (*British Medical Journal*, 1938, 1, 187). Only 4 of 12 subjects showed electrocardiographic changes, the T-wave becoming flattened or inverted. Gilligan and Altschule found increased amplitude of the P-wave in lead II in a few of their cases.

FIG. 6.—Radiograph of Case 14 after 1,200 c.cm. of serum.

There were no P-wave changes in leads RP-R or F in this series. The x-ray findings suggest that enlargement of the heart shadow may involve all chambers in some subjects. Electrocardiographic changes in the direction of preponderance of ventricle would therefore not be expected, since it is well known that cardiac enlargement can take place without them. However, it has not proved possible so far to observe the correlation, if any, between electrocardiographic changes and x-ray appearances after acute overloading of the circulation.

Heart Rate.—The heart rate showed no change or slowing in 9 subjects, while 6 showed an increase.

Symptoms.—Eight subjects had headache; 4 complained of a feeling of constriction in the chest during injection; in 1 it was severe, but not severe enough to necessitate termination of the injection. It passed off rapidly as soon as intravenous fluid was stopped. The most striking sign was vasodilatation; it was present in all subjects with any degree of retention of saline or serum in the circulation. The face became redder in tint, and the peripheral veins became prominent and distended.

Summary

Up to 2,000 c.cm. of saline, serum, and blood was injected into subjects without cardiovascular disease at rates of from 54 to 168 c.cm. a minute. The venous pressure was raised up to 11 cm. H₂O when there was considerable retention of injected fluid in the circulation, as indicated by the fall in haemoglobin. Radiographs showed an increase in the diastolic size of the heart, enlargement of the pulmonary arteries, and prominence of the vascular markings in the lung fields. Vital capacity was diminished, but there was no evidence of pulmonary oedema. In spite of the rise of venous pressure many subjects had no increase in heart rate. Of 12 subjects 4 showed electrocardiographic changes indicating slight right heart stress. Symptoms were absent or unimportant. There was a rapid fall of venous pressure to normal on ceasing injection, except in one subject given blood. There is evidence that the peripheral and pulmonary capillaries and veins dilate to accommodate the increased blood volume.

When the blood volume was first reduced by a large venesection, saline or serum injected in similar amounts and at similar rates caused little or no rise of venous pressure.

We thank Dr. Janet Vaughan for supplies of serum, and Drs. Duncan White and E. J. E. Topham for their help with the radiological measurements. The work was assisted by an expenses grant from the Medical Research Council.

REFERENCES

- Altschule, M. D., and Gilligan, D. R. (1938). *J. clin. Invest.*, **17**, 401.
 Bainbridge, F. A. (1915). *J. Physiol.*, **50**, 65.
 Bayliss, W. M., and Starling, E. H. (1894). *Ibid.*, **16**, 159.
 Caughey, J. L. (1935). *Proc. Soc. exp. Biol.*, N.Y., **32**, 973.
 Eyster, J. A. E., and Middleton, W. S. (1924). *Amer. J. Physiol.*, **68**, 581.
 Glaser, E. M., and McMichael, J. (1940). *Lancet*, **2**, 230.
 Hill, D. K., and Pincock, A. C. (1941). *Ibid.*, **2**, 754.
 Lewis, T. (1937). *Diseases of the Heart*, p. 8, Macmillan, London.
 McMichael, J., and McGibbon, J. P. (1939). *Clin. Sci.*, **4**, 175.
 Meek, W. J., and Eyster, J. A. E. (1922). *Amer. J. Physiol.*, **61**, 186.
 Sharpey-Schafer, E. P., and Wallace, J. (1942). *Lancet*, **1**, 699.
 Wallace, J., and Sharpey-Schafer, E. P. (1941). *Ibid.*, **2**, 393.
 Weiss, S., Wilkins, R. W., and Haynes, F. W. (1937). *J. clin. Invest.*, **16**, 73.
 Wiggers, C. J. (1921). *Arch. intern. Med.*, **27**, 475.
 Wood, P. (1940). *Brit. Heart J.*, **2**, 132.
 — (1941). *Ibid.*, **3**, 21.
 — and Selzer, A. (1939). *Ibid.*, **1**, 29.

SULPHONAMIDE SUNLIGHT ERUPTIONS

A CLINICAL STUDY OF 27 CASES

BY

R. G. PARK, M.B., M.R.C.P.

Captain, N.Z. Medical Corps

AND

W. M. PLATTS, B.Sc., M.B., Ch.B.

Captain, N.Z. Medical Corps

(From a Military Hospital in the Middle East)

The constant sunshine of the Middle East and the presence of considerable numbers of V.D. patients undergoing ambulant chemotherapy in an open-air tented hospital have recently given us unusual opportunities of observing cases of the light-sensitive type of sulphonamide dermatitis.

This form of dermatitis has been classified into two varieties (Tedder, 1939): generalized eruptions showing no predilection for the exposed parts, and light-sensitive eruptions. It is with the latter that this report is concerned.

Cases of sulphonamide photosensitivity were first recorded in 1937 (Newman and Sharlit, 1937; Menville and Archinard, 1937), rashes being described which appeared after from 3 to 5 days of drug therapy and which disappeared after 5 or 6 days of protection from light. Severe dermatitis has also occurred from artificial ultra-violet irradiation during sulphapyridine administration (Hallam, 1939). On the analogy of the sunlight eruptions in haematoporphyria, investigation has shown that sulphonamides cause an excessive urinary excretion of porphyrins, of which one, coproporphyrin I, is a recognized photosensitizer (Rimington and Hemmings, 1938). This occurs more often with sulphanilamide than with sulphapyridine (Wien, 1938). The porphyria varies considerably with the individual, and by exposing unselected patients on sulphanilamide Eidinow (1939) has shown that by no means everyone becomes light-sensitive.

Case Reports

During the 12 months Oct., 1940, to Sept., 1941, 486 cases received sulphanilamide, of which 21 developed eruptions (4.3%); 309 cases received sulphapyridine, yielding 6 eruptions (1.9%). Of the 27 cases 8 occurred during July and 6 during August. As measured by the photo-electric cell, the intensity of visible light in this part of the Middle East in July is about four times that in January. We have divided our cases into two groups:

1. Eruptions Limited to Parts Exposed to Sun during Chemotherapy

There were 17 cases of this kind, 16 with sulphanilamide and 1 with sulphapyridine. In 5 cases the rash appeared on the eighth day of chemotherapy, in 5 on the ninth, and in 7 on the tenth. The eruption consisted of maculo-papules in a morbilliform or roseoliform configuration. It was confluent on the most exposed parts, being distinguished from sunburn only by the discrete lesions at the margins. The demarcation

was very sharp at places such as the singlet area or, where no shirt had been worn, the waist. The axillae were usually spared. The bronzed parts—face, neck, forearms, hands—suffered less than the upper trunk and upper arms, which had been exposed less in the past. Two patients admitted to unusually heavy sun exposure the day before the rash began. Only 5 of the 17 had always sunburned easily.

In addition to the above lesions there was in one case vesiculation and pustulation of the face, in 2 oedema of the hands, and in another oedema of the eyelids. Itching varied from severe to none at all. In one case—an eruption limited to the hands—an experimental exposure of the forearms to the sun was made for 3 hours on the third day of the rash. After an hour a maculo-papular eruption appeared in the exposed area, maximal on the non-bronzed flexor surfaces. In another case, in addition to a roseoliform eruption, the dorsum of the hands showed typical target patches of erythema iris at the sites of some old scars, and the face developed oedema and pustules, going on to extensive desquamation.

Constitutional disturbance was common, fever up to 102° and malaise occurring in half the cases and three rigors in one case. The eruptions faded rapidly—in 2 to 4 days in all cases except one, which lasted a week. Three patients continued chemotherapy for from 1 to 2 days after the rash appeared, with no ill effects, though they kept out of the sun. In one case the rash began 2 days after the drug had been stopped, and in the remainder cessation of the drug either coincided with or occurred on account of the rash.

Of these 17 cases 3 received further chemotherapy at later dates, varying from 1 week to 4 months afterwards. In no instance did the eruption recur, although there was no change in the amount of sun exposure.

2. Eruptions Maximal on Exposed Parts, but also Affecting Others

Case 1 received sulphanilamide for non-specific urethritis, 4 g. daily for 8 days, during which the patient went in the sun with only the face and hands exposed. He had always sunburned easily. On the eighth day an irritable morbilliform rash appeared on the arms, face, and trunk above the waist, principally on the upper part of the back. There was mild cyanosis but no fever.

Case 2 received sulphanilamide for chancroid, 4 g. daily, until it was stopped on the eighth day because of a morbilliform and roseoliform rash, resembling the cases in group 1 except that patches appeared on both thighs over the trochanters. It was slightly itchy, and less pronounced on the bronzed parts. The patient had not previously been sensitive to sunlight. The rash lasted 4 days and was accompanied by a fever up to 103°.

Case 3 received sulphanilamide for chancroid, 4 g. daily for 10 days, during which he went in the sun wearing an open-necked shirt, elbow-length sleeves, long trousers, and rubber shoes. He has always sunburned easily. On the ninth day he developed a slightly itchy morbilliform rash, confluent on the necklace area, back, and outer upper arms. It was also present on the bronzed parts to a lesser degree. Except for a small patch over each great trochanter no rash appeared below the waist. There was no fever or malaise. The rash lasted 3 days.

Case 4 received sulphapyridine for gonorrhoea, 26 g. in 7 days. He had always sunburned easily. On the fifth day he took an open-air shower-bath in the sun, but was otherwise in bed until the seventh day, after which he went out in an open-necked shirt, long trousers, and shoes without socks. On the tenth day a morbilliform and roseoliform rash began. It was confluent on the face, forearms, and dorsum of the feet; discrete on the knees and trunk above the waist; and scanty on the trunk below the waist, upper arms, thighs, and legs. The bronzed parts suffered equally with the others. Itching was slight. The patient was afebrile, but complained of backache. The rash faded in 5 days.

Case 5 received sulphapyridine for gonorrhoea, 26 g. in 7 days. There had been no previous sensitivity to the sun. On the ninth day a dusky-red morbilliform eruption appeared, maximal on the shoulders and outer arms, less marked elsewhere, and especially scanty on the bathing-drawers area and in the axillae. The face was unaffected, but the other bronzed parts suffered equally with the remainder. There had been exposure during chemotherapy. The rash faded in 3 days. A low fever, malaise, vomiting, and headache occurred for 2 days with the rash.

Case 6 received sulphapyridine for gonorrhoea, 26 g. in 7 days. Sun exposure during this time was slight. He had never been sensitive to the sun. On the eighth day a morbilliform