War Medicine Series

The paper printed below is based on a lecture which was one of a series recently given at the British Postgraduate Medical School, Hammersmith

MALARIA IN WARTIME*

BY

Sir P. H. MANSON-BAHR, M.D., F.R.C.P.

The Quartan Parasite

The quartan is a parasite of children rather than of adults. It is a very local infection. We found a certain amount in Salonika and in Palestine in the last war, but among the troops it amounted to only 1% of the total. It is truly the most beautiful of all malaria parasites. The corpuscle in which it lives does not enlarge; its cycle of development is seventy-two hours, in contradistinction to the benign tertian, and therefore it typically produces fever every four days. If we can believe Sydenham, quartan malaria must have been rife in Westminster in his day, and most of the Members of Parliament were often precluded from their duties by recurrences of this fever. He realized too that it was associated with marshy districts such as prevailed in Westminster at that time.

Another point about the quartan is its ability to split the red corpuscle in twain. There are only eight spores formed, instead of the eighteen to twenty-four in the case of the benign tertian and eighteen to thirty-two in the case of the subtertian. The pigment is rather blacker. Little spots develop inside the corpuscular protoplasm, but they are not Schüffner's dots. In this country they are known as "James's dots," which are little blue specks, not red like Schüffner's. In this form of malaria we do not always see the classical quartan temperature chart, and this is due to the fact that there may be two generations of parasites in the blood, producing a rough kind of quotidian fever.

Plasmodium ovale

The latest arrival in the malaria family is *Plasmodium ovale*, so called because the corpuscle in which it lives tends to become oval in shape. It was discovered in 1922, and was at first looked upon as a freak; but now we know it is a common parasite in West and Central Africa, and it had been confused with the benign tertian because of its tendency to produce Schüffner's dots. The corpuscle becomes ragged, frayed, and oval in shape, presently losing its rounded contour. The parasite has only eight spores, and produces a harmless infection. It is said to have been observed in Central Asia and in Central America, but the general idea is that it is peculiar to Central Africa. *Plasmodium ovale* is now much used in malaria therapy, and is the "star turn" at present at the malaria centre at Horton, Epsom, because it produces efficient rigors when artificially inoculated and does not require any prolonged or drastic treatment, but just fades away directly small doses of quinine are administered. I have seen only one patient who was really seriously ill with this variety.

Drugs used in Treatment of Malaria

With regard to drugs in the treatment of malaria, before the last war we were told—from the teaching of Prof. Dixon in Cambridge—that the most specific of all drugs was quinine. Yet as this war progressed we began to disbelieve more and more in its specificity, and had to modify our fundamental ideas. We believe now that quinine is an extraordinarily valuable drug but is definitely selective upon certain stages of the malaria parasite. The species of the parasite on which it acts most quickly is the benign tertian, but we cannot entirely banish it from the human body by one or two or even three doses of quinine. The plasmodium has a way of side-tracking and slipping away into the interstices of the bone marrow and

the spleen, ready to emerge again at the next opportunity. This selective action of drugs was brought out strikingly when plasmoquine (pamaquin) was discovered in 1925; but it is not, as you may have thought from reading the literature, an attempt by Germans to synthesize quinine, but is a by-product of methylene-blue, which was employed by Manson in China as a febrifuge many years ago. He formulated the idea—a very clever one—that if there was a selective action of a stain for a particular blood parasite it could be surmised that the same substance might also exert a therapeutic effect, and compounded a combination with borax (Manson's stain), which he proved to be selective for malaria parasites.

When the blood is full of subtertian crescents and plasmoquine is given in sufficient doses this stage of the parasite, which is not amenable to quinine, rapidly disintegrates. The parasites burst at various points; the chromatin degenerates, and finally they vanish from the blood stream.

Atebrin (or mepacrine), an acridine compound, is another dye derived from methylene-blue. It does not affect this crescent stage at all and has no effect upon exflagellation, but causes the pigment to collect in the centre of the parasite.

Diagnosis

It is the differential diagnosis of malaria which is so important. Malaria usually causes enlargement of the spleen, but whether you can tell that this is so or not depends on your skill in palpation. In subtertian malaria the spleen becomes rounded off rather like a tennis-ball, but it is only after repeated malarial attacks that the classical hard edge can be detected. It is therefore not safe to say that a man has not got malaria because the spleen cannot be palpated. There are various ways of performing palpation, and new-comers have to learn the correct technique; and the question arises: When is a spleen not a spleen? The answer is: When it is a kidney! An enlarged left kidney is sometimes mistaken for an enlarged spleen. I recall a major in the Indian Army who drew a pension for eighteen years on the ground of enlarged spleen when in fact what he had was a large left polycystic kidney; and I have even seen this misdiagnosis made when assisted by uroselectan and radiography. There is also splenic pain, which may be significant.

In the diagnosis of malaria the great point to remember is that in a malarious country it may simulate almost any other disease. Subtertian malaria is the greatest mimic. Quite commonly it produces cerebral symptoms—cerebral malaria, due to the congregation of parasites in the brain, sometimes leading to coma and death. I remember how some soldiers marching from Jaffa to Jerusalem in 1918 fell out by the side of the road waiting for the ambulance, and before it could reach them they were dead. Very often in the Near East the worst forms of malaria are found in the winter-time. Mosquitoes once infected remain so, and in winter tend to congregate in buildings and nooks where there is warmth and then descend upon anyone they can find. These dire cerebral forms may crop up during the winter months and may be mistaken for other forms of cerebral disease. In the summer the patient may go down with hyperpyrexia, and so the case may be mistaken for heat-stroke.

I would especially warn against confusing alcoholism with malaria. In a malarious country a man may be accused of being drunk when he is suffering from malaria.

Mania is another thing which may be misleading: the patient may be suffering from subtertian malaria delirium; and, on seeing a sudden epileptic fit, do not jump to the conclusion that it is true epilepsy, for it may be malaria simulating that condition. Then there are attacks simulating cerebrospinal meningitis with pain and stiffness at the back of the neck. There is an appendicular form of malaria that is due to sporulating forms of the parasite which select that particular part of the intestinal canal in which to congregate. When the patient's temperature rises with rigor to 104° F. as a sequel to an abdominal operation, you will want to ask whether malaria parasites were not in the blood before the patient was subjected to operation or whether the operation actually incited the rigor. Cholecystitis is also a dangerous primary diagnosis. Why should an epidemic of cholecystitis in healthy young men suddenly arise? The explanation lies in

* Concluded from p. 462.

the hepatic congestion in subtertian malaria, with an acutely distended gall-bladder.

Then there is malarial dysentery, or colitis. Malaria is frequently accompanied by diarrhoea; in fact, often there may be diarrhoea and nothing else to guide one. It is very common to see patients from West Africa who have suffered from diarrhoea on the voyage and, owing to this, doctors have prohibited them from taking quinine. I have seen such cases develop cerebral malaria and fatal coma the moment the boat has docked.

In Egypt subtertian parasites have been demonstrated in scrapings made from the bowel. "Malarial dysentery" with blood and mucus in the stools simulates acute bacillary dysentery. It is possible to get such an acute diarrhoea with malaria that it resembles cholera. There were several cholera scares in the last war, but they were due to a choleraic form of subtertian malaria. Then we may meet the icteric forms; but malarial icterus is not a really true obstructive jaundice. In subtertian malaria it is almost diagnostic to get urobilinuria with sherry-coloured urine, and this may sometimes be taken as an index of infection, even in the absence of subtertian parasites in the blood.

I remember in 1916 being asked to see a soldier who had come down from Khartum intensely jaundiced. He was evidently dying. I was then in charge of the laboratory, and found that I had examined his urine and serum, and carried out agglutination tests; but the one specimen that had not been examined was the blood film, which was crammed with subtertian parasites. By the time this was done, however, it was too late! One can meet with forms resembling acute pancreatitis in which the pancreas is affected; others resemble intestinal obstruction, with profuse vomiting. One may even see cardiac forms with generalized oedema, which were frequently encountered in the last war, or, again, pulmonary forms associated with bronchitis or pleurisy. Further, there may be some resembling acute nephritis with blood and cells in the urine; or a cutaneous rash with multiple haemorrhages in the skin, somewhat resembling typhus or measles.

Blackwater Fever

Finally, there is the most dramatic sequel—blackwater fever, solely associated with subtertian malaria. It probably represents an anaphylactic phenomenon due to hypersensitivity to toxins of this parasite. Blackwater fever occurs only in those parts, such as the Jordan Valley, Macedonia, and West Africa, where the subtertian parasite is prevalent. The prevention of blackwater fever is closely linked up with the prevention of malaria, so that if a man does not get infected with the subtertian parasite he will not develop blackwater. If a laboratory service is maintained by which in all cases the blood is examined for malaria the moment the patients fall ill, then the onset of blackwater fever will be prevented. In the years immediately succeeding the last war I never treated less than 4 or 5 cases of blackwater fever here in London every winter; they were usually acute dramatic cases, and the death rate figured between 25% and 50%; but since the rule has been established that all must have their blood examined directly they land in this country from West Africa, blackwater has not been observed. A patient is exposed to the danger of blackwater fever once he has had acute malaria—usually up to nine months. It is apparently a case of supersaturation with this parasite, so that blackwater seldom results after a primary infection. It occurs mostly in the old residents of the Tropics who, on returning to this country, have been exposed to cold and wet or some shock which causes a sudden multiplication of parasites. Then the red blood corpuscles break down and become haemolysed, and the victim may die from blocking of the kidneys with products of their disintegration. If, therefore, this process can be prevented from taking place, then blackwater fever will not ensue. Treatment depends upon the alkalization of the blood and upon prompt blood transfusions.

The Importance of Blood Examination

The diagnosis of malaria depends upon correct blood examination. The diagnostician must be familiar with all stages of the malaria parasites: he should be so sure of his premisses

that he can recognize them as readily as Smith, Brown, or Robinson. But can ordinary unsophisticated folks learn to distinguish these appearances in a reasonably short time? The answer is that they can be taught to do so. In Central Africa there are native dressers who are proficient in spotting parasites. In the last war when put to the test we had to improvise an organization for the detection of malaria at the earliest possible moment for treatment in the field. There originated the diagnosis stations which have now been included in the Army service. Orderlies were chosen from the ranks, some of them with little previous education, and by six weeks' intensive training, by making them look at and stain blood films, we got them sufficiently accurate in discerning malarial parasites to be of great service. A number of these men are still employed here in London at the present time. Those thirty men who were trained were responsible for diagnosing 417,000 blood slides in the last year of the war in Palestine.

Prevention and Treatment

How are we going to prevent malaria in the field? Quinine in large doses is a poison; it may produce amblyopia in some people, and they may become blind. Quinine in enormous doses, as a means of preventing a malarial attack, is a broken reed. It has been shown that one can inject quinine together with malaria parasites into the blood of the recipient and yet produce malaria. On the West Coast of Africa most officials cling to the belief that 5 grains of quinine will prevent malaria and blackwater. It probably modifies the acuteness of the malarial attacks and thereby the proclivity to blackwater fever, but when you give soldiers who are operating in open country 5 grains of quinine a day it has little visible effect in reducing the incidence of malaria. For prevention on a large scale the chief line of defence is the mosquito net. The next best method is to kill the mosquitoes.

With regard to drug treatment of the actual attack, however, quinine still remains the standard. It is most effective in benign tertian malaria and it has also a great effect upon subtertian. But it is necessary to remember that quinine given in big doses to a patient with a large number of subtertian malaria parasites is likely to precipitate blackwater fever by the sudden dissolution of the parasites, and the patient may die within twenty-four hours. For benign tertian malaria, quinine should be given in doses of 10 grains three times a day, and the bowels should be freely kept open. One must go on giving the quinine in decreasing doses for about six weeks, though even this continuous therapy will not extirpate the infection from the body entirely; but to inject quinine intramuscularly to everybody indiscriminately is to ask for trouble, as abscesses and indurations are apt to form in a certain number. At the same time I never hesitate to give quinine intramuscularly when it is necessary as a remedial measure in severe infections. The results are magical if it is given in the right way and injected deeply into the gluteal muscles, but it should never be injected below the folds of the buttocks.

You cannot treat malaria scientifically unless you control treatment by frequent blood examinations and can recognize the changes in the parasites when acted on by quinine or other drugs. Intravenous quinine is reserved for the most severe forms of subtertian malaria. From 5 to 7 grains dissolved in 5 c.c.m. of distilled water is quite enough. With regard to plasmoquine, remember also that this is poisonous. When it was first introduced we gave it in too big doses, and were thereby producing poisonous effects. It is now recognized that small amounts of plasmoquine in tablet form—known as plasmoquine compound—are beneficial, each tablet containing 1/6 grain, with 2 or 3 grains of quinine. This, given as a regular course, has been found by the Army in India to be the best method of treating benign tertian malaria and of preventing further relapses. In such cases a start is made with the quinine and continued with plasmoquine compound, the dose of which is two tablets twice a day for a week, amounting to four courses of seven days, with a break of four days between each. This means that about 2/3 grain of plasmoquine is given a day, in addition to 20 grains of quinine. With this method the relapse rate in benign tertian malaria has fallen from 40% to something like 8% in the Army in India.

Plasmoquine and atebirin go badly together, but atebirin and quinine work well in combination. The dose of atebirin is 1/10 gramme three times daily for seven days. It is a yellow tablet which is given by the mouth; but there is also a soluble form which can be injected. It is active especially in the treatment of subtertian malaria, and I have seen only one case in which blackwater fever supervened after atebirin, so it is useful in this direction. I do not believe, however, that it is nearly as effective in treating benign tertian malaria as the method I have outlined. Certain disadvantages of atebirin have been pointed out, and it is well to warn people that they may acquire a somewhat yellowish tinge. I have never seen anybody go "off his head" when taking atebirin, but acute mania in native races has been described after taking it in therapeutic doses, though this certainly does not seem to be common in Europeans.

RECEPTION OF CASUALTIES IN AN E.M.S. HOSPITAL

BY

STANLEY J. FIRTH, M.D., M.R.C.S., D.P.H.

Medical Superintendent

AND

HAROLD PARK, M.B., F.R.C.S., F.R.C.S.Ed.

*Resident Surgeon**The Municipal Hospital, Brighton*

The sudden influx of a number of wartime casualties, whether Service cases or civilian, calls for a procedure different from that used for ordinary peacetime admissions. These casualties are the immediate concern of several separate hospital departments, and it is therefore important that each section should be supplied with the necessary data in the simplest form. With this in view we prepared an urgency admission sheet which has worked very well in actual practice and which, having been thought deserving of praise in high medical quarters, we consider worthy of publication.

This Sheet to be issued to the Ward and used on admission of Casualty.

Name of Patient		Age	
Date admitted	Time	a.m./p.m.	Religion
Clerical Details recorded:			
<div style="display: flex; justify-content: space-around;"> </div>			
CONDITION ON ADMISSION	T.	P.	R.
			R.P. 1.
			2.
			3.
PROVISIONAL DIAGNOSIS.		Dangerously ill Seriously ill Slightly ill	
IMMEDIATE TREATMENT.		Morphia Radiant Heat Intravenous therapy - Blood Plasma Saline A.T.S. Oxygen Catheter.	
X-RAY			
OPERATIVE TREATMENT.		PRIORITY (red label) Required EARLY (yellow label) DELAYED (green label) Not required	
PREMEDICATION & ANAESTHETIC.			
Examining M.O.'s initials			

This sheet, the front page of which is here reproduced, is hung at the foot of the bed in a temperature-chart holder, and consists of a single piece of paper about the size of the printed page of the *Journal*, printed on both sides. A description of the back of the sheet can be given in a few words. It is divided into two sections, one half being for the operative

findings, the other for the post-operative instructions. This side is completed by the anaesthetist during the operation.

The front of the sheet is a record made up of the following sections. It is headed by the name, the Service No. of Service cases, age, date and time of admission, and religion. Immediately below this is a space for the use of the clerical staff, who, having recorded the details which concern them, denote this by affixing their stamp. Next comes a section in which are entered the temperature, pulse, respiration, and blood pressure of the patient, together with a pictorial record of the injuries, which are roughly sketched on a series of anterior, posterior, and right and left lateral diagrams of the body. The section that follows provides a space for the provisional diagnosis, together with three headings denoting the severity of the patient's condition. The immediate treatment occupies a section below this, and consists of a list of modern resuscitative procedures which the surgeon orders by eliminating those not required. Radiographic instructions are provided for, and the succeeding section indicates whether operation is required or not and gives the preferential order for cases to be taken to the theatre. A coloured label is affixed according to whether this preference is (a) priority (red), (b) early (yellow), or (c) delayed (green). This classification has proved to be a great asset in dealing with a ward full of casualties. It is a guide to the radiographer as to the cases to take first and as to the order of their transfer to the theatre. Thus, those with the red label are first to go, being followed by the yellow label cases; the green label cases can usually be left till next morning for operative treatment, the immediate dressings being done in the ward.

Value of a Simple Sheet in Hospital Practice

The advantages of this type of sheet to the various hospital departments can best be shown by a brief statement of how the system works in our own hospital.

Medical.—We have found that a surgical team can, on an average, deal with 25 casualties. The idea of rushing new arrivals to the operating theatre has been generally discarded since the days of Dunkirk, it being realized that methods of resuscitation and procedures to counteract shock are of paramount importance. The surgeon in charge of a group of casualties has therefore adequate time to go the round of his cases, provided the treatment can be clearly indicated with the least possible delay. For this reason copious note-taking at this time must be dispensed with. As will be seen, the subdivisions of the sheet allow fully for this to be done. The surgeon makes his examination and in a few seconds completes the sheet, thereby giving the provisional diagnosis, instruction for radiography, the resuscitation procedures he requires, and the grading of cases for time of operation. He also assesses the degree of severity, which indicates to the clerical staff the procedure which instructions state shall be followed for E.M.S. cases.

The other members of the team follow up and, taking the red label cases first, carry out the ordered treatment which requires their personal attention—for example, intravenous therapy, blood pressure, and haemoglobin estimations. The treatment having been completed the sheet is initialled in the space provided at the side. Thus the surgeon is enabled to look though his own cases and then proceed with the operative work, each patient arriving at the theatre in the correct order and with the preliminary treatment carried out.

Radiography.—It has been found that practically all cases admitted under war conditions require an x-ray examination. The radiographer, working with a portable apparatus, takes the cases in order according to the coloured label. He has not to keep asking which to take first; he can proceed unhindered with his work, his instructions having been given in the appropriate column. When the films are completed they are hung at the foot of the bed and go with the patient to the theatre.

Nursing Staff.—Injections ordered are given by the sister, who, with the help of the concise recordings of the sheet, gains a clear and quick conception of what is happening to patients in her ward. Anaesthetic premedication is given, and there is little likelihood of cases being sent to the theatre in the wrong order, thereby delaying treatment of more urgent