Rh-positive and the mother's serum contained an Rh antibody. Moreover, in one of these cases the mother had previously given birth to an infant affected with icterus gravis neonatorum. This finding suggests that certain cases usually diagnosed as physiological jaundice would be more properly regarded as mild cases of erythroblastosis. The supposition that the aetiology of some cases of mild jaundice of the newborn is connected with the passage of incompatible agglutinins across the placenta receives further support from the findings in the other mothers and infants investigated. In these the incidence of cases in which the mother's serum was incompatible with the infant's erythrocytes (due to the presence of anti-A or anti-B agglutinins) was considerably higher than the expected incidence in a random sample of mothers and infants.

(The concluding part, with a list of references, will appear in next week's issue)

TREATMENT OF SHOCK BY DIRECT ACTION ON THE VEGETATIVE NERVOUS CENTRES*

BY

LENA C. STERN, M.D.

Professor of Physiology, Second Moscow State Medical Institute

The Mechanism of Traumatic Shock

Efforts to produce shock experimentally play an important part in the elucidation of the mechanism of traumatic shock. The criteria of shock conditions in all these experiments were: (1) sudden and persistent fall in blood pressure, with a rapid and feeble pulse, (2) shallow breathing, (3) the absence of reaction on irritation, (4) lowering of the temperature. All this points to a disturbance of the vegetative nervous system, particularly the sympathetic, which in itself leads to a diminution of volume of the circulating blood, due to the accumulation of blood in the blood depots and diminution in the liquid part of the blood from plasmorrhoea. This disturbance of the vegetative nervous system is also a cause of chemical changes which characterize the course of shock (appearance of acidosis, diminution of glucose, etc.). This work also showed that with the weakening of the sympathetic nervous system there is a stimulation of the parasympathetic system which shows itself by a number of inhibitive effects.

Experiments made by us and our colleagues (Rosin, Chvoles) have proved that sensitivity, especially to pain, plays an important part in the development of shock. They showed that irritation of the skin by an induction current for 2 to 6 hours causes shock, whereas with preliminary anaesthetizing of the skin with novocain, much longer electrical stimulation is without effect. Given the same conditions, shock is most easily produced by irritation of the skin on the inner surface of the thigh, the peritoneum, and the periosteum, whereas irritation of the dura even for as long as 8 to 12 hours does not cause shock. The rhythm of irritation is important, shock usually appearing much more quickly when the rhythm is quicker.

The significance of the preceding conditions for the development of shock in man has been proved by clinical observations. Excessive tiredness, prolonged insomnia, insufficient nourishment, fear, exposure, loss of blood, and, not least, painful irritations, lead, after transient overstimulation of the sympathetic nervous system, to its exhaustion and thus facilitate shock. The results of these experiments show that shock is caused, apart from loss of blood, by a whole complex of circumstances connected with a definite condition of the nervous centres. Only when all these conditions are present does shock develop. In clinical studies of shock, attention has been mainly directed to the functional state of the peripheral organs, such as the cardiovascular and respiratory organs. A more detailed analysis, however, has shown that all these

cardinal changes, including the changes in metabolism and in the physical and chemical state of the blood and the tissues, are not actually specific to shock. In fact, similar changes, such as a fall in blood pressure, diminished nervous reactions, etc., can be caused by a depression of the sympathetic nervous system—such as occurs also in other pathological states. All these changes form a syndrome which characterizes shock and shock-like conditions and can be caused by action on the vegetative centres; this points to the important part played by the central nervous system in the production of shock. It therefore follows that in order to be able to understand the mechanism of shock and to apply useful therapy and prophylaxis one must concentrate on the functional conditions of the nervous centres, on their metabolism, and, in conjunction with this, on the character of the cerebrospinal fluid—i.e., their immediate nourishing medium.

Function of Cerebrospinal Fluid

The importance of the cerebrospinal fluid to the functioning of the nervous centres has been shown by observations in laboratories and clinics for many years. These observations have proved that the cerebrospinal fluid (taken in its broad sense as the fluid which fills not only the ventricles of the brain and the subarachnoid space, but the whole of the perivascular and pericellular spaces of the brain tissue) is the immediate nourishing medium of the central nervous system. The chemical composition, as well as the physical, chemical, and biological character, of the cerebrospinal fluid mainly depends on, and is determined by, a special mechanism—the so-called haemato-encephalic barrier. The anatomical substratum of this barrier is, apart from the vascular plexus, the capillaries and precapillaries of the brain, in the first place their endothelium. The haemato-encephalic barrier has a great selective capacity, on account of which not all substances which normally circulate in the blood, or get there accidentally, pass from the blood stream into the c.s.f. There takes place "a certain choice" as the result of which a relative constancy of the composition of the c.s.f. is maintained, and there is a certain independence of possible changes in the consistency and quality of the blood.

It has been shown that not only the introduction of abnormal substances from the blood into the c.s.f., but even a simple disturbance of interrelations between the normally existing substances in the c.s.f., such as between certain electrolytes, causes a more or less severe reaction in the nervous centres which sometimes leads to the death of the animal. In certain cases we were able to notice that definite changes in the c.s.f. are accompanied by some features which resemble the complex of symptoms characteristic of shock and similar states. This particularly applies to the ions of Ca and K. It was shown that change in the concentration of these ions in the c.s.f., leading to a change in the coefficient K/Ca , has a very strong influence on the condition of the central nervous system, particularly on the vegetative centres. For example, the introduction of minimum doses of Ca salts directly into the c.s.f., especially into the ventricles, brings about in the animal a condition of severe depression, whereas the introduction of minimum doses of K salts produces pronounced excitement. This suggests that the direct cause of the change in the functional condition of the cerebrospinal system may be a corresponding change in the composition of the c.s.f. Comparing the occurrence in shock of changes in the different physiological systems, particularly in the cardiovascular and respiratory systems, with the action which causes a stimulation of the vegetative nervous system, we conclude that the sympathetic and parasympathetic nervous centres play the most important part in the production of shock. The cause of changes in the vegetative nervous centres may be strong irritation coming from the periphery. The intensity of effect brought about by such irritations depends on the preceding condition of the vegetative centres (their greater or lesser excitability and reactivity), which in itself is closely connected with the chemical composition of the c.s.f.

The possibility of varying the reaction of the sympathetic and parasympathetic centres by introducing into the cerebrospinal canal certain substances such as K salts and Ca salts points to the influence of the electrolytic composition of the c.s.f. on the condition and activity of the vegetative nervous

* Translated by Dr. H. W. Swann.

system. Diminished tone of the sympathetic centres causes those disturbances of function characteristic of shock, and especially of the haemodynamics, the disturbance of which is quite sufficient to bring about in different physiological systems those changes which characterize shock. It is therefore understandable that all efforts to combat the signs of shock are mostly directed to the re-establishment of circulation and the normal action of the cardiovascular system.

Failure of Shock Therapy

However, the ordinary methods of fighting shock generally do not give positive results, because they do not influence the condition of the central nervous system. These negative results are explained to a great extent by the presence of the haemato-encephalic barrier.

As regards certain medicinal substances, it has been noted that, introduced into the circulation, they have no effect on the central nervous system, whereas the introduction of minimal quantities of the same substances directly into the cerebrospinal canal, particularly into the ventricles, often has a marked effect. The conclusion, therefore, is that in order to obtain a positive effect one should introduce the substance intrathecally. The action of a substance admitted into the general circulation differs from the action of the same substance introduced straight into the c.s.f. This applies particularly to those substances which are normally present in the c.s.f. and the blood, such as K and Ca salts and some hormones. Many experiments have proved that there is an antagonism between the central and peripheral parts of the vegetative nervous system as regards their reaction to the same substance. This antagonism more or less explains the absence of a positive anti-shock action of the medicinal substance when introduced straight into the circulation. Actually, if these substances do penetrate into the central nervous system, particularly the vegetative centres, they may produce an opposite effect and thereby neutralize the immediate action which they have produced on the peripheral vegetative organs. It follows, therefore, that in order to understand the action of a given substance one must take into consideration not only the permeability of the haemato-encephalic barrier for this particular substance, but also the fact that the substance which easily passes through the haemato-encephalic barrier into the c.s.f. may, on contact with the vegetative centres, cause an action which is opposite to the one on the peripheral organs. There is, therefore, an interference, and the final effect may be negligible or even opposite to the one expected. To produce a direct action on the vegetative nervous centres one must administer the substance directly into the c.s.f.

One must add that a positive effect can be obtained only in cases in which the given substance is introduced into the ventricles of the brain, whereas the introduction of the substance into the subdural or subarachnoid spaces is usually without effect because it passes rapidly from the c.s.f. into the blood stream and does not come into contact with the nervous centres.

Intraventricular Injection of K Salts

Laboratory experiments (Chvoles) have shown that in many cases of experimental traumatic shock the direct injection into the c.s.f. of a solution of K salts gave a positive result. In most of the experiments the signs of shock disappeared very quickly, the blood pressure rose rapidly to normal, the rhythm of respiration became normal, as also the reactivity and excitability of the animal.

It is interesting to note that in those cases in which trauma was accompanied by a severe loss of blood (up to 50%) the introduction of potassium phosphate into the c.s.f. without a preliminary blood transfusion brought the animal out of shock, whereas in control experiments the introduction of a physiological solution as well as a transfusion of blood had no effect. It appears, therefore, that under the influence of increased excitability of the sympathetic centres there is a mobilization of the blood from the depots and simultaneously a transference of fluid from the tissues into the blood, as a result of which the volume of the circulating blood is increased.

The best results have been obtained by injecting potassium phosphate directly into the lateral ventricles or into the cisterna magna; provided that definite rules are followed. If these rules

are not observed, the injected fluid often does not reach the cerebral ventricles, but, following the normal flow of the c.s.f., passes into the subarachnoid space without reaching the vegetative centres. The introduction of the fluid into the subarachnoid space does not produce the necessary effect, because it passes into the blood stream very quickly without coming into contact with the nervous centres. It can now be positively stated that by fulfilling the necessary conditions as regards concentration, temperature, and reaction of the solution of potassium phosphate, as well as the rules of technique of its introduction into the cerebrospinal canal (i.e., the ventricles), a definite effect can as a rule be obtained. It is obvious that a positive action of potassium phosphate on the vegetative nervous centres is possible only if the centres have still retained the ability to react. Completely exhausted vegetative centres which have lost their reactivity cannot be re-established by this method.

Taking into consideration the fact that the introduction of potassium phosphate into the cerebrospinal fluid increases the tone of the sympathetic centres, its use is indicated in those cases in which symptoms of weakened tone are present. Irrespective of whether we deal with shock or a shock-like condition, we cannot possibly expect any useful effect in cases in which there is an anatomical injury to the vegetative centres. An effect can be expected only in a purely functional disturbance of the vegetative nervous system.

The time factor is of great importance. One obtains the best results in cases when the potassium phosphate is introduced into the ventricles of the brain within a short time after the appearance of shock. This applies not only to traumatic shock but also to anaphylactic, peptone, and histamine shocks. This can be explained by the fact that the exhaustion of the vegetative centres gradually increases, and when it reaches a certain stage the possibility of stimulating them by any method disappears.

The results obtained by laboratory methods have been fully corroborated by clinical experience and at the front. To obtain a positive effect 1 c.cm. of a solution of potassium phosphate in a concentration of 1/6 gramme-molecule, with pH of 7.6, was found best. Introduction of potassium phosphate into the lateral ventricles not only increases the concentration of potassium but also diminishes the concentration of the Ca ions, which leads to a considerable increase in the coefficient of the K/Ca ions. When the potassium phosphate is injected, one must remember the necessity for rapid penetration into the ventricles so that direct contact may be obtained with the vegetative nervous centres. Laboratory experiments, as well as clinical observations, have shown the importance of the preceding condition of the sick or wounded, particularly the increased excitability of the nervous system in the development of shock. In prophylaxis it is rational to counteract the increased excitability so as to prevent exhaustion, which causes shock, by giving sedatives, and blood transfusion when there is loss of blood. To prevent a possible relapse this prophylaxis can be successfully applied also in the initial stages of shock, as well as after the shock has disappeared. The first condition for preventing a repetition of shock syndrome in traumatic shock is the removal of the direct cause of shock, particularly the source of painful sensations, as, for example, by local anaesthesia. Fluid can be introduced into the ventricles of the brain by way of cisternal puncture. The following rules must be observed.

Technique of Cisternal Injection

The patient must be in a horizontal position (preferably on his left side); the head must be lower than the body and bent on to the chest; the lower limbs must be flexed (a position similar to Trendelenburg's). The landmarks are the external occipital protuberance and the first cervical spinous process. These landmarks are felt on the back of the head, which has been shaved and thoroughly disinfected. One fixes with a finger the skin between these two points and inserts the needle exactly in the midline at the level of the external auditory meatus. Sliding along the occipital bone one reaches the atlantoid membrane. The moment this membrane is perforated one hears a crackling sound. For this puncture one may use a graduated cisternal needle or an ordinary lumbar puncture needle. As a preliminary measure one estimates the length which corresponds with the depth of the puncture (from 4 to 5 cm.). The criterion that the needle has penetrated into the cisterna magna is the appearance of translucent c.s.f., which escapes

from the needle after removal of the stylet. On account of weak pressure in shock conditions the c.s.f. very often does not appear spontaneously and has to be drawn out by a syringe. After 1 to 2 c.cm. of the c.s.f. have been withdrawn with a syringe, 1 c.cm. of the solution of potassium phosphate, which has been warmed to body temperature, is injected with a syringe under a definite pressure, so that it penetrates into the ventricles. This method of treatment is very easy and can be applied at the front. The preparation of the potassium phosphate solution is not complicated, but must be done very carefully. The glass of the ampoules must be neutral. The solution is a mixture of the monometallic and bimetallic potassium phosphate (concentration 1/6 gramme-molecule), and can be prepared either from these salts (K_2HPO_4 and KH_2PO_4) or by adding the orthophosphoric acid to a solution of corrosive potassium up to pH 7.6.

As a rule the effect of this injection takes place in 1 to 2 minutes or even sooner (i.e., in the first place the blood pressure rises and the respiration becomes deeper). The absence of effect within a few minutes is generally due to the fact that the fluid has not reached the ventricles, as a result of the incorrect direction of the needle or of insufficient pressure. In these cases one must repeat the operation. When toxæmia is threatening—i.e., when toxic substances which form in the injured tissues enter the blood—the further formation of these substances must be stopped by excising the injured parts where possible; positive results, which are obtained by the direct action of potassium phosphate on sympathetic nervous centres, do not exclude the possibility of the appearance of toxæmic shock later on.

Conclusions

1. In the last phase of traumatic shock there is first of all a considerable fall in the tone of the sympathetic nervous system and a partial rise in the tone of the parasympathetic nervous system.
2. The most effective way of combating traumatic shock in its later phase is to raise the tone of the sympathetic centres by direct action on the vegetative centres; potassium phosphate simultaneously raises the tone of the sympathetic centres and lowers the tone of the parasympathetic.
3. By direct action on the nervous centres the ionic potassium causes an excitability of the sympathetic nervous centres, whereas the ionic phosphates, by concentrating the ionized Ca, cause a diminution of the tone of the parasympathetic nervous centres.
4. A direct introduction of potassium phosphate into the cerebrospinal canal contributes to a quicker action of this substance on the nervous centres and at the same time obviates the possibility of its opposite effect on the peripheral part of the vegetative nervous system.
5. The use of potassium phosphate as an anti-shock remedy is indicated in all those cases in which there is a diminution of the tone of the sympathetic nervous centres, such as a fall of the blood pressure, a lowering of cardiac activity and respiration, diminution of excitability, reactivity, etc.
6. Potassium phosphate must never be used in cases in which there is increased excitability of the sympathetic nervous centres, such as sometimes occurs in the first stage of shock.
7. If haemorrhage is present it must be arrested by all available means, because the considerable rise of the blood pressure through the introduction of potassium phosphate will definitely increase haemorrhage.
8. To avoid a repetition of shock it is necessary after the introduction of potassium phosphate to remove the actual cause of shock, in particular painful stimuli.
9. To ensure the direct action on the vegetative nervous centres, which are situated in the ventricles of the brain, potassium phosphate must be introduced directly into the ventricles. This is done by means of cisternal puncture. The fluid must be introduced with a definite pressure and the direction of the needle must be correct. The position of the patient must be horizontal, preferably on the left side, and the head must be lower than the body.
10. One must act quickly, without loss of time, so as to avoid a complete exhaustion of the vegetative centres.
11. The introduction of potassium phosphate does not exclude all the other measures against shock, especially blood or plasma transfusion.
12. Blood transfusion must be done in all cases where there is marked loss of blood. In other cases a blood transfusion is not absolutely necessary.

RECENT TRENDS OF SOME INFECTIOUS DISEASES

BY

W. J. MARTIN, Ph.D.

(Of the Medical Research Council Statistical Staff)
London School of Hygiene and Tropical Medicine

The results obtained by the introduction of new methods of treatment in certain types of infectious diseases have in hospital experience been almost dramatic. It seemed worth while to find whether any change could be shown in the number of deaths for the whole country from the numerically more important causes for which control has been attempted. The estimated saving in lives following the introduction of new methods forms an interesting speculation. The death rates cannot be used as a statistical index, since the populations at risk in 1940 and 1941 are not definitely known, so the ratio of deaths to notifications has been adopted.

Cerebrospinal Fever

The employment of chemotherapy in cerebrospinal fever has resulted in a very large decline in the fatality rate from this cause. A large increase in incidence of cerebrospinal fever occurred after the outbreak of war, an experience similar to that of 1914-18. The statistics for recent years in England and Wales are:

Year	Deaths	Notifications	Deaths per 100 Notifications
1934	729	1,080	68
1935	617	865	71
1936	635	967	66
1937	698	1,113	63
Average of 1934-7	670	1,006	67
1938	652	1,258	52
1939	517	1,500	34
1940	2,584	12,767	20
1941	2,163	11,073	20

If the average of the years 1934-7 is taken as representing the endemic level immediately before the general use of sulphapyridine, it will be seen that a great improvement has been effected. The fatality rate was halved in 1939, and in 1940 and 1941 it was less than one-third of the average value. On the basis of the average fatality rate of 66.60% during 1934-7, the expected number of deaths in the three years 1939-41 would have been 16,876. The number actually registered was 5,264, or 11,612 fewer than expected. This is probably an over-estimate, since in a period when a disease is prevalent fewer mild cases are missed and there tends to be a greater proportion of wrongly diagnosed cases than in non-epidemic periods—factors that produce a decrease in the fatality rate. The most recent pre-war epidemic occurred during 1931-3, when 5,919 notifications with 3,595 deaths were reported, giving a fatality rate of 60.74%. The expected deaths in 1939-41 on the basis of this fatality rate would be 15,392—i.e., 10,128 in excess of the observed. An objection that could be raised to this comparison is that the conditions of life were vastly different in the two periods. To obtain approximately similar conditions of life—the aggregation of large sections of the population—it is necessary to go back to the last war. During the epidemic of 1915-17 there were 5,222 deaths and 8,068 notifications, yielding a fatality rate of 64.72%. Taking this rate as a standard, the expected deaths in 1939-41 would have numbered 16,400, or 11,136 more than were recorded.

It seems reasonable to suggest that as a consequence of the discovery and adoption of treatment by sulphapyridine about 10,000 lives were saved during 1939-41.

Pneumonia

The response of cases of pneumonia, particularly lobar pneumonia, to treatment with sulphapyridine has been very successful, and large declines in hospital mortality have been