War Medicine Series

The paper printed below is based on a lecture which was one of a series recently given at the British Postgraduate Medical School, Hammersmith

TETANUS

BY

LESLIE COLE, M.D., F.R.C.P.

The causal organism of tetanus, *Clostridium tetani*, is a rod-shaped Gram-positive organism. Its most important attributes are that it produces spores which are highly resistant, that these spores germinate in wounds only under anaerobic conditions, and that it produces an exotoxin which is absorbed and has a strychnine-like action on the cells of the central nervous system. It is prevalent in soil and is a common inhabitant of the digestive tract of many domestic animals, particularly horses. Perhaps for this reason the disease is most often found in agricultural districts, where the spores are spread by manure and horse ploughing.

Predisposing Causes

The incidence of tetanus in a district or on a battlefield depends partly on the prevalence of spores in the soil and partly on the type of wound. Deep, penetrating, or lacerated wounds allow spores to develop under anaerobic conditions, and growth is also favoured by pyogenic or other anaerobic infection and the presence of tissue necrosis or foreign bodies. Such conditions are a commonplace of modern war. On the other hand, uncleanness of slight or superficial wounds with scab formation, sealing of slight wounds, or infection of body cavities may also lead to tetanus, and these are now the chief sources of infection in civilian practice. Whitlows in agricultural workers are a not uncommon cause of severe tetanus. Other, though unusual, sources of infection are intramuscular injections, grey wool used in padding plaster-of-Paris splints and infecting the wound under the plaster, felt applied over a pressure sore, infected catgut, puerperal sepsis, and umbilical infection in the newly born.

With the widespread use of prophylactic antitoxin the type of wound which causes tetanus has been profoundly modified. After the obvious wound it is prevented, but it tends to crop up when least expected.

Prevention

With thorough organization it should be possible to prevent tetanus entirely, but once the disease has developed only a certain proportion of patients can be saved. Prevention is therefore of supreme importance, and two methods are now available.

Passive Immunization.—This is attained by injecting a small dose (3,000 I.U.) of antitoxin soon after a wound has been received. The value of this was proved when it was first introduced on a large scale during the early months of the war of 1914-18. The figures given by Bruce show that when this was done the incidence of tetanus fell from 9 per 1,000 wounded in September, 1914, to 1.4 per 1,000 in the November. Even when this does not prevent the disease entirely it usually lessens the severity and lowers the mortality. After prophylactic antitoxin cases generally have a long incubation period, relatively frequent local tetanus, and a slow onset of reflex spasms or sometimes none at all. It is important to remember that the passive immunity conferred by one such injection lasts only between two and three weeks, so that a wound in which infection with tetanus persists will produce the disease at the end of this time if no further prophylaxis is given. For this reason the injection should be repeated at weekly intervals after severe wounds, and an injection should always be given

before operation or manipulation of old wounds because these have been known to stir up a latent infection. After some very severe heavily infected wounds the first prophylactic dose may with advantage be doubled.

Active Immunization.—Ramon and Zoeller showed that this could be produced by injections of formalized toxin (toxoid), and a modification of their method is now used in the British Army. Two injections of 1 c.cm., at an interval of six weeks, are given to the soldier before he goes on active service, and this procedure seems to confer immunity for a year and probably much longer. It is very important to remember that the rise in antitoxin titre is much greater after the second injection than after the first, so that a single injection has comparatively little immunizing power and the second must be given. It has been shown that the immunizing effect of toxoid is increased if it is given with T.A.B., and the two may therefore with advantage be combined.

The Passive and Active Methods Compared.—The disadvantages of the passive method are that antitoxin has to be given soon after a man is wounded, which is not always possible in modern war; that, occasionally, severe reactions follow or sensitivity to horse serum, given later, is produced; and that even when repeated the antitoxin does not always completely protect. The active method, on the other hand, only rarely produces slight reactions, has the great advantage that it can be completed before the soldier goes on active service, and does not need to be repeated until a year or more. Its efficacy has not yet been tested out on a mass scale over a long period, but experimental evidence and practical experience in this war suggest that it is reliable. For reliance to be placed on this method alone in an army it is clear that every man must be immunized; therefore it is important that exact records of injections should be included in his papers. At present both methods are used, prophylactic antitoxin being given to all wounded even if they have already been actively immunized.

Description of the Disease

At onset typical tetanus is characterized by fleeting pains, chiefly in the neck and back; these are followed by the gradual onset of stiffness, first affecting the masseters so that there is difficulty in opening the mouth, and later the muscles of the trunk so that the spine is rigid and slightly arched and the abdomen board-like. In the early stages the eyebrows are drawn up, with wrinkling of the forehead, and the corners of the mouth are drawn back, producing a peculiar facial expression called the risus sardonicus. Dysphagia is sometimes an earlier symptom than trismus. The severer the case the more rapidly rigidity and trismus increase, and they tend to be more and more increased by manipulation or movement. This reflex excitability gradually becomes greater until sudden reflex spasms occur in which the jaws are forcibly clenched, the chest is fixed so that respiration is temporarily stopped, and the spine arched backwards in the position called opisthotonos. These reflex spasms may come on any time from twelve hours to seven days after the first onset of trismus, and the earlier they begin the more severe and fatal is the disease likely to be. The time between the first appearance of trismus and the establishment of reflex spasms (period of onset) can generally be estimated to within six hours, and is a good indication of prognosis. If it is less than forty-eight hours, death usually takes place within six days whatever is done; but if it is longer than this there is a good chance of recovery, which becomes greater as the period lengthens. In severe cases which recover, reflex spasms may continue to occur over a period of three weeks, increasing to a maximum, waxing and waning for a time, and then gradually getting less severe until they cease, leaving only tonic rigidity, which may persist for several weeks more, but eventually passing off. Reflex spasms are dangerous because they stop respiration, sometimes for several minutes, make feeding impossible, and exhaust the patient. Tonic rigidity in itself is harmless. In fatal uncomplicated cases death usually takes place within six days either from prolonged respiratory spasms, if these are not controlled, or from cardiac failure when toxin reaches vital centres in the medulla. It is often preceded by hyperpyrexia.

Clinical Types

For convenience tetanus may be divided into five types, as follows:

Type 1, in which at onset there is local tetanus in the wounded limb with rigidity or twitching, and in which the typical generalized picture appears later.

Type 2, in which there is generalized tonic rigidity without reflex spasms.

Both these types are mild, and usually occur after prophylactic antitoxin.

Type 3, like *Type 2*, but the stage of tonic rigidity passes into the stage of reflex spasms described above.

Type 4—splanchnic tetanus—in which the muscles of deglutition and respiration are involved early and generalized symptoms are slight or do not have time to appear. This form is very rare and fatal, and usually follows a visceral infection.

Type 5—cephalic (Kopf) tetanus—in which paralysis or irritation of cranial nerves, most commonly the seventh or oculomotor, appear early and generalized tetanus only later. This type usually follows wounds of the head or neck, and may be regarded as a form of local tetanus. If the wound is not severe the prognosis as a rule is good.

Diagnosis

"Lockjaw" is a good name for tetanus because it emphasizes the most significant early symptom. A small girl was told that if she cut herself between her thumb and first finger she would get lockjaw. A few weeks later she did this, and a few days afterwards, finding that she could not open her mouth, she told her parents she had lockjaw. It was only after she had been seen by three doctors, and after a good deal of delay, that her diagnosis was confirmed and treatment begun. The onset of painless stiffness of the jaw after a wound should always suggest the possibility of tetanus. Other important confirmatory signs in the early stages are: rigidity of the spine, with slight arching of the back so that the flat of the hand can be passed between the bed and the small of the back without difficulty, and almost painless but board-like abdominal rigidity, increasing when the trunk or limbs are manipulated or the patient attempts to move. The facial expression is also an important diagnostic sign.

Differential Diagnosis

From the early occurrence of trismus, tetanus has to be distinguished from local affections of the mouth such as an impacted wisdom tooth, tonsillitis, peritonsillar abscess, or glands about the angle of the jaw. All these conditions are painful, while the trismus of tetanus is comparatively painless; and this fact, along with a careful examination of the mouth and throat, is usually sufficient to solve the difficulty. Local tetanus may be mistaken for neuritis or arthritis, but, in the former, stiffness or twitching is more marked and pain is less. Cephalic tetanus with cranial nerve palsies has to be distinguished from Bell's palsy, polio-encephalitis, and meningitis, and the generalized form from meningitis and encephalitis also. Tuberculous meningitis may simulate it very closely. In these latter, diagnostic lumbar puncture may be necessary. When abdominal rigidity comes on before trismus, tetanus may be mistaken for acute abdominal conditions such as appendicitis or perforated gastric ulcer, but pain is a more pronounced early symptom in the latter.

Finally, the reflex spasms have to be distinguished from other forms of convulsion such as hysterical or epileptic fits and those of strychnine poisoning, in which trismus is a less marked feature. In these a careful history will often solve the problem.

Treatment

This is best considered under four headings: Antitoxin, treatment of wound, control of reflex spasms, and feeding and general management.

Antitoxin

When toxin is exerting its effect on the nervous system it is extremely doubtful if it can be neutralized by antitoxin; consequently if a lethal dose is so acting before antitoxin is given, this treatment cannot save the patient. Happily in most cases antitoxin can be given early enough to prevent the

absorption of a lethal dose. The first need, therefore, is to flood the circulation with antitoxin as soon as possible after the first symptoms have appeared, so that toxin in the blood and lymph which is still accessible to antitoxin is neutralized and any further absorption from the wound prevented. A single large dose (200,000 I.U.) should be given by the intravenous route or, if this amount is not available, as large a dose as possible. In the case of wounds of a limb 10,000 units may also be injected into the subcutaneous tissues round the wound or deeply girdling the limb above the wound.

Repetition of Doses.—In cases with slight wounds I am very doubtful if anything is to be gained by giving further antitoxin. With severe wounds 50,000 units should be repeated at weekly intervals or before any operation, until reflex spasms have ceased. It is difficult to see what is to be gained by daily repetition of injections, for it has been shown that antitoxin continues to circulate in large amounts for at least ten days after a big initial dose, and repeated injections are a worry to the patient and tend to increase spasms. To continue to give antitoxin after the reflex spasms have ceased and only tonic rigidity remains is a sheer waste.

Other Routes.—Antitoxin is also given intrathecally and cisternally, usually in combination with intravenous injections, and good results are claimed by some workers; the evidence, however, is conflicting. In my opinion nothing is gained and a great deal may be lost through the excitation of spasms by such procedures, and they are better avoided, particularly under Service conditions. If for any reason antitoxin cannot be given intravenously it should be given intramuscularly.

Before using antitoxin inquiry should be made as to previous serum therapy and for any history suggesting a tendency to allergy or anaphylaxis, and in cases of doubt desensitization should be carried out. If no contraindications are found the serum may be injected direct into a vein and need not be diluted or warmed. A solution of 1 in 1,000 adrenaline should be at hand in case of reactions.

Treatment of Wound

The wound should be touched as little as possible for an hour after antitoxin has been given. The aim should then be to convert an anaerobic into an aerobic infection by thorough drainage and removal of necrotic and infected tissue and foreign bodies. Wounds should be irrigated four-hourly with hydrogen peroxide through Carrel's tubes if for other reasons this is suitable, and dressings should not interfere with free drainage. It is thought that the practice of early excision of wounds, which was introduced in 1917, lowered still further the incidence of tetanus. The occurrence of tetanus may also be a factor in favour of amputation when this is considered for other reasons.

Control of Reflex Spasms

Tonic rigidity in itself is not dangerous; but reflex spasms exhaust the patient and stop respiration, and it is these which need to be checked. The rate at which they develop is the best indication of their approaching severity. If they begin within forty-eight hours of the first onset of trismus they usually increase rapidly both in frequency and in intensity, and may be very difficult to control. In such cases it is well to make preparations to deal with a sudden prolonged spasm should one occur without warning.

In mild cases and in the early stages before reflex spasms have begun, doses of potassium bromide up to 20 grains two-hourly or sodium luminal up to 1 grain four-hourly usually suffice for sleep and rest and to lessen the excitability of the nervous system. When reflex spasms come on they are most easily controlled by avertin or paraldehyde, given rectally as for basal anaesthesia. The dose of avertin should be between 0.07 and 0.1 c.cm. per kg. of body weight (estimated), and this usually stops spasms for from four to six hours. After this the dose has to be repeated, and it is sometimes necessary to give three such doses a day, and often two a day for some days consecutively. The effect in the individual patient and the frequency and severity of the spasms when they recur are the best indications for dosage. The dose of paraldehyde should be at least 1/2 drachm per stone dissolved in normal saline (1 drachm to 2 oz.). Both avertin and paraldehyde tend to produce cyanosis, and this should be controlled by

warmed oxygen given by catheter or mask. Evipan intravenously is useful, particularly for sudden severe spasms which stop respiration completely; as also is nembutal, $1\frac{1}{2}$ grains by mouth and repeated. Morphine and heroin should be employed sparingly, but are useful at times. Nitrous oxide and oxygen, chloroform, and ether are also occasionally of value, but if spasms are severe they cannot be inhaled. With all sedatives the risks of overdosage have to be weighed against the risks from the disease. If there is moisture in the lungs atropine should be given subcutaneously.

Feeding and General Management

Reflex spasms are excited by external stimuli of all kinds, and these must be reduced to a minimum. Patients should be kept as quiet as possible in a darkened room or at least be screened from others. The bedclothes should be cradled, and all necessary manipulations, such as feeding and the giving of injections, done while the patient is most deeply under the influence of sedatives. In severe cases a sister or orderly should be near the patient all the time. It is important to maintain strength by giving plenty of nourishing fluids and feeds whenever opportunity occurs. Glucose or cane sugar in water or weak lemonade, milky foods; egg and milk, broth, and bovril are all good, and the last two tend to make up any loss of chlorides from excessive sweating. When trismus is severe it is sometimes necessary to extract teeth before feeding is possible, and occasionally feeding by stomach tube is necessary. Avertin is often of great value in relaxing the jaw, and the attempt to feed should be made from one to three hours after the drug has been given. Feeding by rectal drip is also occasionally useful.

On active service the question may arise whether patients should be evacuated. The decision may be a difficult one and, apart from the wound, depends chiefly on the severity of the reflex spasms and the stage which the disease has reached. Patients who have not developed any reflex spasms four days after the onset or whose spasms have almost subsided can usually be moved with safety. Those who are developing them within four days or who are still having severe spasms should stay.

Complications and Prognosis

Pneumonia is a not uncommon and a fatal complication, and, if suspected, sulphapyridine should be given. Retention of urine occasionally occurs, and hyperpyrexia is usually a terminal event. The latter should be treated by antipyretics and tepid sponging. Sequelae of tetanus are rare.

In arriving at a prognosis the factors that have to be considered are: whether there has been active or passive immunization; the severity, degree of infection, and site of the wound; the length of the incubation period and period of onset; the age and physique of the patient; the presence of intercurrent disease or injury, loss of blood, or exhaustion; the severity of reflex spasms; and the conditions under which treatment can be carried out. An incubation period of more than seven days, a period of onset of more than two days, the presence of local tetanus, and a slight wound only, are all in favour of recovery. Other things being equal, the prognosis is better with wounds of the leg than with those of the head, neck, or arm. In war, intercurrent weakness from the severity of the wound itself, from haemorrhage, and from exhaustion has to be weighed against the beneficial effects of active or passive immunization.

- * The 34th annual meeting of the Medical Benevolent Society for the East and North Ridings of the County of York (including the City of York) was held in the rooms of the York Medical Society on Sept. 30, Dr. C. H. Milburn of Harrogate presiding. The treasurer's report showed the society to be in a sound financial position, and it was decided to make further investments in Government securities. Since the last meeting two members had died, and sympathy was expressed with the relatives. One new member was elected. The annual grant to the dependants of an invalid member was renewed. The following officers were elected for the year 1942-3: Dr. M. Jacobs, Hull, President; Dr. Howard Jackman, York, President-Elect; Dr. G. F. Longbotham, Middlesbrough, Vice-President; Dr. D. Matheson Mackay, Hull, Hon. Treasurer; and Dr. W. W. A. Kelly of 92, Micklegate, York, Hon. Secretary, to whom inquiries may be addressed.

SOME OBSERVATIONS ON SENDING ASSISTANCE TO BOMBED TOWNS

BY

F. WILFRED WILLWAY, M.D., M.S., B.Sc., F.R.C.S.

*Temp. Hon. Assistant Surgeon, Bristol Royal Hospital;
A.R.P. Officer, Bristol Royal Infirmary*

I put forward these observations with some diffidence though they are written dogmatically. Many may be more experienced, but I have not had the good fortune to read their papers on the subject. If this paper provokes a discussion it will have served some useful function. Apologies are offered for the overuse of the "first person singular," but this is a personal report, and the observations and deductions are in no sense official.

In the winter of 1940-1 Bristol received its average share of air raids. I was active in all save one of these and learnt first-hand lessons. As the winter of 1941-2 approached we were all keyed up to face the bombardments that never came. Instead raids were made on smaller neighbouring towns, notably Weston-super-Mare and Bath. It is obviously possible to view a raid on a neighbouring city with considerable detachment. Such vision may be useful, and I had the good fortune to assist in the "good-neighbour policy" in all these raids. In each case we spent part of the time in the "blitzed city" and part of the time organizing in the "rescuing city." This was possible as the raids were repeated on the second night in each town.

On the whole the good-neighbour system worked well, but it is felt that many improvements could be made. I do not propose to detail what was done—this might be invidious—but the following commentary may be of interest. Admittedly some of it is painfully obvious, but most British cities found that it took actual bombs to drive the lessons home!

Communications

It is clearly vital to establish some sort of contact between the city needing rescue and the rescuer. It is true that rescue parties can be sent fully equipped and self-contained to a considerable degree, but usually fresh supplies, relays of helpers, etc., are necessary. As a busy voluntary hospital, heavily worked and with a war-depleted staff, we do not carry supplies sufficient for "self-contained" as opposed to "shuttle or circulating" rescue work. A fully self-contained unit would obviously need rations, a field kitchen, water, etc. None of this is available, nor is it considered desirable, though certain improvements along these lines could be made. In the case of the Weston raid, phone communication with Bristol was good though the line was sadly overworked (raid excitement) and messages were repeated unnecessarily—for example, a request for more sulphapyridine was separately phoned up by four officials.

In the Bath raids communication was bad. (This is written in terms of the hospital personally visited.) I arrived there at 2 p.m. on the day after the first raid. (I was then leading a surgical team under orders from the Group M.O.) We saw cases and operated till 12 midnight, staying at the hospital for the second raid. At 4.45 a.m. casualties were again arriving. Since leaving Bristol we had been unable to contact Bristol or the other Bath hospitals. The sole contact was the visit of a dispatch rider at 4 p.m. on the first day. We had then been there only two hours and were in no position to know what extra help might be needed, as the problem had only just been surveyed—e.g., some 60 casualties in an unfamiliar hospital with facilities for two teams to operate. At that time we were not *au fait* with the laundry position or the drug and dressing position.

At 6 o'clock on the morning of the second raid I returned to Bristol in my own car to obtain further aid. Clearly this was wrong. It should not be necessary to detach a surgeon to act as a dispatch rider. It is now obvious that when the dispatch rider called at 4 p.m., the aid I should have requested was the honour of a further inspection at, say, 10 p.m., when we would have understood the situation better. We should then have known enough to appeal for rations. The large team