Current Practice

CHILD CARE IN GENERAL PRACTICE

Deafness

EDITH WHETNALL,* M.S., F.R.C.S.

Methods of detection, diagnosis, and management of deafness in children depend so entirely on a knowledge of the processes involved in communication by speech and in the acquisition of this skill that I propose to consider them first. It will be found that the practical steps to be taken in dealing with patients flow naturally from this knowledge.

Hearing and Speech

Until recent years it was thought that the hearing of the child born deaf was, in most cases, of no value in enabling him to learn to talk and this idea was so ingrained that little attempt was made to use what hearing he had. Now it is known that very few, if any, of these children have no hearing at all and that even the smallest amount of hearing is of the greatest value. The explanation of this depends on realizing how a language is learnt. Too few people know how this is done, for an individual takes in his own native language at such an early age and so painlessly that he is completely unaware of his knowledge. A vast store of linguistic information is laid down in the auditory

areas and it is on memories of this that speech depends. This store varies from one language to another; thus an Englishman does not understand French without having first "learnt to speak French" though he can hear the sound plainly enough. To learn a language involves learning its linguistic system, for it is this system which varies from language to language.

The development of hearing, speech, and language in the child is very closely bound up with his mental development. In the first years of life he is developing the ability to perform the skilled movements which constitute speech and to recognize and interpret the acoustic signals which accompany these skilled movements when made by other people. At this time the child is developing his mental powers in various ways, and there is plenty of evidence that this development of mental capacity is very much dependent on the learning of speech.

Development of speech involves not only the learning of skilled articulatory movements but the mastery of cues for the recognition of speech, the building-up of the store of linguistic knowledge, and the gradual organizing at the psychological level

of many processes that form the basis for both the production and the reception of speech. These processes are illustrated by the speech communication chain shown in the Figure.

It is easier to consider hearing and speech in the adult before considering the learning process in the child.

Speech is organized by the speech centres in the brain, and the events that follow the activity in a speaker's brain are well-known—psychological, physiological, physical, and then in the reverse direction to the listener's brain. No matter what these transformations are, the listener can decode the message back to the form in which it was originally sent only if both he and the speaker have a large memory store of language material in common. When this is the case the speaker and listener are said to speak the same language.

Organization of Language in Brain

The psychological processes make up what is called "language." Every language is a system—a system of units of differing size that combine together in a manner regulated by the structure of the system. The smallest unit in spoken English is the phoneme, the equivalent of the single letter in writing, and the total number of English phonemes is just over 40. The next linguistic unit in order of size is the morpheme, a term used to refer to units that have a mainly grammatical function (*dictate*=one morpheme, in *dictating* another morpheme is added). The combining of morphemes produces the units recognized as words, and they in turn form sentences.

The knowledge of all these units and a great deal of statistical information about the way in which they combine is stored in the memory of every speaker and listener, though they are not usually aware of it. Acquisition of this information is not a random process and the brain acquires at a subconscious level a vast store of linguistic information as the result of practice in listening early in life. Once the system is learnt it is not necessary to hear every word in order to understand speech. The listener is able to guess successfully words he has not heard because he knows the rules that string words together to make sentences, and at any point in a sentence he knows what word is more or less likely to come next. Very considerable knowledge is available to him of the probabilities governing the combinations of phonemes and morphemes with one another.

Whether or not the brain is in possession of this store of linguistic knowledge constitutes the fundamental difference between congenital deafness and deafness acquired after speech has been learnt.

Control of Speech Movements

The function of speech is to convert language into a transmissible form, and in this process speech appears as a predominantly motor activity which has acoustic consequences.

* Director, Nuffield Hearing and Speech Centre, the Royal National Throat, Nose, and Ear Hospital, Institute of Laryngology and Otology, University of London.

The speech mechanism is fed with a flow of language from the speaker's brain which must be converted into acoustic signals. The movements of the articulators maintain a more or less continuous modification of the shape of the resonating cavities of the pharynx, mouth, and nose, and these movements are controlled by means of feed-back circuits. During the production of any speech-sequence information is constantly relayed to the brain concerning the progress and the effect of these movements.

The feed-back circuits important in speech are auditory and kinaesthetic. By far the most important during the learning period is the auditory feed-back. The wave motions set up by the speech movements are transmitted not only to the listener but also to the speaker's own hearing mechanism and thence to the speaker's brain. Thus the distance that the tongue moves, the speed at which it moves, the exact moment at which it changes direction—all these are determined largely on the basis of information supplied by the auditory feed-back.

Once speech is learnt the kinaesthetic feed-back becomes of greater importance as it is through this that the habit of speech is established, and it is for this reason that an adult who becomes deaf will not lose his speech. But during the learning period every vestige of hearing is of the greatest possible importance for the deaf child in order that he may learn to acquire the habit of kinaesthetic feed-back.

Reception of Speech

The reception of speech consists in recovering language units from the transmitted signals. The listener, if he is to understand a message, must possess a memory store of linguistic information similar to that held by the speaker and be able to decode it. The brain is presented with a continuous sound-wave which the brain segments into discrete language units. An incoming pattern is compared with a stored repertory of patterns and the knowledge of what is coming next or is likely to come next is all-important, for the listener is all the time filling the gaps by using his linguistic knowledge. It is important to realize that learning to recognize sound is carried out entirely in the brain. The capacity of the brain to organize information is quite extraordinary, and, while the amount of information sent to the brain may be less from a deaf ear, it is what the brain does with this information that is important in learning to recognize words. Provided that the deaf child is exposed to sounds that are loud enough and occur often enough his brain will learn to recognize them in the same way as the brain of the child who hears normally.

Learning to Hear

Normal hearing implies the ability to understand sounds. Hearing in this sense is not present at birth but is learnt during

the first years of life provided that certain conditions are present. It is because the existence of these conditions is not realized that speech is said to develop spontaneously. If these conditions are provided for the deaf child he will learn in the same way as the hearing child.

During the first year of life changes take place in the hearing which are dramatic. The hearing of the infant at birth is reflex; it is quite different from the comprehension hearing of the adult or child who has learnt to speak, and is analogous to that of lower animals, the primary function of which is protective. The analogue of this primary reflex in the newborn infant is the "startle reflex" in response to a loud unaccustomed sound.

The process of learning comprehension hearing starts at birth and consists of learning to recognize sound and interpret it. Sounds differ from one another by the different patterns of frequency and intensity of the sound-waves composing them and sounds come to be associated with meaning. As the result of experience, reflex hearing becomes inhibited as the infant learns to ignore a sound that neither harms him nor does him any good. At 5 months the infant may not respond to a loud bang or at most looks up once and then ignores it. With another but new sound the infant will keep still and listen, look up or towards the sound once, but not again. By 8 months the reflex startle is inhibited and a new response develops to a sound heard dozens of times and associated with pleasure—his mother's voice, sounds of food, his own rattle, or rustling paper. He looks in the direction of the sound; an expression of pleasure indicates understanding. The greatest response is to speech. The infant also uses his own voice, copying first the intonation patterns and gradually improving the imitation until it is recognized as speech. Any approximation to a word is received with approbation and so the infant is encouraged to repeat the sound and perfect his performance. These stages of development are shown more fully in the Table below.

All learning depends on the presence of certain essential conditions. The conditions for learning to hear are that the sounds must be heard (1) loud enough, (2) often enough, (3) in association with their meanings, and (4) at the right age—the first months and years of life when there is a special facility for learning to hear. Less hearing is required once speech is learnt than during the learning process. The hearing loss in many adults is such that, if they had been born deaf with a similar loss, they could not have developed speech without special help.

Detection and Diagnosis

Detection and diagnosis of deafness are based on knowledge of the normal infant's response to sound (see Table). Any delay or abnormality in these responses should cause suspicion.

Normal Responses to Sounds in Infancy at Different Stages of Development

Approximate Age	Startle Reflex (to sounds over 75 db.)	Reaction to Sound	
		Unfamiliar	Familiar
Immediate post-natal period	Present	Nil	Nil
1 month	Present	If spontaneous activity present, sound of voice, bell, or rattle causes activity to cease	
3 months	Starting to become inhibited	If spontaneous activity present, sound of voice, bell, or rattle causes activity to cease	
4-5 months	Varying degree of inhibition	Many sounds unfamiliar, looks only once. If spontaneous activity present sound causes this to cease	Turns eyes in direction of sound—mother's voice, bunch of keys, rattle, cup tapped with spoon, rustling paper
8 months	Usually inhibited; may be one response	Very few sounds unfamiliar	Localizes—i.e., turns briskly and looks at—many sounds. Voice acquiring intonation of normal speech. Responds to mother's voice by imitation.
1 year	If child fails to localize familiar sounds deafness should be suspected. Failure in speech development is also a danger sign.		
2 years	If child fails to understand speech or to talk deafness must be suspected.		

It is best to make the tests with the infant sitting on the mother's lap, so that the sounds can be made from behind. Sounds such as described in the Table are used, testing alternate ears first with each new sound, for the infant may respond only once, while the sound is unfamiliar, and this can give rise to the idea that the ear tested second is deaf.

The term detection implies setting out to find cases of deafness even where deafness is not actively suspected. The disability can be effectively overcome provided the infant is taught during the physiological period of learning to hear. Detection should be at the beginning of this period. To the question "Who should detect?" the answer is, ideally, the parents, who should be made aware of the fact that the infant is learning to listen and hear in the first year of life.

The mother is frequently the first person to suspect that her infant is deaf and she must be given help straightaway in obtaining advice from an otological clinic. Mothers are regrettably still told that the diagnosis cannot be made in a child so young and that even if it could be there is nothing that can be done until the child is older.

The reasons often given by mothers for suspecting deafness are (1) Failure to respond to sound; the infant does not jump or turn to a loud sound, or smile on hearing the mother's voice. Failure to respond to sound has been observed as early as two days after birth and is commonly noted by 3 to 6 months of age. This failure to respond to sound is the complaint most frequently made by mothers. (2) Failure to use the voice in response to the mother's voice, failure to talk or, if talking, defective speech, are the next most common complaints. (3) Lack of facial expression and lack of awareness of events taking place are frequently noted.

Much information about the degree of deafness can be obtained from the history of the infant's reaction to sound—is sound ignored altogether, or will he look up to some sounds if not to others? Does he use his voice in babbling and does the tone approximate to normal? Behaviour is important, as a delay in reaching milestones indicates backwardness rather than deafness.

No test method can be set up to ascertain in a single test the exact extent of an infant's hearing loss, but the temptation must be resisted to delay screening tests until too late an age, when it is safe to be definite. What is required are tests made soon after birth which should be repeated on subsequent occasions during the first and second year if any doubt is present. Post-natal clinics can do a great deal to spread this awareness but the family doctor has an even more important part to play. Cases "at risk" should of course be looked for, but these account for only 50% and a routine test should be made in every infant—after all, it only takes a few minutes and a lifelong disability may be avoided.

Assessment of Hearing

It is essential to remember that the deaf child always has more hearing than is indicated by the first tests of hearing. Sounds have been heard too infrequently for the brain to learn them and so no meaning is attached to them. There may be no response at all, the startle reflex may not occur, the infant merely looks up on hearing the sound. Very few sounds are familiar—that is, the infant responds once to a few sounds. Localization is not possible.

Few children, if any, have no hearing at all. Most have some, but it is not until training of the hearing has been carried out for at least a year that hearing tests begin to indicate the true ability to hear. The distance from the ear at which speech sounds are heard at the first test is the distance at which speech can be understood after training. Even if the child hears only at the meatus the help of a hearing-aid will give levels of hearing which together with lip-reading will enable the child in

many cases to go to an ordinary school. Deafness reduces the occurrence of sounds which are loud enough for hearing to occur. To assume that because a child has no speech he cannot develop speech is hopelessly incorrect. Bring the sounds within reach and the child will hear them. Even if no response is obtained at first, total deafness should not be diagnosed until training with an aid has been carried out for at least a year.

Differential Diagnosis

Difficulty will arise in children who for reasons other than deafness fail to respond to sound, fail to talk, or wear a blank expression. The effects of sound deprivation or lack of individual attention, of illness, dullness, or prematurity, must be carefully observed and understood by those responsible for the diagnosis of deafness. Any of these conditions may result in a delay in hearing development and thus in talking. The commonest difficulties are:

1. Failure to respond to loud sounds is a common finding in a child with normal hearing at the age of 6 to 8 months.
2. Delay in development of response in relation to age is common in children who are born prematurely or suffer a severe illness.
3. The child who is neglected fails to respond to sound, and may appear deafer than the deaf.
4. Response to faint sounds occurs only when learning has taken place. Absence of response in the third or fourth month does not indicate deafness.
5. Failure to respond to sound is sometimes seen when parents suspect deafness and stop talking to their infant, thinking it is of no value. Amateur tests of hearing may also have been used—loud bangs to see if the child will turn. This inhibits any response at all.
6. Delay in hearing and speech development is common in mental retardation. In these cases responses may be those of an infant months younger, arrest of learning taking place at the mental age of the child. Responses to faint sounds may be normal but the ability to learn to understand speech or to talk is not present. If there is doubt as to the diagnosis these children should be trained as though deaf in order to see if they can make progress. Tests of hearing and of intelligence must, during this time, be repeated and some time may be needed to make a final decision. The assistance of a psychologist trained in the problems of the deaf child is quite indispensable in these cases.
7. Delay in speech development due to late maturation is commonly seen and can be correctly diagnosed with the assistance of a psychologist.

Management

As soon as deafness is diagnosed the parents must be instructed about the process of learning to hear so that they can understand how to help their child. Only they can give the individual attention that is essential. Every mother can give this attention provided that her role is explained to her and she is seen from time to time to help her to deal with problems and to remind her of the principles of learning to hear. The provision of a home-help may make all the difference and is as necessary and reasonable as in cases of disability from other causes.

The mother must understand that evidence of even a vestige of hearing is of the greatest value in learning to speak. Even the child who appears totally deaf must be treated as though

some hearing were present. The mother must be reassured that first tests will indicate less hearing than is really present and that the hearing will appear to improve with training. She must realize that once speech is learnt it is heard more easily and the deafness is thus less of a handicap. Some mothers can make easy conversation but others must be shown how to seize and take every opportunity to use speech themselves in order to encourage the child's use of speech. Mothers should attend with their children once a week or once a fortnight at a hearing and speech centre or clinic for deaf children so that they can be taught by a clinical teacher of the deaf, or by a speech therapist, how to carry on helping their children. A hostel for mothers who live too far away for such visits is available at the Royal National Throat, Nose, and Ear Hospital. Here, in a course lasting two weeks, the mother learns about the management of her deaf baby.

Hearing-aids

The function of a hearing-aid is to make sounds louder. Once the infant gets his aid he must still go through the lengthy process of learning to hear the sounds which now reach him through it. The child with normal hearing starts this learning process at birth and does not produce speech until he has been listening to it for a year. The child with impaired hearing must be given at least a year with a hearing-aid before he is expected to produce speech. The parents must be encouraged to resist the temptation to give up just before the child is about to start to talk. This is another way in which the family doctor can give invaluable help in the care of the child who is born deaf.

Two ears are better than one and so two aids clearly should be better than one. This, in fact, is found to be so, for summation occurs, enabling the very deaf child to hear sounds louder than with one ear. Almost more important than this is the ability to localize sound and as a result of this to be able to hear more easily in a noise. The young child with two aids learns more easily and more quickly and acquires better speech. Clearly it will be of the greatest use, too, for the deaf child in the noisy classroom to have two aids.

The aids should be worn continually, and it will be found that the child soon likes his aid and does not want to be without it, and will protest if it is taken off for any reason.

If a child fails to make good progress in learning to talk it may be the hearing-aid which is at fault in one or other of the following ways.

(1) Insufficient amplification may result in poor quality of the voice and slow progress. A more powerful aid is necessary. (2) Intolerance of sound may make a child refuse to wear his aid, and an aid incorporating automatic volume control is necessary to reduce sounds to a comfortable level. Deafness due to maternal rubella, Rh incompatibility, or meningitis are associated with this intolerance of loud sounds. (3) A badly

fitting insert causes "howling," and the remedy is to obtain a new insert as quickly as possible, as the aid cannot be used with sufficient amplification while it is howling.

The child's whole personality will alter in a few moments when changing from a wrong aid to a good one. The child hears and understands at levels which are comfortable and he starts at once to use his voice. Speech deteriorates rapidly with the wrong aid.

Schooling

Pre-school Children

Admission to a day nursery at the age of about 3 years with children who hear normally is helpful, for young children use simple speech in loud voices—ideal material for so-called auditory training. Moreover, children at this age mix well. The nursery teacher will be a most valuable ally, and her help must always be obtained.

Children Over 5 Years Old

Those children who make the grade to attend an ordinary school will still need the continued help of the otologist, general practitioner, and teachers in order that there is no strain and no falling behind. In this way the child's confidence and his progress are ensured.

Children Diagnosed Late

Facilities must be available for children who have useful hearing, but have not learnt to hear. A hostel where individual training is given is available for such a child, again at the Royal National Throat, Nose and Ear Hospital.

A small "partially hearing unit" provides a temporary solution for some of these children. It must be small, for the idea of such a unit is that the child, while receiving special auditory training and extra help in education in the unit, is gradually transferred for more and more of his lessons to a class with children who hear normally.

I would like to thank Dr. Raymond Greene for reading this paper and for his valuable criticism and comments. The figure illustrating the speech communication chain is reproduced from *The Deaf Child*, by Edith Whetnall and D. B. Fry, by permission of William Heinemann (Medical Books) Limited.

Correction.—We much regret an error in Dr. D. H. Garrow's article on "Acute Respiratory Infections in Childhood" (30 January, p. 297). Under the heading of "Whooping-cough" the third sentence in the first paragraph should have read: "A pernasal swab is the best way to isolate *Bordetella pertussis*, which is, however, a delicate organism."