CHARLOTTE M. ANDERSON: LONG-TERM SURVIVAL WITH SIX INCHES OF SMALL INTESTINE

FIG. 1

FIG. 1.—Barium x-ray film at zero time, showing large stomach.

FIG. 2.—Barium x-ray film after 15 minutes, showing extent of small intestine and barium already in caecum.

FIG. 2

H. GARLAND *ET AL.*: IATROGENIC VERTEBRAL ARTERIOVENOUS FISTULA

FIG. 1

FIG. 2

FIG. 1.—Right vertebral arteriogram via axillary artery. Contrast has passed directly into greatly dilated vertebral veins and into the deep cervical vein. There is already some contrast in the vertebral veins of the opposite side. Note the dilated vertebral artery, which narrows suddenly at C 4 level, the probable site of the fistula.

FIG. 2.—Arteriogram taken one second after Fig. 1. Contrast is draining downwards through dilated vertebral and deep cervical veins on the right side. Note also the small veins extending across the midline at mid-cervical level and the outlining of the left vertebral vein (arrowed).

our patients had a hyperparathyroidism without bone disease; but this is the most provocative aim of this article, as they had none of the known diseases with elevated calcium plasma level (sarcoidosis, vitamin-D poisoning, and so on).

Summary and Conclusions

In six stone-forming patients the intestinal absorption of ^{45}Ca was studied: it was found that all six patients examined were calcium-hyperabsorbers.

It is concluded that determination of the intestinal absorption of ^{45}Ca in patients with formation of calcium concretions proves such patients to be calcium-hyperabsorbers.

The principal metabolic abnormality in some patients with calcium nephrolithiasis could be an increased intestinal absorption of calcium.

REFERENCES

- Albright, F., Bauer, W., Claflin, D., and Cockrill, J. R. (1932). *J. clin. Invest.*, **11**, 411.
 Anderson, J., Dent, C. E., Harper, C., and Philpot, G. R. (1954). *Lancet*, **2**, 720.
 Aub, J. C., Tibbets, D. M., and McLean, R. (1937). *J. Nutr.*, **13**, 635.
 Caniggia, A., Gennari, C., Bianchi, V., and Guideri, R. (1963). *Acta med. scand.*, **173**, 613.
 Fourman, P. (1960). *Calcium Metabolism and the Bone*. Thomas, Springfield.
 Henneman, P. H., Dempsey, E. F., Carroll, E. L., and Albright, F. (1956). *J. clin. Invest.*, **35**, 1229.
 Jaworski, Z. F., Brown, E. M., Fedoruk, S., and Seitz, H. (1963). *New Engl. J. Med.*, **269**, 1103.
 Lafferty, F. W., and Pearson, O. H. (1963). *J. clin. Endocr.*, **23**, 891.
 Lenzi, F., and Caniggia, A. (1962). *Fisiopatologia e Clinica delle Osteoporosi diffuse*. 63rd Congr. Soc. Ital. Med. Int., Roma. Pozzi, Rome.
 Lichtwitz, A., and Parlier, R. (1959). *Presse méd.*, **67**, 134.
 — de Sèze, S., Hioco, D., Miravet, L., Lanham, C., and Parlier, R. (1963). *Ibid.*, **71**, 165.
 Morgan, H. G., Mitchell, R. G., Stowers, J. M., and Thompson, J. (1956). *Lancet*, **1**, 925.
 Stapleton, T., Macdonald, W. B., and Lightwood, R. (1956). *Ibid.*, **1**, 932.

Medical Memoranda

Iatrogenic Vertebral Arteriovenous Fistula

[WITH SPECIAL PLATE]

Brit. med. J., 1965, **1**, 429–430

Arteriovenous communications between the vertebral vessels are usually a result of trauma. The common mechanisms are stab wounds, gunshot wounds (Aronson, 1961), or, occasionally, indirect injury to the cervical spine (Faeth and Dueker, 1961). There is one report of a "spontaneous" vertebral arteriovenous fistula (Gooddy and Schechter, 1960). Attention is drawn to a further cause—namely, percutaneous vertebral arteriography. We can discover in the English literature only two reports of a similar occurrence.

CASE REPORT

The patient, a 37-year-old housewife, had since the age of 18 suffered from a variety of psychogenic symptoms. At the age of 23, when in the second trimester of pregnancy, she developed pain and impairment of vision in both eyes, which had regressed by the end of pregnancy. Seven years later she developed an unsteady gait, tending to stagger to the right. This persisted, but two years later her left leg became numb and weak and she tended to trail the left foot. Two years ago she complained of increasing deadness of the left leg, associated with tingling in the left arm and leg, and renewed aching and blurring vision in the right eye. These and many non-organic symptoms led to her subsequent admission to another hospital for investigation. It was thought that she was suffering from disseminated sclerosis, but percutaneous right carotid and bilateral direct vertebral angiography were attempted to exclude a possible vascular cause for her symptoms. The radiographs were reported as being normal, but the patient was informed that the puncture of the right vertebral artery had been unsuccessful, though a successful examination had been obtained on the left. Five days later she experienced a terrifying, loud, rushing noise in the occipital and right temporal region, accentuated by stooping and by turning her neck. This remained unchanged, and six months later she was admitted to the Neurological Unit of the General Infirmary at Leeds for investigation because the noise was becoming increasingly difficult to tolerate.

Examination revealed pallor of the right optic disk, a central scotoma, and visual acuity of Jaeger 8 in the right eye. She had pyramidal signs in the left arm and leg, and a mild sensory impairment, of dubious significance, on the left side of the body. The most striking finding, however, was an easily palpable thrill and a very loud continuous bruit audible over the whole neck, but maximal on the right side; this was loudest over the upper end of the internal carotid artery, and was diminished by digital compression of the right subclavian artery. No bruit was audible over the orbits or skull. The blood-pressure was 130/80.

It was felt that in addition to disseminated sclerosis the patient had sustained an arteriovenous fistula. It was not possible to decide with absolute certainty which vessels were involved, but the fact that the bruit was reduced by pressure on the right subclavian artery but not by occlusion of the right carotid, and the history of a failed right vertebral puncture, were thought to be strongly suggestive of a lesion affecting the vertebral rather than the carotid artery. Radiological investigation was undertaken to confirm the diagnosis in the hope of demonstrating an operable lesion.

Radiological Investigation

Preliminary plain film examinations of the skull and neck showed no abnormality. A right axillary arteriogram was carried out under local anaesthesia and a red Kifa catheter positioned with its tip in the vertebral artery. Two injections each of 8 ml. of 45% Hypaque were made at hand pressure and serial films taken. The patient complained of transient slight discomfort in the right side of her neck after each injection.

An arteriovenous fistula between the vertebral artery and its venous plexus was demonstrated. The vertebral artery was enlarged and its proximal few centimetres were more tortuous than normal. Contrast medium passed rapidly into a much-dilated vertebral venous plexus, draining rapidly through enlarged deep cervical and vertebral veins. Some contrast passed across the midline at C4 level through the external vertebral venous plexuses and outlined the plexus of veins around the left vertebral artery below this level. Drainage was rapid and the left vertebral vein could be identified even before injection of contrast into the right vertebral artery had been completed.

The exact site of the fistula could not be identified, but it was thought to be at C4 level, where the vertebral artery narrowed abruptly and maximal venous filling occurred. Later films showed irregular dilatation of several small arterial anastomoses between the vertebral and the deep cervical, and between the vertebral and

ascending cervical arteries. The vertebral artery could not be identified above C 3 level and no contrast entered the posterior fossa. Subsequent aortography demonstrated normal left vertebral and carotid arteries.

Subsequent to investigation the patient had started to tolerate the noise more easily. After careful consultation with a neurosurgeon, a vascular surgeon, and a thoracic surgeon, it was decided to withhold surgical treatment, although this would be reconsidered should her symptoms increase.

DISCUSSION

Sutton (1962) noted this complication on only one occasion in a total of over 10,000 arteriograms, including all sites; this patient "became conscious of a loud murmur in her right ear synchronous with the pulse" soon after several attempts at right percutaneous vertebral puncture. The only other report discovered is that of Olson *et al.* (1963). The presenting features in this instance were boring pain in the right shoulder and arm, motor and sensory loss in C 5 distribution, and a continuous bruit in the neck. The symptoms had begun five weeks after bilateral carotid and direct right vertebral arteriograms.

The development of an arteriovenous fistula is a recognized complication of arterial puncture. It has been recorded after attempted catheterization of the brachial artery and also after femoral arteriography (McAfee, 1957). That it is extremely infrequent, however, is witnessed by the fact that Lang (1963), in a survey of over 11,000 percutaneous retrograde femoral arteriograms, did not report the occurrence of a single example of arteriovenous fistula. Its rarity is perhaps surprising when one considers that most of the arteries punctured lie in close relation to veins. This close relationship is especially marked with the vertebral artery, which is surrounded by a plexiform network of veins in its course through the transverse foramina. One might also suppose that the inability to apply direct pressure at the puncture site after vertebral arteriography could predispose to this complication.

One of us (J. T. L.) has not encountered this complication in over 600 vertebral arteriograms performed by percutaneous direct puncture.

Although it is possible that an arteriovenous fistula in this situation might escape notice, we consider this unlikely, since the symptoms produced are so severe and dramatic in their onset. An understandable reluctance to publicize unfortunate complications may account for a few unreported examples; we have recently heard of at least four of these (North of England Neurological Association, Leeds, 1964).

It is interesting to note that the vertebral puncture was unsuccessful in our case and also in that reported by Sutton (1962), and it may be that trauma to the artery from repeated attempts at puncture predisposes to this complication. It is clearly wise, where difficulty is encountered, to avoid repeated insertions of the needle at the same level. The complication involved the right vertebral artery both in this patient and in the two reported examples; although this is probably coincidental, the fact that the right vertebral artery tends to be smaller than the left (Stopford, 1916), and thus more difficult to puncture, may be a contributory factor.

In our patient there was a five-day interval between vertebral arteriography and the sudden development of a loud bruit; in

Sutton's patient the bruit developed "soon after" the examination, while in Olson's there was an interval of five weeks between arteriography and the onset of symptoms. The reason for this interval is not clear, but it is recognized as being not infrequent in the development of arteriovenous fistula after other forms of trauma at other sites (Callender, 1920). It is possible that the thrombus formed after the initial injury is subsequently dislodged, allowing the fistula to open. A point of further interest in this report was the partial drainage of the fistula via the opposite vertebral vein, and the dilatation of several of the small anastomotic channels between the vertebral and cervical arteries. (Special Plate, Figs. 1 and 2.)

The management of patients with this complication can be extremely difficult. In the first place the diagnosis should be based upon the development of a loud and continuous bruit in the neck, usually audible to both the patient and the doctor. The latent interval between the onset of symptoms and the performance of arteriography should not deter the physician from relating these two events.

If suspicion of this complication arises, it is imperative from both diagnostic and therapeutic standpoints to proceed to indirect vertebral arteriography by either catheterization or subclavian puncture. This may demonstrate a fistula amenable to direct surgical attack. It is also vital to demonstrate the adequacy of the opposite vertebral artery and the patency of the carotid arteries by aortography prior to ligation.

Although it is possible that spontaneous closure of the fistula may occur, it seems probable that most patients will require surgical treatment. A review of operations on the vertebral artery up to 1946 disclosed a mortality rate of over 50% (Aronson, 1961). It would appear that recurrence, or incomplete suppression of symptoms, is likely if proximal ligation of the artery is performed. The procedure of choice is to expose the fistula fully by the lateral approach (Elkin and Harris, 1946), and then, guided by the arteriographic findings, to obliterate the fistula directly. Elkin and Harris successfully treated 10 examples of traumatic vertebral arteriovenous fistula in this way without mortality. It is the general consensus of neurological opinion that operative treatment is indicated in the majority of such patients.

HUGH GARLAND, T.D., M.D., F.R.C.P.

J. T. LAMB, F.F.R.

JOHN PEARCE, M.B., M.R.C.P.

Departments of Neurology and Diagnostic Radiology,
General Infirmary at Leeds.

REFERENCES

- Aronson, N. I. (1961). *Neurology (Minneapolis)*, **11**, 817.
- Callender, C. L. (1920). *Ann. Surg.*, **71**, 428.
- Elkin, D. C., and Harris, M. H. (1946). *Ibid.*, **124**, 934.
- Faeth, W. H., and Duckert, H. W. (1961). *Neurology (Minneapolis)*, **11**, 492.
- Goody, W., and Schechter, M. M. (1960). *Brit. J. Radiol.*, **33**, 709.
- Lang, E. K. (1963). *Radiology*, **81**, 257.
- McAfee, J. G. (1957). *Ibid.*, **68**, 825.
- Olson, R. W., Baker, H. L., and Svien, H. J. (1963). *J. Neurosurg.*, **20**, 73.
- Stopford, J. S. B. (1916). *J. Anat. Physiol.*, **50**, 131.
- Sutton, D. (1962). *Arteriography*. Livingstone, Edinburgh.