It seems somewhat surprising that a period of four weeks in-patient treatment for a relatively severe illness should have been considered suitable by such an authoritative body as the Clinical Psychiatry Committee.—I am, etc.,

Bangour Village Hospital,
Broxburn, West Lothian.

J. WATT.

Paracetamol or Aspirin?

SIR,—In your leader under this heading (24 April, p. 1083) I read "recent ingestion of aspirin has been found in several large series of hospital patients admitted for acute gastro-intestinal bleeding." As one who has recently published just such a series¹ I fear that some may be misled by the rather ambiguous second statement—"in individual patients it is often difficult to prove a connexion between bleeding and taking salicylates." No one, of course, supposes that all aspirin-taking causes overt gastric bleeding, but does your writer mean to imply that he is unconvinced by the connexion between gastric bleeding and aspirin-taking in these long series to which he refers?

Though direct clinical assessment of patients is now held in lowered esteem, what better proof in this matter can we have than that which comes from careful and extended observation? My conclusion, similar to that of Grossman, Matsumoto, and Lichter² ("the gun must be loaded for an explosion to occur when salicylates pull the trigger") was that in many conditions for which aspirin products are commonly taken (nervous headaches, organic and nervous dyspepsias, respiratory infections, etc.) they are particularly liable to erode the gastric mucosa.—I am, etc.,

G. H. JENNINGS.

Edgware General Hospital,
Edgware, Middlesex.

REFERENCES

- Jennings, G. H., *Gut*, 1965, 6, 1.
- Grossman, M. I., Matsumoto, K. K., and Lichter, R. J., *Gastroenterology*, 1961, 40, 383.

Intravenous Regional Analgesia

SIR,—Drs. B. R. Kennedy and others (10 April, p. 954) state in their summary that they do not feel justified in using the intravenous regional analgesia technique because of the high incidence of toxic phenomena they describe.

Neither do I, but may I point out that this does not necessarily apply if the technique is modified? The answer to their problem is to allow the local analgesic to enter the general circulation in small doses, where it will be destroyed quickly; this may be achieved by releasing the tourniquet at the end of the operation and immediately reinflating the cuff for a few minutes. This may be repeated several times if large doses of analgesic have been used.—I am, etc.,

The General Infirmary,
Stafford.

D. EYRE-WALKER.

SIR,—We would like to comment on the two articles on intravenous regional anaesthesia by Dr. B. R. Kennedy and others and Mr. Charles Sorbie and Mr. P. Chacha (10 April, pp. 954 and 957) in view of our own findings on 130 successive cases performed in this casualty department.

In these cases a technique basically similar to that described by Sorbie and Chacha was used, but no patients were premedicated, only upper limbs were anaesthetized, and all were discharged post-operatively. The age range was 12–87 years, with females predominating. Most cases required 30–50 ml. of 0.5% lignocaine, with three requiring only 20 ml. and one 60 ml. In six cases Hyalase (hyaluronidase) 1,500 units was added.

The commonest procedure, by far, performed under this type of anaesthesia was manipulation of a Colles fracture (91 cases); manipulation of dislocated elbows and fingers and other fractures, wound and tendon suturing, and exploration for foreign bodies made up the remainder of the cases. No injection-release times were recorded, but as can be seen from the list above no prolonged procedures were performed.

There were only four failures of satisfactory anaesthesia. One of these four provided the only case of definite toxic phenomenon when the tourniquet accidentally became displaced; the patient had a brief episode of loss of consciousness and some muscle twitching—she was also a known epileptic.

During the period these 130 cases were collected, 110 patients had general anaesthetics, for the same type of case as above, for reasons of patient preference, personalities unsuited to the procedure, age (i.e., too young), or poor veins. Thus, there has been some selection of cases which may account for the low failure and toxicity rate which we have experienced.

While, as Dr. Kennedy and his colleagues have shown, there may be some cases developing slight bradycardia and a fall in blood-pressure (perhaps partly due to pre-medication) we cannot agree that "we no longer feel justified in continuing to employ it." This is a simple, safe, and satisfactory form of anaesthesia, admirably suited to minor limb surgery in a casualty department, on a patient who may have a full stomach or one who can be discharged almost immediately post-operatively.—We are, etc.,

DAVID H. GORDON.
HUGH PERKINS.

Paddington General Hospital,
London W.9.

Potentially Dangerous Forms of Drug Economy

SIR,—Dr. K. Zinnemann's letter (17 April, p. 1067) raises a number of points regarding the chemotherapy of chronic bronchitis. Dr. Zinnemann is concerned lest the long-term use of small doses of tetracycline and ampicillin should encourage the widespread development of resistant strains of *Haemophilus influenzae*. Recent work, however, by May and Delves¹ has shown that the requisite conditions for this to happen, at any rate with ampicillin, may occur with any dose that does not produce a bactericidal effect. When the infection is controlled and the sputum becomes mucoid, the concentration of ampicillin in the bronchial secretions falls to a very low level. Consequently, whatever the dose, continued therapy results in prolonged exposure of surviving organisms to sub-inhibitory concentrations, with their attendant risk of the encouragement of drug-resistant organisms.

Within this context, Dr. Zinnemann's letter should be re-worded as a caution against all long-term chemotherapy in chronic bronchitis. However, whether or not one should deny to a patient the undoubted relief that this form of treatment often provides, merely because resistant organisms *might* be produced, is open to considerable doubt. In fact the development of resistant strains of *H. influenzae* is rare. Although an insignificant trend towards tetracycline-resistance was found in the second, but not the third, trial sponsored by the British Tuberculosis Association,^{2,3} no change whatever occurred in the sensitivity to penicillin; and the risks of ampicillin-resistance should be assessed from behaviour to penicillin, not tetracycline as Dr. Zinnemann has done.

My own procedure, as regards small doses, is to prescribe them only as maintenance therapy in patients whose sputum can be kept free from pus by them; and there is no doubt that the infection in some patients can be controlled completely by 250 mg. twice daily. In these circumstances it seems that the risk of breeding resistant strains of *H. influenzae*, which is undoubtedly very small, is no greater than for any other dose.—I am, etc.,

J. ROBERT MAY.

Institute of Diseases of
the Chest,
London S.W.3.

REFERENCES

- May, J. R., and Delves, D. M., *Lancet*, 1965, 1, 929.
- Francis, R. S., May, J. R., and Spicer, C. C., *Brit. med. j.*, 1961, 1, 979.
- — — — — *ibid.*, 1964, 1, 728.

Abortion Law

SIR,—The leader on abortion law (17 April, p. 1009) ends with the sentence, "Surely the proper way to put out of business the back-street abortionist, who flourishes mainly on the distress of married women, is through cheap and reliable contraception." This is perfectly true, and it is the impression of many general practitioners that "the pill" has considerably reduced the numbers of illegal abortions in their practices. Unfortunately, the side-effects of the pill render it unsuitable for all women. Similarly, the intrauterine contraceptive devices receive high praise from those who can tolerate them.

Regrettably there is evidence that the number of abortions is increasing, and the E.B.S. reported recently that the demand for abortion beds had risen by 40% in the past eight years. In 1964 over 5,800 such admissions were made, and although it is not suggested that all of these abortions were procured previous surveys of hospital admissions show that a relatively small percentage of abortions are spontaneous. I agree with your leader writer that gynaecologists are far from being doubtful about the law "in practice," since anyone who decides that termination of pregnancy is indicated in a particular instance is obliged to rely upon his own interpretation of the law. In the tenth edition of *Midwifery* by Ten Teachers¹ it states categorically that every time a pregnancy is terminated a felonious act is committed. A recent questionnaire on abortion sent to 2,237 London doctors elicited 751 replies. Half the doctors replying said that the present law made it difficult for them to know when abortion is legal.

Your leader writer fails to point out that until 1803 it was not a criminal offence in