It was unnecessary for Professor Conn to select cases of primary aldosteronism from our published results,⁶⁻⁸ since we had reported these conclusions in the same group of cases—in Professor Conn's presence—more than a year before.⁹ Professor Milne's case was not one of these. Our recent paper⁹ was a progress report on a patient described in some detail earlier.⁶ We said of this case in 1963 that the association of low plasma-renin concentration and increased aldosterone secretion suggested an aldosterone-producing adrenal tumour, and that we were studying the effects of spironolactone treatment before removing the tumour.⁶ It thus seems obtuse of Professor Conn to misconstrue² the intention of our paper⁹ as "final proof" of his more recent "thesis."

We should further make it clear that neither we nor Professor Conn and his colleagues can rightly claim priority for the prediction of low plasma renin in primary aldosteronism, or for the removal of a tumour from a patient investigated in this way. Any such credit belongs to others.^{10, 11}—We are, etc.,

J. J. BROWN.
D. L. DAVIES.
A. F. LEVER.
W. S. PEART.
J. I. S. ROBERTSON.

St. Mary's Hospital,
London W.2.

REFERENCES

- 1 Brown, J. J., Davies, D. L., Lever, A. F., and Robertson, J. I. S., *J. Amer. med. Ass.*, 1965, **191**, 867.
- 2 Conn, J. W., *Brit. med. J.*, 1965, **1**, 656.
- 3 Boucher, R., Veyrat, R., de Champlain, J., and Genest, J., *Canad. med. Ass. J.*, 1964, **90**, 194.
- 4 Conn, J. W., Cohen, E. L., and Rovner, D. R., *J. Amer. med. Ass.*, 1964, **190**, 223.
- 5 Veyrat, R., de Champlain, J., Boucher, R., and Genest, J., *Canad. med. Ass. J.*, 1964, **90**, 215.
- 6 Brown, J. J., Davies, D. L., Lever, A. F., and Robertson, J. I. S., in *Boerhaave Course Hypertension*, edited by J. de Graeff, 1963, p. 44. Leyden University.
- 7 ———, *Canad. med. Ass. J.*, 1964, **90**, 201.
- 8 ———, in *Aldosterone*, a symposium organized by C.I.O.M.S., edited by E. Baulieu and F. Robel, p. 453, 1964.
- 9 ———, Peart, W. S., and Robertson, J. I. S., *Brit. med. J.*, 1964, **2**, 1636.
- 10 Laragh, J. H., Cannon, P. J., and Ames, R. P., *Ann. intern. Med.*, 1963, **59**, 117.
- 11 Kirkendall, W. M., Fitz, A., and Armstrong, M. L., *Dis. Chest*, 1964, **45**, 337.

Urinary Infection in Childhood

SIR,—We would like to add our support to the comments of Dr. H. Braude, Professor J. O. Forfar, and Dr. J. C. Gould (28 November, p. 1393) on the paper by Dr. Jean M. Smellie and her collaborators (14 November, p. 1222). We also have found that growths of potential pathogens exceeding 10^5 organisms per ml. (loop/blood-agar surface count) in bag-urine specimens from babies are frequent in the absence of significant pyuria or other urinary abnormalities which would support the diagnosis of urinary infection.

Our colleagues from Edinburgh do not indicate how they are dealing with this problem, and we would like to describe two urinary investigations which we have found helpful in making a positive diagnosis of urinary infection in some babies.

The first is to do a cell-count in a white cell counting chamber on the unspun specimen.^{1, 2} In contrast to an estimation of the number of cells per high-power field in the centrifuged deposit this method eliminates the

main technical variables, although not that due to different rates of urine secretion. A count of 200 leucocytes per c.mm. or more in a clean specimen is good evidence of urinary infection at any age. Also a count of five or more red cells per c.mm. is abnormal and warrants an examination of further specimens for more definite evidence of urinary infection or other urinary-tract abnormality.

The second method we have been using over a shorter period is to dilute the specimen 1 in 100 and spread two separate loopfuls (each delivering 2 c.mm. of urine) on half-plates of blood agar. In this way an estimate of bacterial counts in the range of 10^5 to 10^7 per ml. is obtained. Our present data suggest that counts of 5×10^6 per ml. or over indicate urinary-tract infection even in bag-urine specimens.

Finally, when the urinary findings are equivocal the value of examining repeated specimens before starting chemotherapy cannot be overstressed. We consider that in spite of the difficulties of diagnosis in babies catheterization should continue to be avoided because of the danger of introducing infection in this age group as well as at other ages.—We are, etc.,

PETER L. MASTERS.
IAN C. LEWIS.

Princess Margaret Hospital,
Perth, Western Australia.

REFERENCES

- 1 Masters, P. L., *Guy's Hosp. Rep.*, 1953, **102**, 76.
- 2 Stansfeld, J. M., and Webb, J. K. G., *Arch. Dis. Childh.*, 1953, **28**, 386.

Asymptomatic and Significant Bacteriuria

SIR,—Kass¹ uses the expression "asymptomatic bacteriuria" in respect of those patients with bacteriuria who "were not regarded by those responsible for their care as having active infection of the urinary tract." Kass was working with out-patients of the Boston City Hospital. Other authors use other equally local and subjective criteria, if indeed they attempt to define the condition at all. Clarity would be gained if authors using the term asymptomatic bacteriuria were to state which of the local and general symptoms and signs of inflammation of the urinary tract were expressly inquired after and found to be absent. As a basis for discussion may I suggest that the following should be looked for: pain on micturition, frequency, urgency, double micturition, dyspareunia, hypogastric pain or tenderness, renal pain or tenderness (elicited by percussion over the renal angle), abdominal bloating, pallor (common), a red, glossy complexion (less common), frontal headache, depression, debility, insomnia, epigastric discomfort, nausea, bad-smelling urine, and an ammoniacal smell to the patient. In children other symptoms are important—for instance, anorexia, bouts of vomiting, constipation, failure to gain weight, irritability, and screaming attacks.^{2, 3}

Significant bacteriuria is usually deemed to be present when there are 100,000 or more organisms per ml. of urine. The significance of this bacteriuria is a bacteriological one, and, while it is perfectly legitimate for bacteriologists to isolate a group of patients

with this urinary feature for special investigation, clinicians should not accept significant bacteriuria as being synonymous with inflammation of the urinary tract. Many (possibly most) patients presenting with inflammation of the urinary tract do not have significant bacteriuria, though they are not otherwise clearly distinguishable from those which have. For the clinician, bacteriuria is something of a red herring, and to him inflammation of the urinary tract is a disease of urinary tract tissues, which may or may not be accompanied by loss of organisms in the urine. We do not know if the presence or absence of bacteriuria implies a difference in kind between these two classes, or whether they are to be regarded as "open" and "closed" cases of urinary tract infection similar in principle to "open" and "closed" cases of pulmonary tuberculosis.—I am, etc.,

Lowestoft, Suffolk.

N. B. EASTWOOD.

REFERENCES

- 1 Kass, E. H., *Trans. Ass. Amer. Physns*, 1956, **69**, 56.
- 2 Stansfeld, J. M., *Proc. roy. Soc. Med.*, 1954, **47**, 631.
- 3 Smellie, J. M., Hodson, C. J., Edwards, D., and Normand, I. C. S., *Brit. med. J.*, 1964, **2**, 1222.

Alcohol in the Driver

SIR,—In an editorial (27 February, p. 540) you refer to the B.M.A. Council's approval of the B.M.A. Alcohol and Road Accidents Committee recommendation "making it an offence for a person with a blood-alcohol concentration in excess of 80 mg./100 ml. to drive a motor vehicle on the public highway."

There are two threshold levels of quite a different order as regards performance impaired by ethanol, reasonable allowance being made for dose-response variations. The first threshold, the higher one, indicates the level above which absence of impairment is unthinkable in the majority of subjects. The second threshold indicates the level above which the safety of road users would be measurably reduced. The upper threshold is often clearly associated with drunkenness, which in turn is associated with serious social and moral implications. (However uncertain pharmacologists are as regards the meaning of drunkenness, there is every certainty that, used or misused as a label, it may be most damaging.) On the other hand, the lower threshold is not, and should not be, associated with drunkenness in normal circumstances. I think it should be stressed that the recognition of two sharply differentiated offences,¹ one comparable to a serious moral offence and the other to a speed-limit infringement, is essential. Apart from the desirability of grading penalties to fit crimes, the reputations of drivers as well as the safety of road-users are involved.

A physiological threshold is something which varies. But the concept is real and invaluable. Pain, convulsion, and head-drop thresholds therefore vary. But the variation has its measurable limits to a standard range and an ED50 (median effective dose) level of drug can be established. Readers will accept without question the implication of the ED50 of drugs in general, but allow variation in the response to ethanol to fog their appreciation of ethanol thresholds.

The maximum blood-alcohol level consistent with the safety of other road-users has

been stated by the B.M.A. Special Committee to be 50 mg./100 ml.² In 1960 the Association was concerned in a statement that its relevant Committee "cannot conceive of any circumstances in which it could be considered safe for a person to drive a motor vehicle on the public roads, with an amount of alcohol in the blood greater than 150 mg./100 ml." Above this level was considered (in 1936) by committees of the American Medical Association³ and the U.S. National Safety Council⁴ to be *prima facie* evidence of being "under the influence." Here, then, are figures suggested for two important thresholds. But what threshold does 80 mg./100 ml. suggest? Is it a muddled compromise between 50 mg. and 150 mg.? One could see 80 mg. more effective in destroying reputations than saving lives.

Failure to appreciate the statistical basis of physiological thresholds, which are in turn the basis of so much legislation in regard to all sorts of activities, robs blood-alcohol levels of much of their meaning. The man who claims that speed limits should not be imposed because A drives at 40 m.p.h. less dangerously than B at 30 m.p.h. has the mentality of the man who cannot see why the carrying of firearms should be illegal in peacetime.

It should be clearly understood that, as regards what might be termed the high-threshold offence, the fact that moral and social considerations seriously enter the picture makes it important not to rely on chemical testing alone. With the low-threshold offence the chemical test would offer the sole criterion for conviction. It is difficult to see a satisfactory role being played by chemical testing without recognition of two offences, viewed as differently as the unlawful carrying of firearms and murder.

Considerations of space make it difficult to state the argument in less dogmatic terms.—I am, etc.,

PAUL J. CANNON.

Pharmacology Dept.,
University College,
Dublin.

REFERENCES

- 1 Cannon, P. J., and Pennington, G. W., *J. Irish med. Ass.*, 1962, 50, 137.
- 2 *Relation of Alcohol to Road Accidents*, p. 7, 1960. B.M.A., London.
- 3 A.M.A. Preliminary Report of Committee to Study Problems of Motor Vehicle Accidents. *J. Amer. med. Ass.*, 1937, 108, 2137.
- 4 *Ibid.*, 1939, 112, 2164.

Integration of Medical Schools

SIR,—Professor W. C. W. Nixon's letter (26 December, p. 1657) came as a surprise. Isolationism between medical schools does not exist to any extent in the U.S.A. At a recent conference in Massachusetts General Hospital the chairman of one of the departments remarked on the great co-operation and good will that exists between the many hospitals associated with Harvard Medical School. He became departmental chief a few years ago, after many years of training and teaching at a medical school on the west coast. What a privilege it was, he said, to work where the love of learning was a close second to the care of the patient.

At the many conferences at Boston hospitals, physicians from any hospital—civilian, Army, or Navy—are welcome. Whenever there

is a well-known medical personality in town or some eminent Bostonian is a guest speaker, all the hospitals with a training programme in Boston are informed and their staff are invited.

But this is not just confined to Boston. It is very typical of American Medicine. It would be a pity if the medical schools associated with the University of London did not make the most of their close proximity and eradicate their isolated policies, for isolationism is negative.—I am, etc.,

Boston, Mass.,
U.S.A.

G. J. O'KELLY.

Classification of Drugs

SIR,—During the past six months the Secretariat of the Committee on Safety of Drugs have developed, and are now using, a simple code for classification of drugs, as an integral part of their Register of Adverse Reactions to Drugs.

This code is based on the therapeutic uses of various drugs, but, to a limited degree, also takes account of pharmacological properties and chemical structure. It is not intended to be a substitute for the complex pharmacological and chemical codes which are being developed for research purposes by the Committee.

The Committee would like to draw attention to the existence of this code and is prepared to supply copies to interested persons. Any correspondence arising from this letter should be addressed to the Medical Assessor, Committee on Safety of Drugs, Queen Anne's Mansions, Queen Anne's Gate, London S.W.1.—I am, etc.,

D. A. CAHAL,
Medical Assessor,
Committee on Safety of Drugs.
London S.W.1.

P.H.L.S. Committee on Hospital Disinfectants

SIR,—On 13 February 1965 you published a report by this committee (p. 408), of which I am secretary. The recommendations for phenolic disinfectants for general use included clear soluble fluids of the Lysol type having a Chick-Martin coefficient of at least 1.5. There is also a reference to chlorinated phenols. This reference was intended to exclude those agents which consist entirely of such substances and which may have a narrow antimicrobial spectrum and an undue sensitivity to organic matter. It was not intended to exclude those fluids which may merely contain some chlorinated phenols. Clear soluble fluids are in fact very suitable for general use in hospitals, but no standard specification—or indeed agreed definition—for such products exists. In order to avoid any confusion it may be helpful to list the fluids of this type known by me to be at present available, and which meet the requirements of the committee:

Hycolin—Wm. Pearson Ltd.

Printol }
Sudol } Printar Industries Ltd.

—I am, etc.,

Central Public Health
Laboratory,
London N.W.9.

J. C. KELSEY.

Drug Advertisements

SIR,—A careful perusal of your issue of 27 February fails to reveal even one advertisement that would fit Professor Owen L. Wade's blanket indictment in that issue, p. 585. Many of these advertisements, indeed, are models of restraint, giving little beyond factual information, immediately useful to the doctor.

Constructive criticism should always be welcomed, but surely Professor Wade weakens his argument by overshooting the mark and introducing elements of bias. Tranquillizers, for example, have without question had a profoundly beneficial effect in the treatment of disturbed patients beyond the reach of other drugs, and not all of Professor Wade's colleagues will agree they are no better than barbiturates, which, valuable as they are, have often come under attack. Again, accepting the need for restraint in giving antibiotics, to insist upon the guidance of a bacteriologist in every case would mean to withhold these life-saving drugs unreasonably in many situations where their prompt administration can do the most good.

It is not necessary to defend every advertisement and every piece of promotional literature on behalf of drugs. The remarkable fact is that so very few of them offend, while the great majority perform the indispensable function of keeping the doctor abreast of a multitude of important developments on which he could inform himself with equal promptitude and accuracy in no other way—a service which most doctors fully appreciate.

Like all to whom advertising is directed, Professor Wade and whoever of his fellows may be displeased have a simple remedy. They are under no obligation to read the promotional material prepared for them with such care and, for the most part, responsibility. With respect, however, they will be better doctors for giving it the attention it merits.—I am, etc.,

STEWART M. RITCHIE,
Managing Director,
Intercon International Marketing Ltd.
London W.1.

The Scientific Method

SIR,—I regret I do not find Dr. A. K. Clarke's views on the nature of truth (20 February, p. 526) convincing. My point was not to examine why the circulation of the blood was not discovered before Harvey's time, but to assert that Harvey did in fact discover something the truth of which every investigation since then has confirmed. Certain views may be mistakenly accepted as true (such as that of the blood ebbing and flowing) and hamper advance, but the possibility that people can sometimes be wrong should not be made to prove that they can never be right. I do not suggest we should ever accept any statement on authority, but if some piece of knowledge fits in with the evidence of the senses and the rules of logic and is confirmed by every possible test that man can devise then it is reasonable to accept it as true, constituting a truth as absolute as can be attained. That, at least, is what I understand by absolute truth; scientists (from *scire*—people who do know or seek to know what is or could be the truth) certainly proceed on the assumption that certain truths are true