Aetiology of Multiple Sclerosis

SIR,—By using a new method, recently referred to in your columns¹ by Sir Macfarlane Burnet, we find that suspected idiopathic autoimmune disorders can be classified by statistical criteria into two categories. This division also corresponds, interestingly enough, to a fundamental anatomical distinction regarding the location of the target tissue.

In both categories it appears^{2,3} that the disease process is initiated in genetic carriers by a small number (from 1 to, perhaps, at least 12) of specific random events, the average rate of which is effectively constant in a given genotype, and in normal environments throughout post-natal life. Following Burnet⁴ we have suggested that these "random initiating events" are a form of somatic gene mutation in lymphoid stem cells, and that they trigger-off the growth of "forbidden clones." When such a disease affects both sexes we find the average rate of a specific initiating event in carrier females is either the same as or double that in carrier males, and we have so far found no significant departures from this "rule." An interval or "latent period" intervenes between the last random initiating event and the onset of symptoms or signs. It is here that the statistical distinction between the two classes of spontaneous autoimmunity is manifested.

The duration of the latent period is evidently governed, in part, by an endogenous defence mechanism directed against forbidden clones, and this is vulnerable to certain infective agents, certain drugs, and to mental stress.^{5,6-7} In one class of autoimmunity (inflammatory polyarthritis and systemic lupus erythematosus are classical examples), where the primary pathogen is probably a cell-bound (lymphocytic) autoantibody, defence is twice as efficient, on the average, in females as in males. For a given environment the average latent period in females is twice as long as in males.⁸ Defence is probably mediated through immunoglobulins,^{2,6-7} and in this class of disease gross abnormalities in these plasma proteins (for example, hypergammaglobulinaemia, rheumatoid factor, anti-nuclear factor) are common.

In the other class of suspected autoimmunity (multiple sclerosis, some peripheral neuropathies, and psoriasis are typical examples) the sex- and age-distributions show that the average latent period, in a given environment, is the same in males and females. We conclude that there is no sex-differential in the efficiency of the related endogenous defence mechanism. Furthermore, the target tissues (for example, central and peripheral nervous systems, and epidermis) are located behind a blood-tissue barrier and they are not normally infiltrated by small lymphocytes. There are various indications⁹ that the primary pathogen, or autoantibody, in these diseases is humoral (in some instances, at least, it appears to migrate with the α_2 -globulin fraction) and that it can diffuse through blood-tissue barriers. Disturbances in the immunoglobulins are generally inconspicuous in this class of disease—as Wright *et al.*⁸ confirm.

We therefore propose, in contradiction with your leader (20 February, p. 466), that multiple sclerosis may prove to be a

disturbed-tolerance autoimmune disease in which the primary pathogen is a humoral (non-immunoglobulin) autoantibody.—We are, etc.,

P. R. J. BURCH.
R. G. BURWELL.
N. R. ROWELL.

The University of Leeds,
The General Infirmary,
Leeds 1.

REFERENCES

- 1 Burnet, F. M., *Brit. med. J.*, 1965, 1, 338.
- 2 Burch, P. R. J., *Lancet*, 1963, 1, 1253.
- 3 Burch, P. R. J., *Brit. J. Psychiat.*, 1964, 110, 825.
- 4 Burnet, F. M., *The Clonal Selection Theory of Acquired Immunity*, 1959. Cambridge University Press, London.
- 5 Burch, P. R. J., and Rowell, N. R., *Lancet*, 1963, 2, 507.
- 6 ———, *Amer. J. Med.*, 1965, in press.
- 7 ——— and Burwell, R. G., *Quart. Rev. Biol.*, 1965, in press.
- 8 Wright, R., Morton, J. A., and Taylor, K. B., *Brit. med. J.*, 1965, 1, 491.

Oral Contraceptives and Liver Damage

SIR,—The communication on this subject by Dr. Antti Eisalo and his colleagues (15 August, p. 426) reports abnormal liver-function tests in every member of a small group of post-menopausal patients receiving 1–2 tablets of Lyndiol (5 mg. 17 α -ethynyl-oestrenol and 0.15 mg. 3-methoxy-17 α -ethynyl-oestradiol) daily. Similar changes have been noted by Drs. I. P. Palva and O. O. Mustala (12 September, p. 688) in post-menopausal patients receiving one tablet daily of Anovlar (4 mg. of norethisterone acetate and 0.05 mg. of ethinyloestradiol), which contains a different progestogen and a different oestrogen. On the other hand, Dr. L. I. Swaab (19 September, p. 755), Dr. G. Linthorst (10 October, p. 920), Dr. E. Rice-Wray (17 October, p. 1011), and Dr. E. T. Tyler (14 November, p. 1264) emphasize that large-scale, long-term investigations of premenopausal patients given Lyndiol or Conovid (5 mg. norethynodrel and 0.075 mg. mestranol) for contraception showed no evidence of liver dysfunction. We agree with these authors that Dr. Eisalo's report may reflect a different metabolism for these compounds in post-menopausal as opposed to premenopausal patients. We would, however, like to record our experience apropos of Dr. Rice-Wray's comment—"the evidence is insufficient to label this as a hepatotoxic effect," and Dr. Tyler's comment—"laboratory tests, *per se*, are not enough to prove a clinical diagnosis."

Since Dr. Eisalo's communication we have carried out routine liver-function studies and also liver biopsy in four post-menopausal patients receiving six tablets of Lyndiol daily in the treatment of breast cancer. In all cases there was a rise in serum glutamic oxalacetic transaminase and serum isocitric dehydrogenase levels well beyond normal, the changes in liver-function tests appearing after two weeks' drug administration. The serum glutamic oxalacetic transaminase level was above 200 units (Reitman-Frankel method; normal range 10–40 units) in all cases, the highest value being 450 units. The serum isocitric dehydrogenase values at this stage ranged from 77–165 units (normal range 3–10 units).¹ In addition, clinical jaundice was

noted in two patients associated with a serum-bilirubin level of up to 4.1 mg./100 ml. The alkaline-phosphatase and serum-protein levels remained normal in all cases.

Liver biopsy in all four patients showed parenchymal cell necrosis which was strictly limited to the centrilobular zones with accumulation of lipofuscin. In the two with raised serum bilirubin these areas showed also cholestasis in the hepatic cells and biliary canaliculi. In two cases where therapy was stopped after two months the abnormal levels in the liver-function tests fell to normal within two weeks of stopping in one case, but in the other, although reduced, have persisted for six weeks. In two cases where Lyndiol was continued beyond two months (currently six months in one case) the abnormal levels fell spontaneously while under treatment but did not return to normal.

We note that Dr. Eisalo found that the progestogen component of the drug (Lynestrenol or 17 β -hydroxy-17 α -ethynyl-oestrenol) given alone causes no changes in liver function whereas the oestrogen component (Mestranol or 3-methoxy-17 α -ethynyl-oestradiol) did so in one out of three cases. This is an interesting point, as Sherlock² has pointed out that steroids with an alkylated group at C17 are liable to induce some hepatic damage and cholestasis when given in sufficient dosage for prolonged periods. Both the progestogen and the oestrogen components of our drug possess this configuration.

Our experience therefore confirms Dr. Eisalo's findings, and in addition we have shown, by liver biopsy, the presence of hepatic impairment in four cases. We emphasize again that our patients were post-menopausal and were receiving six tablets daily.—We are, etc.,

BASIL A. STOLL.
J. T. ANDREWS.
R. MOTTERAM.
J. UPFILL.

Peter MacCallum Clinic,
Melbourne C.1, Australia.

REFERENCES

- 1 Bell, J. L., and Baron, D. N., *Clin. Chim. Acta*, 1960, 5, 740.
- 2 Sherlock, S., *Diseases of Liver and Biliary System*, 1963, 3rd ed., p. 307. Blackwell, Oxford.

Somatic Mutation

SIR,—In the second paragraph of his paper "Somatic Mutation and Chronic Disease" (6 February, p. 338) Sir Macfarlane Burnet writes, "For many years it has been recognized that . . . (for) . . . several types of cancer . . . if both age-specific incidence and age were plotted logarithmically the points fell on a straight line. This can be interpreted *only* on the assumption that the condition is determined by the random occurrence of events occurring with equal likelihood over the time-span being considered" (our italics). This, in our opinion, is not true.

The demonstration by Nordling¹ and Armitage and Doll² that the "random occurrence of events" hypothesis produces the required age-specific incidence pattern may be generalized as follows: Suppose that the probability distribution function of the age