Current Practice

CHILD CARE IN GENERAL PRACTICE

Trauma in Childhood

ROBERT ROAF,* M.CH., F.R.C.S.

Trauma in the neonatal and intranatal period will not be considered in this article. It should, however, be noted that the results of neonatal trauma are not always immediately apparent, and indeed some conditions—for example, spasticity and Erb's palsy—are often recognized only at a later date when they manifest themselves as chronic orthopaedic conditions. Once the newborn child has survived the trauma of birth one would expect him to be relatively immune to trauma until he reaches the toddling stage, but two aspects of modern life have altered this state of affairs—the development of the internal combustion engine and the increasing recognition over the last few years of the "battered baby" syndrome.

Mechanical Injuries

The babe in arms, whether as a passenger or as a "pedestrian," is liable to be quite badly injured in accidents involving motor-cars. Such trauma does not follow any definite pattern, but like other severe direct injuries may lead to very serious consequences in the young babe.

The recognition of fractures and dislocations in young babies by radiography may be difficult, and if the mother and child are brought into hospital either as a result of being knocked down by another car, or because the car in which they were passengers was involved in an accident, it is essential to examine the baby very carefully, particularly looking for evidence of gross swelling or haematoma formation, as this may well indicate serious underlying damage. It is important to realize that injuries and haematoma formation in small children may closely mimic bacterial infection. Locally there is a hot, red, painful fluctuant swelling; the baby is pyrexial and fretful, appetite is impaired, and there is a leucocytosis. In addition, a haematoma acts as an ideal culture medium and may become secondarily infected. Any hot, red swelling in a child should be aspirated under aseptic conditions. I have seen a swelling in a haemophilic boy closely resemble an abscess, so it is wise to be sure before using a knife. The injection of chymotrypsin into the haematoma after aspiration may aid in resolution.

In these days of increasing mechanization on the roads, in the home, and on the farm serious mechanical injuries are increasingly common. These do not follow any set pattern and each case must be treated on its merits. However, compared with adults, the powers of restoration in children are amazing. This means that these lesions must be treated with the utmost conservatism—for instance, a guillotine amputation of the index finger in an adult would need trimming of bone and formal skin closure, but the same injury in a child could be safely treated with a sterile dressing, when the raw area will become epithelized, thus preserving the maximum possible length.

Battered Baby Syndrome

The battered baby syndrome has become more commonly recognized in the last few years (Griffiths and Moynihan, 1963).

The child is usually teething and is often brought to hospital by the parents with one or more grossly swollen limbs, or perhaps a grossly swollen face, with evidence of haematoma formation. The parents deny that it could have been injured, and it is common to suspect some serious condition—for instance, a blood dyscrasia, infection, or even tumour formation. In the early stages radiographs are not always informative, but those taken a few days later will often show subperiosteal formation of new bone at the ends of the long bones (Fig. 1) and the syndrome is characterized by fracture separation of non-ossified cartilaginous epiphyses. Usually, but not invariably, these children suffer from a certain amount of malnutrition. By now a number of these patients have been described and it is recognized that the syndrome is nearly always due to injury inflicted by the baby's own parents or by baby-watchers. The condition was first clearly described and made widely known by Caffey (1957), but it must be differentiated from the other syndrome described by Caffey and Silverman (1945)—namely, "infantile cortical hyperostosis," in which there is swelling of the jaws and often of the scapulae and long bones, and in which subperiosteal new bone formation is also increased. Alternatively, the child may present with cerebral symptoms—vomiting and "inexplicable" coma—due to an intracranial haematoma. If undiagnosed, a cranial injury from parental assault can prove fatal, and in the absence of a history of injury diagnosis may be very hard.

FIG. 1.—Bone changes at upper end of humerus in battered baby syndrome.

Unless one is aware of this syndrome it may be difficult to recognize it, and even when it is suspected it is not always easy to prove that trauma is the only cause; indeed, it is almost mandatory that the child should be admitted to hospital and fully investigated, and other causes excluded. When the condition

* Professor of Orthopaedic Surgery, University of Liverpool.

has been diagnosed it is extremely hard to know what is the right course of action to take in the interests of the child, for it is not unusual, when the child's condition has improved and he has been sent home, for the parents to assault him again.

The treatment of the physical injuries does not usually present any serious problem, but preventing the recurrence of the assaults is a far more complex procedure. It is essential to have full consultation between the parents' own general practitioner, the medical officer of health, and the various welfare services. Probably in the long run the parents' own doctor is in the best position to advise about the best course of action to take in the ultimate interests of the child.

Common Fractures

Greenstick Fracture.—At a slightly older age, when the child is a toddler, he is likely to fall, and at this age the most frequent injury is a greenstick fracture of both bones of the forearm, or a spiral fracture of the tibia or femur. Treatment is usually extremely simple, and, even if the fracture is not fully reduced, the powers of growth and remodelling in the small child are

FIG. 2a

FIG. 2b

FIG. 2.—Remodelling of supracondylar fracture: (a) radiograph after reduction; (b) two years later.

FIG. 3a

FIG. 3b

FIG. 3c

FIG. 3.—Fracture of capitellum: (a) before reduction; (b) after reduction; (c) untreated fractured capitellum.

very considerable. Within a year or two even 30° of angulation or 1 in. of shortening will be corrected by nature alone (Figs. 2a and b).

Elbow Fractures

Pulled Elbow.—Another common syndrome at this age is a "pulled elbow." The history is usually that the child has been lifted by one hand, after which the elbow becomes painful and the child is unable to rotate his forearm. Radiographs do not show any abnormality. It is usually believed that this syndrome is due to the head of the radius being pulled down into the orbicular ligament and becoming impacted in this position. A push on the head of the radius associated with a pronation-supination movement will usually cure the condition.

Fracture of Capitellum.—When the child is slightly older the most common elbow injury is a fracture of the capitellum after a fall on the outstretched hand (Figs. 3a and b). The elbow becomes swollen and a radiograph reveals considerable outward displacement and rotation of the capitellar epiphysis. It is extremely important to recognize this injury as, if the capitellum is left unreduced, non-union, failure of growth, and gross deformity will result (Fig. 3). It is usually impossible to reduce the capitellum satisfactorily by conservative means, and open operation and fixation are required—either with a Kirschner wire or a screw, which require removal later, or a catgut suture—to hold the capitellum in perfect alignment. This will ultimately restore the normality of the elbow.

Supracondylar Fracture of Elbow.—Another important injury of the elbow in children is a supracondylar fracture of the elbow. As with so many injuries its real importance lies not in the fracture itself but in the complications which go with it (Fig. 4). The most important of these is Volkmann's ischaemic contracture. As a result of a fall on the outstretched hand the elbow becomes swollen and the classical signs of impaired circulation appear. Of these the first and most important is pain in the hand aggravated by passive extension of the fingers; the child is also unable to extend the fingers voluntarily. At a later stage there will be absence of the radial pulse, impaired peripheral capillary circulation, and either pallor or cyanosis of the hand with swelling of the fingers. Later still blistering and complete paralysis of the forearm muscles appear. However, the condition should be suspected long before these signs are apparent.

The treatment of a threatened Volkmann's ischaemic contracture is a matter of some controversy. It is of course important that all splints and bandages should be removed and the child's general blood-pressure maintained. With the help of these measures the collateral circulation will very often be adequate to maintain a good blood supply to the muscles of the forearm and hand even though the brachial artery has been occluded by direct pressure or spasm. However, if, after having elevated the arm, released all constricting bandages, and placed the elbow in a semi-extended position, there is still evidence of impaired circulation it will usually be wise to perform an operation. Two main types of operation have been described. The first method, associated with the name of Professor George Perkins, is to approach the bone fragments posteriorly and replace them by direct open reduction so that the lower fragment, instead of lying posterior to the upper fragment, lies in line with it, or even in front. The other operative approach is to expose the brachial artery anteriorly and if it contains a

thrombus or blood clot then this should be removed. If the arterial wall is hopelessly damaged it is probably best to resect the affected portion and ligate the artery above and below it.

FIG. 4.—Nipping of brachial artery in supracondylar fracture of elbow.

Owing to the small size of the artery in children it is only rarely possible to do an excision and direct suture of a damaged segment of the artery or to perform a replacement graft.

It will be clear that even after heroic measures such as these function may not be completely restored. Obviously far more important than the ability to undertake major and complicated surgery is that of recognizing the threat of Volkmann's ischaemia and taking immediate steps to prevent its occurrence. If these measures are taken early enough it is rare for the child to develop serious contracture at a later date. This is undoubtedly the most serious complication of closed fractures of the limbs in children.

Monteggia Fracture.—Another serious fracture occurring around the elbow in children is the so-called Monteggia fracture—that is, dislocation of the head of the radius associated with a fracture of the ulna. In this condition there is usually tearing of the interosseous and orbicular ligaments and widespread separation of the radius and ulna. Though it may be possible to reduce these injuries by conservative means, it is usually wise to operate. The head of the radius is reduced and fixed in place by a reconstruction of the orbicular ligament. Internal splints are then applied to the ulna by means of either a plate or an intramedullary nail.

Naturally in any fracture or injury around the elbow there may be damage to one of the surrounding nerves. Of these the median nerve is the most important, though the radial nerve is the most commonly damaged. Again it is of the utmost importance to recognize such a complication at the earliest possible date and point out to the parents that it is caused by the injury and not by the treatment.

Most other fractures in children have a strong tendency to unite, and even if there is some mal-union growth usually restores normality; the nearer the fracture is to the growing end of the bone the more quickly this natural correction occurs. The opposite occurs if there has been damage to the epiphysis; under these circumstances progressive deformity and shortening will result. The epiphysis may be damaged in early childhood—usually from infection or at a later stage from injury—for example, a fall on the outstretched hand may damage the lower radial epiphysis and produce a Madelung's deformity, or damage to the trochlear epiphysis often produces progressive cubitus varus or "gun-stock elbow."

Ankle Injury

At an older age—around 12 or 13—an inversion injury to the ankle may damage the lower tibial epiphysis; as a result there is progressive varus deformity of the ankle due to failure of growth of the medial part of the lower tibial epiphysis, and continued growth of the fibula.

General Considerations

The characteristics of a child's reaction to injury are his amazing powers of repair and restoration, both of function and appearance, even when a fracture has been imperfectly reduced. It requires considerable judgment and experience to know when a given position is "acceptable" or when strenuous efforts should be made to improve the position, and the surgeon must always bear in mind that most injuries in childhood have a favourable outcome provided that they are not over-treated. This applies particularly to elbow injuries, where excessive manipulation may delay recovery and cause serious complications such as Volkmann's ischaemia. In fractures of the shafts of long bones in children open operation and internal fixation are rarely necessary.

A possible source of diagnostic confusion in children is that the child may have been suffering from some constitutional disturbance before the injury—for instance, classical rickets is not uncommon among the immigrant population in big urban areas. It may also be necessary to exclude renal rickets and other conditions such as fragilitas ossium, especially in cases of multiple fractures. In this connexion a greenstick fracture of both bones of the forearm in children of 5 or 6 presents a special problem. For some ill-understood reason re-fracture is not uncommon and the parents frequently suspect either that the child has some underlying bone disease or that the fracture has not been treated properly. It is therefore wise to warn them of this possibility when treating a fracture of the forearm.

Nerve, Tendon, and Blood-vessel Injuries

With the exceptions mentioned above closed injuries of children's limbs do not usually produce serious consequences. On the other hand, cuts—especially round the wrist—may lead to severe loss of blood or division of tendons and nerves. Provided these are promptly repaired the prognosis with regard to ultimate function is usually excellent in children, but immediate diagnosis is all-important.

Sensory testing is exceedingly difficult and often unreliable in children—even a complete division of the median nerve in a child may not be easily detected by sensory testing alone, as the area of autonomous distribution is relatively small. If there is any possibility of a nerve or tendon being damaged and if there is an open wound at one of the important anatomical sites—for example, the wrist or round the elbow—it is better to explore the wound and examine the tendon and nerves rather than fail to diagnose. Primary repair of severed tendons is usually desirable, but, unless the wound is exceptionally clean, nerves should usually be sutured six weeks later when the skin has healed.

Burns

Burns in children are exceptionally serious, not only from the point of view of producing deformity but also as a danger to life, and in this respect the skin loss due to avulsion of skin or crushing is very similar to that due to burns. The management of extensive burns or other forms of skin loss is of course a complicated problem requiring the services of a plastic surgeon, but it is all-important in the early stage that the area should be protected from infection by applying a large sterile dressing and that the child should be treated for shock—if necessary by intravenous therapy—as soon as possible.

REFERENCES

- Caffey, J. (1957). *Brit. J. Radiol.*, **30**, 225.
— and Silverman, W. A. (1945). *Amer. J. Roentgenol.*, **54**, 1.
Griffiths, D. LL, and Moynihan, F. J. (1963). *Brit. med. J.*, **2**, 1558.