contain two amphetamine tablets (10 mg.) and two Parnate (tranylcypromine) tablets (10 mg.).

Her temperature continued to rise to 109.4° F. (43° C.). She was treated by immersion in ice water and the infusion of intravenous salicylate. Therapy was instituted in order to correct the marked dehydration, acidosis, hyperkalaemia, and subsequent hypotension that she suffered. Her blood-pressure remained unobtainable in large doses of metaraminol, but did respond to L-norepinephrine. After about 20 hours of therapy the patient began to support her own blood-pressure, carry on normal respiration, and respond feebly to simple commands. During the second day in the hospital she rallied dramatically and by its end was recognizing and conversing with relatives. Once her sensorium was clear, the patient related a detailed story of her ingestion of amphetamines and tranylcypromine. She had apparently been using amphetamines intermittently for both diet and mood-elevating effects. A few days before she became ill she started to ingest 10 mg. tranylcypromine tablets once or twice a day. She had taken one amphetamine and one tranylcypromine tablet just before going on duty the night that she took ill. This was the first time that both drugs had been taken together. The patient had obtained the drugs as samples in the hospital clinics.

The patient recovered completely. Her mental status during convalescence was normal and she gave us no reason to doubt her story.

At the time of her admission extensive blood, urine, spinal-fluid, and stool cultures were negative. A lumbar puncture had been normal. A variety of drugs were looked for, including salicylates. None were found. Unfortunately no assay for amphetamine or tranylcypromine was done.

I believe that this case represents an example of near-fatal hyperpyrexia caused by the ingestion of therapeutic amounts of tranylcypromine and amphetamine. In the full appraisal of the clinical usefulness of antidepressant agents, such cases must be given proper consideration.—I am, etc.,

Johns Hopkins Hospital,
Baltimore,
Maryland, U.S.A.

EDMUND LEWIS.

REFERENCES

- Stanley, B., and Pal, N. R., *Brit. med. J.*, 1964, 2, 1011.
- Bowen, L. W., *ibid.*, 1964, 2, 1466.
- Lockett, M. F., and Milner, G., *ibid.*, 1965, 1, 921.

Oral Contraceptives and Liver Injury

SIR,—In your leading article (29 May, p. 1391) you refer to the enormous numbers of women who are now taking oral contraceptives, and, of course, very many of these women are taking them without medical indications. Those of us who are concerned about the possible long-term effects of this wholesale use of hormones will not be very greatly reassured by your article. It may be true that liver damage is the least of the risks (I think there are more important possibilities), or it may be that there is no danger at all, but surely the evidence is insufficient to warrant the conclusions you draw.

The investigations reported by Dr. G. I. M. Swyer and Dr. V. Little (29 May, p. 1412) are extremely valuable, but we cannot say the risk is negligible merely because twelve women have taken these pills for a few years without demonstrable liver damage. Would it not have been as logical to have argued, say ten years ago, that because twelve men had smoked heavily for three to six years and none had developed cancer, there was there-

fore no harm in smoking? Such a conclusion would plainly have been dangerously incorrect, but it would have had this in common with the other conclusion: it would have been in accordance with what people wanted to hear and to believe.—I am, etc.,

Banbury, Oxon.

J. M. GATE.

Action of Bismuth Aluminate

SIR,—The paper by Dr. S. G. Flavell, Matts and his colleagues (20 March, p. 753) demonstrates statistically significant clinical results in favour of bismuth aluminate. Although the exact mode of action of this new bismuth compound is not understood, the authors' hypothesis suggesting that the good therapeutic results may be due to a coating effect providing a local antipepsin action at the site of the ulcer is an interesting possibility.

Bismuth aluminate has been used in this department in following up our studies on the effect of aspirin on gastric secretion.¹ Our initial objective was to determine whether bismuth aluminate would provide protection against gastric mucosal damage caused by the administration of aspirin. Our results in this respect were negative, but some of our experimental findings may have a bearing on the hypothesis put forward by Matts and his colleagues.

In one series of our cat experiments the animals were given either bismuth aluminate or aspirin, or a combination of the two drugs, prior to histamine infusion and tests for gastric function. The experimental technique was basically the same as that reported in our paper on histamine secretory tests in cats.² The bismuth aluminate or aspirin was suspended in about 50 ml. of water and injected into the stomach. After 1½ hours the stomach was opened and all residue of the drug was removed.

Intense histamine stimulation was started and the gastric juice collected for the next 3½ to 4 hours. At the conclusion of the experiment the stomach was opened and in 13 cats where bismuth aluminate had been administered alone a film of the bismuth compound was found to be clinging to the gastric mucosa even though the drug had apparently been previously removed from the stomach.

In another series of experiments gastric juice was obtained from dogs; peptic activity was measured and subsequently bismuth aluminate was added at a concentration of 0.05 g. per ml. Pepsin estimations were repeated at various pH levels and it was found that peptic activity was reduced by bismuth aluminate at all pH levels from 0.5 to 6.5. The gastric juice used in these experiments was obtained from denervated Heidenhain pouches and either because of this or because of species difference there was much less peptic activity than in man. This suggests that to demonstrate the antipeptic effect of bismuth aluminate on gastric juice from the lumen of the stomach in man a larger dose would be needed than that used by Matts and his colleagues. Nevertheless, our results provide some confirmation of the antipeptic action previously demonstrated *in vitro*.³

Thus it appears that bismuth aluminate could certainly provide a coating of the gastric mucosa, and this may exert a local antipeptic effect of a sufficient order to prevent auto-digestion at the site of the ulcer, even though it may not be detectable by biochemical analysis of a sample of gastric juice from the lumen. In this context it is of interest that surface mucus can have a pH of up to 9 or 10 while the pH of the stomach contents may be less than 5.

As Matts and his colleagues suggest, hyperacidity alone may be relatively unimportant in connexion with the origin of peptic ulceration which is almost certainly due to auto-digestion by acid-pepsin and possibly involving other gastric processes. Peptic inactivation is thus important, and if bismuth aluminate exerts a local antipeptolytic action at the site of the ulcer, following coating of the gastric mucosa, this could certainly account for the good clinical effect demonstrated in the double-blind trial.—I am, etc.,

St. Vincent's Hospital,
Sydney,
New South Wales, Australia.

G. W. MILTON.

REFERENCES

- Lynch, A., Shaw, H., and Milton, G. W., *Gut*, 1964, 5, 230.
- Milton, G. W., Skyring, A. K., and George, E. P., *Gastroenterology*, 1963, 44, 642.
- Bateson, P. R., *J. Pharm. Pharmacol.*, 1958, 10, 123.

Sodium Fluoride and Cell Growth

SIR,—Professor W. D. Armstrong and his colleagues (29 May, p. 1435) have now broadened the scope of their earlier observations (20 February, p. 486) which refuted the evidence of Berry and Trillwood¹ for the inhibition of cell growth *in vitro* by small amounts of sodium fluoride incorporated in the culture medium. Dr. Berry and Mr. Trillwood (20 March, p. 793) had disputed the validity of the earlier observations of Armstrong and his colleagues on the ground that their own data were based upon cultures extending over three- to four-cell duplications, which was a better test of growth inhibition than the one-cell duplication originally examined by Armstrong and his colleagues.

The more recent observations of Armstrong and his colleagues go some way toward answering these objections, although only a small number of selected HeLa-cell clones were studied. Nevertheless, since Berry and Trillwood had maintained that their original biological observation had still not been validly refuted, it seemed important to repeat their own experiments—i.e., growth tests of HeLa cells growing as a monolayer attached to glass for at least four cell duplications during a week. I have now performed five such experiments with cells cultured for over six cell duplications during a week and I am unable to confirm their observation of a growth inhibition by sodium fluoride.

Five growth experiments have been performed in this laboratory with HeLa cells

Number of HeLa Cells ($\times 10^6$) after 1 Week's Growth With and Without Sodium Fluoride (1 p.p.m.)

Experiment Number	Control Cultures	Fluoride Cultures
1	3.77 3.75	3.87 4.02
2	4.65 4.90	4.82 4.80
3	5.02 4.17	4.22 4.12
4	4.30 4.80	4.50 4.00
5	4.90 4.60	4.50 4.30
Mean	4.486	4.315

Difference between means = 0.171 ± 0.120 (4% $\pm$ 2.8%).