- 3 Lance JW, Smee RI. Partial seizures with visual disturbance treated by radiotherapy of cavernous hemangioma. *Ann Neurol* 1989;26:782-5.
- 4 Jampel RS. Convergence, divergence, pupillary reactions and accommodation of the eyes from faradic stimulation of the macaque brain. *J Comp Neurol* 1960;115:371-400.
- 5 Chiappinelli VA, Wolff KM, Feng C, Yum L, McMahon LL. Different responses to opioids measured in terminals and somas of Edinger-Westphal neurons. *Neuroscience* 1993;57:425-32.

Parkinson's disease and tumour in the supplementary motor area: a re-evaluation

In 1988 Straube and Sigel¹ reported on a 56 year old patient with a bilateral Parkinson's syndrome, including resting tremor, rigidity, bradykinesia, and a favourable response to levodopa medication, starting at the age of 51 years. This patient was discovered to have a tumour (low grade astrocytoma grade II-III) in the left hemisphere, mainly involving the supplementary motor area. Based on the low chance that both Parkinson's disease and astrocytoma occurred at the same time in this young patient (estimated probability 0.005%–0.1%), it was postulated that the tumour in the supplementary motor area induced the bilateral parkinsonian syndrome, although the levodopa responsiveness was unusual. Importantly, however, the supplementary motor area is considered to represent one of the critical motor areas which are establishing the motor circuit through the basal ganglia.^{2,3} We now report that this patient has idiopathic Parkinson's disease independently of astrocytoma grade II.

The patient is now 65 years of age. She has typical Parkinson's disease, featured by bradykinesia, rigidity, and resting tremor predominantly on the left side. She has responded favourably and continuously to levodopa over the past five years.

Her daily medication at the report of 1988 was 400 mg levodopa (plus decarboxylase inhibitor) and 12.5 mg bromocriptin. Her medication now consists of a combination therapy of 400 mg levodopa (plus decarboxy-

lase inhibitor) and 1 mg pergolide per day. Currently, she has developed biphasic motor fluctuations, and biphasic and peak of dose dyskinesia on both sides (predominantly left) and nocturnal akinesia. Testing with apomorphine (3 mg subcutaneously) disclosed a positive dopaminergic response according to the criteria previously published.^{4,5}

Her astrocytoma grade II was treated successfully with iodine-125 seeds in the tumour. She currently presents with a cyst in her supplementary motor area. Over the past two years there was regrowth of the tumour (anaplastic astrocytoma, grade WHO III). She was retreated with a course of γ rays and cortisone. Under this treatment, her parkinsonian syndrome (shuffling gait, resting tremor, on-off fluctuations, freezing episodes) deteriorated transiently for about 10 weeks. This deterioration seemed to distinctly exceed potential γ ray/cortisone induced side effects, as for instance manifested by fatigue and equilibrium disturbances. After cessation of the treatment, the parkinsonism gradually improved to the previous stage over a period of six months in combination with increased levodopa/dopamine agonist dosages.

Single photon emission computed tomography (SPECT) investigations performed with [123 I]IPT (N-(3-iodopropen-2-yl)-2 β -carbomethoxy-3 β -(4-chlorophenyl) tropane, a cocaine analogue with high affinity for the presynaptic dopamine transporter,^{6,7} displayed greatly reduced striatal binding of the radioligand. Compared with age matched controls specific [123 I]IPT binding was significantly reduced with more pronounced decrease of tracer accumulation in the putamen than in the caudate. Two SPECT scans were performed at an interval of 12 months. Both studies disclosed findings compatible with idiopathic Parkinson's disease (figure, table).⁸ Multiple system atrophy, progressive supranuclear palsy, and other atypical Parkinson's disease syndromes were excluded by clinical and MRI criteria. Additionally, an iodobenzamide (IBZM) SPECT showed normal, striatal D2 receptor binding, compatible with the diagnosis of idiopathic Parkinson's disease.⁹

In conclusion, the proposal of the previous report¹ that a tumour in the supplementary motor area may cause a parkinsonian syndrome is withdrawn. At present we are not

Spect-IPT values

	March 1995	March 1996	Age matched controls
Striatum left	3.1	3.1	
Striatum right	3.0	2.6	
Mean (SD)	3.0	2.9	7.3 (1.3)
Caudate left	4.5	3.9	
Caudate right	4.2	3.7	
Mean (SD)	4.4	3.8	8.6 (1.4)
Putamen left	1.8	2.7	
Putamen right	2.2	1.7	
Mean (SD)	2.0	2.2	6.5 (1.4)

SPECT was performed 90 to 120 minutes after injection of 150 MBq IPT. After reconstruction by filtered backprojection transverse slices corrected for attenuation were realigned parallel to the AC-PC line. For semiquantitative evaluation of specific [123 I]IPT binding ratios between striatum, caudate, putamen, and background regions were calculated. Specific uptake ratios were defined as mean counts per pixel in the respective region of interest minus mean counts in a background region divided by the mean counts in the background region. The uptake ratios obtained in the patient studies and in age matched controls are listed in the table. The increased IPT binding in the left putamen in the 1996 study is most probably due to an artefact caused by a spot of high background activity accidentally located around the putamenal region of interest in this investigation.

Compared with a representative control (A) the SPECT study of the patient showed profoundly reduced striatal IPT uptake (B). The decrease of specific IPT uptake was more pronounced in the putamen than in the caudate.

aware of a case reporting a levodopa responsive parkinsonian syndrome with a tumour in the supplementary motor area. Certainly, the criteria "responsiveness to dopaminergic drugs" may help to differentiate tumour induced parkinsonism from the concomitant idiopathic disease.^{4,5} Interestingly, Dick and coworkers reported on a patient with an infarct in the supplementary motor area, who had bradykinesia, but not from rigidity, which may suggest that the symptom "rigidity" may be suitable to differentiate tumour-induced versus concomitant parkinsonian syndrome.¹⁰ Furthermore, in rare cases, in which patients with parkinsonism with CNS tumours respond favourably to dopaminergic medication (to our knowledge only one case report¹¹), modern imaging techniques with specific ligands to presynaptic terminals of the nigrostriatal pathway further aid in differentiating between a tumour induced syndrome and comorbidity.

ANDREAS KUPSCH
OLIVER POGARELL
JOHANNES SCHWARZ
ANDREAS STRAUPE
WOLFGANG H OERTEL

Department of Neurology,
Ludwig-Maximilians-Universität, Klinikum
Großhadern

KLAUS TATSCH
RAINER LINKE

Department of Nuclear Medicine, Marchioninistrasse
15, 81377 München, Germany

OLIVER POGARELL
WOLFGANG H OERTEL

Department of Neurology, Philipps-Universität
Marburg, Rudolf-Bultmann-Strasse 8, 35033
Marburg, Germany

ANDREAS KUPSCH

Department of Neurology, Humboldt-University Berlin,
Charité-Virchow, Augustenburger Platz 1, 13359
Berlin, Germany

Correspondence to: Dr W H Oertel, Department of
Neurology, Philipps-Universität Marburg, Rudolf-
Bultmann-Strasse 8, 35033 Marburg, Germany.

- 1 Straube A, Sigel K. Parkinsonian syndrome caused by a tumour of the left supplementary motor area. *J Neurol Neurosurg Psychiatry* 1988;51:730-1.
- 2 Alexander GE, Crutcher MD, DeLong MR. Basal ganglia-thalamocortical circuits: parallel substrates for motor, oculomotor, "prefrontal" and "limbic" functions. In: Uylings HBM, van Eden CG, de Bruin JPC, Corner MA, Feenstra MGP, eds. *Prog Brain Res* 1990;85:115-46.
- 3 Myagi Y, Morioka T, Otsuka M, Fukui M. Striatal glucose and (18)fluorodopa uptake in a patient with tumor-induced hemiparkinsonism. *Neurosurgery* 1993;32:838-41.
- 4 Gasser T, Schwarz J, Arnold G, Trenkwalder C, Oertel WH. Apomorphine test in patients with previously untreated Parkinson's disease. *Arch Neurol* 1992;49:1131-4.
- 5 Hughes AJ, Lees AJ, Stern GM. Challenge tests to predict the dopaminergic response in untreated Parkinson's disease. *Neurology* 1992;41:1723-5.
- 6 Kung M-P, Essman WD, Frederick D, Meegalla S, Goodman M, Mu M, et al. IPT: a novel iodinated ligand for the CNS dopamine transporter. *Synapse* 1995;20:316-24.
- 7 Mozley PD, Kim HJ, Gur RC, Tatsch K, Muenz LR, Mc Elgin WT, et al. Iodine-123 IPT SPECT imaging of CNS dopamine transporters: non-linear effects of normal aging on striatal uptake values. *J Nucl Med* 1996;37:1965-70.
- 8 Tatsch K, Schwarz J, Mozley PD, Linke R, Pogarell O, Oertel WH, et al. Relationship between clinical features of Parkinson's disease and presynaptic dopamine transporter binding assessed with [I-123]IPT and single-photon-emission tomography. *Eur J Nucl Med* 1997;24:415-21.
- 9 Schwarz J, Tatsch K, Arnold G, Gasser T, Trenkwalder C, Kirsch CM, Oertel WH. 123I-iodobenzamide-SPECT predicts dopaminergic responsiveness in patients with de novo parkinsonism. *Neurology* 1992;42:556-61.

- 10 Dick JP, Benecke R, Rothwell JC, Day BL, Marsden CD. Simple and complex movements in a patient with infarction of the right supplementary motor area. *Mov Disord* 1986;1:255-66.
- 11 Lhermitte F, Agid Y, Serdaru M, Guimares J. Syndrome parkinsonien, tumeur frontale et L-DOPA. *Rev Neurol (Paris)* 1984;140:138-9.

Hereditary neuromyotonia: a mouse model associated with deficiency or increased gene dosage of the PMP22 gene

Neuromyotonia is characterised by increased muscle stiffness caused by hyperactivity of motor units. Clinical characteristics are increased stiffness and cramping on intended muscle contraction, slowed relaxation, and myokymia.^{1,2} Neuromyotonia is a heterogeneous condition.³ In the acquired form, autoimmunity against peripheral nerves seems to play a pathogenic part. Neuromyotonia may occasionally follow immune mediated polyneuropathy.⁴ Antibodies against voltage gated potassium channels have been detected in some cases.⁵ In the hereditary forms, neuromyotonia may occur in isolation⁶⁻⁸ or in association with hereditary neuropathies.^{3,9,10} We here report on neuromyotonia developing in aged mice homozygously deficient for or carrying an increased gene dosage of PMP22. Null mutants display a severe dysmyelinating neuropathy characterised by tomacula formation and subsequent myelin degeneration resembling human hereditary neuropathy with pressure palsies (HNPP).¹¹ Nerve conduction studies show a profound slowing¹¹ and repetitive compound muscle action potentials (CMAPs). Despite these profound

peripheral nerve abnormalities, mice did not show overactivity up to 12 months. Between the ages of 12 and 14 months, we found progressive overactivity with tonic stretching of the hind limbs and increased cramping of the small foot muscles augmented on voluntary contraction, and generalised myokymia including the whisker muscles in all three PMP22 deficient mice examined. These signs persisted during general anaesthesia. On needle EMG, we found serial high frequency bursts of motor units in the gastrocnemius, quadriceps, and small foot muscles (figure, A). Some of the discharges were doublets, triplets, or multiplets (figure, A and B). In addition, we found continuous myokymic muscle fibre activity (figure, B). Sciatic nerve transection at the sciatic notch did not abolish the overactivity indicating a peripheral nerve generator. In five 14-18 month old mice heterozygously deficient for the expression of PMP22 and with the typical electrophysiological signs of a dysmyelinating neuropathy, we found neuromyotonia only in one out of three animals at a single site in the gastrocnemius muscle. We also found severe neuromyotonia in a transgenic mouse model with an increased PMP22 gene dosage at one year of age (data not shown). These overexpressing mice have a severe hypomyelinating neuropathy in which Schwann cell development is arrested in a hyperproliferative and premyelination-like state.¹² This suggests that the occurrence of neuromyotonia is linked to the PMP22 gene dosage. We did not find abnormal nerve conduction, repetitive CMAP or neuromyotonia in four 13 month old control mice with normal PMP22 expression.

EMG recordings from a 14 month old mouse homozygously deficient for the expression of the peripheral myelin protein PMP22. The mouse was anaesthetised with Hypnorm (Janssen, Beerse, Belgium) and investigated using a Tönnies electromyograph with a concentric needle electrode (Medelec DFC25, 0.3 mm diameter, recording area 0.019 mm²). (A) Multiplet discharges in the small foot muscles. (B) Representative recording of myokymic discharges and doublets from another mouse (PMP22 -/-). There was abundant myokymic motor activity, but no voluntary limb movements during these recordings.